

Accepted refereed manuscript of:

Fathi M, Dickson C, Dickson M, Leschen W, Baily J, Muir K, Ulrich K & Weidmann M (2017) Identification of Tilapia Lake Virus in Egypt in Nile tilapia affected by ‘summer mortality’ syndrome, *Aquaculture*, 473, pp. 430-432.

DOI: [10.1016/j.aquaculture.2017.03.014](https://doi.org/10.1016/j.aquaculture.2017.03.014)

© 2017, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Highlights

- Summer mortalities in tilapia have occurred in Egypt for 3-4 years
- Epidemiological study finds 37% of tilapia farms affected
- TiLV PCR positive tilapia found on three of the seven farms with summer mortality syndrome
- Partial sequence (255bp) shows 93% homology to published TiLV segment 3

Identification of Tilapia Lake Virus in Egypt in Nile tilapia affected by ‘summer mortality’ syndrome

Mohamed Fathi^a, Cathryn Dickson^a, Malcolm Dickson^a, William Leschen^b, Johanna Baily^b, Fiona Muir^b, Kristina Ulrich^b, and Manfred Weidmann^b

^a WorldFish, Abbassa Research Center, Sharkia, Egypt

^b Institute of Aquaculture, University of Stirling, Stirling, Scotland, UK

Corresponding author

Manfred Weidmann
University of Stirling
Institute of Aquaculture
Stirling FK9 4LA
Scotland, United Kingdom

tel: +441786467873

e-mail: m.w.weidmann@stir.ac.uk

Running Title: Tilapia lake virus in Egypt

Keywords: Tilapia lake virus, Tilapia, Mullet, Egypt

Word count text: 994 words

1 **Abstract**

2

3 Egyptian fish farms have faced unexplained mortality of tilapia during the summer

4 months in recent years. Epidemiological surveys indicated that 37% of fish farms

5 were affected in 2015 with an average mortality rate of 9.2% and a potential

6 economic impact of around US\$ 100 million. Despite a number of researchers and

7 organizations investigating potential causes results so far have been inconclusive.

8 Meanwhile recent reports emerged of the presence of a new orthomyxovirus, Tilapia

9 Lake Virus (TiLV) in Israel, which shares a border and migrating avifauna with Egypt.

10 Tissue samples from seven farms affected by 'summer mortality' were tested at the

11 University of Stirling for TiLV. Samples from three of seven farms tested positive

12 using PCR; the first time that TiLV has been identified in Egypt. Sequence analysis

13 yielded a TiLV sequence with 93% homology to the published TiLV sequence

14 described from Israel. More research is required to determine if TiLV is linked to

15 'summer mortality'.

16

17 **Introduction**

18 The Egyptian aquaculture sector is the largest producer of farmed fish in Africa (1.14
19 million tonnes in 2014) and the third largest global producer of farmed tilapia after
20 China and Indonesia (FAO, 2016; Fitzsimmons, 2016). Until recently, there were no
21 major disease concerns but over the past three to four years farmers have faced
22 unexplained, significant mortalities of medium (>100g) and large sized pond reared
23 Nile tilapia (*Oreochromis niloticus*) during the summer months (June-October) when
24 water temperatures typically rise to over 25°C. These disease outbreaks, termed
25 ‘summer mortalities’, have been investigated by a number of research teams in an
26 attempt to identify causative agents but with inconclusive results. Another study
27 linked the opportunist pathogen *Aeromonas veronii* biovar *sobria* to disease outbreaks
28 of pond reared tilapia in Egyptian fish farms but this has not been confirmed as the
29 primary causative agent (Eissa, et al., 2015).

30 Meanwhile a new orthomyxovirus, termed Tilapia Lake Virus (TiLV) was isolated in
31 Israel in 2014 after an investigation into tilapia mortalities in wild and farmed stocks
32 and has also been associated with mortalities of farmed tilapia in Ecuador and
33 Columbia. The full TiLV genome was described and improvements to isolation and
34 detection methods reported. (Bacharach, et al., 2016; Eyngor, et al., 2014; Kembou
35 Tsofack, et al., 2016).

36 In 2015 WorldFish conducted an epidemiological study of ‘summer mortalities’ on 68
37 fish farms in the three most important Egyptian aquaculture governorates; Kafr El
38 Sheikh, Behera and Sharkia. The aim of this study was to identify the effect of various
39 production factors on the incidence of disease outbreaks in Egyptian pond based
40 aquaculture. Also, as Egypt shares a border with Israel and is on an important bird
41 migration route passing through Israel it seemed likely that TiLV could also be

42 present in Egypt and might be linked to ‘summer mortalities’. Accordingly, in
43 summer 2016 tilapia from Egyptian farms affected by ‘summer mortalities’ were
44 sampled and analysed for TiLV.

45

46 **Materials and Methods**

47 **Epidemiological Study**

48 The study was conducted on 68 randomly selected fish farms from the WorldFish
49 database in the governorates of Kafr el Sheikh (32 farms), Behera (15 farms), and
50 Sharkia (21) farms. These areas are responsible for around 85% of Egyptian
51 aquaculture production (GAFRD, 2016). A structured questionnaire and monthly field
52 visits by WorldFish staff recorded fish mortalities over the growing season. Data were
53 also collected on; farm size, stocking density, production systems, and economic
54 impact of losses due to ‘summer mortalities’.

55

56 **Sampling for TiLV**

57 After consultation with the Egyptian authorities, liver, brain, kidney and spleen tissues
58 from moribund fish from 7 ‘summer mortality’ affected farms (3 fish per farm) were
59 sent to the Institute of Aquaculture at Stirling University to test for TiLV. Tissue
60 samples from healthy fish from 4 non-affected farms were also sent as controls.
61 Samples for PCR assay were transported in RNA preservation reagent while viral
62 transport medium and sterile tubes, kept chilled, were used to transport samples for
63 virus isolation.

64 Tissues from all sampled fish were fixed in neutral buffered formalin for histological
65 analysis. Tissue homogenates of these samples were also inoculated onto BF-2, EPC

66 and CHSE-214 cell lines and RNA was extracted from tissue samples using TRI-
67 reagent and tested by RT-PCR for segment 3 as described (Eyngor et al, 2014) .

68

69 **Results**

70 **Epidemiological study**

71 Out of 68 randomly selected fish farms, 37% (25 farms) were affected by ‘summer
72 mortalities’ with an average mortality rate of 9.2% (range 5-15%). Incidence rates
73 were significantly higher in farms with higher stocking densities and in large farms
74 compared to smaller farms (Tab. 1). Many farms grow mullet (*Mugil cephalus* L and
75 *Liza ramada* Risso, 1827) in co-cultivation with tilapia (85% tilapia: 15% mullet)
76 because of the higher prices obtained for these species. Notably only 3 / 38 farms
77 practicing monoculture of tilapia were affected whereas 22 / 30 farms practicing
78 tilapia / mullet polyculture were affected. Further studies are required to investigate
79 the possible role of mullet (which appear to be resistant to summer mortalities) in
80 disease transmission (Tab 1). The overall impact of summer mortalities in 2015 was
81 estimated at 98,000 tonnes of lost production (US\$ 100 million).

82

83 **Analysis of clinical samples**

84 the following pooled tissue samples (n=3, brain = B, kidney = K, liver =L, spleen = S)
85 of three farms tested positive: L and S samples from farm 1, S from farm 2 and B, K,
86 L, S from farm 3 - a first time detection of TiLV in this laboratory and in tilapia from
87 Egypt. Histologic lesions in tissues of the PCR positive fish included multiple foci of
88 gliosis, encephalitis and mild perivascular cuffing by lymphocytes in the brain; mild
89 chronic meningitis; mild perivascular and multifocal, chronic hepatitis and moderate,
90 multifocal interstitial haemorrhage in the kidney (Fig. 1.). The epidermis was

91 moderately disrupted with superficial haemorrhage, inflammation, and oedema. In
92 comparison, unaffected, PCR negative fish from uninfected negative control farms
93 showed very mild perivascular hepatitis and occasional encysted trematodes within
94 the gut wall. Moderate diffuse congestion/haemorrhage of the spleen was common to
95 both PCR positive and PCR negative fish.

96 Cycle sequencing of the PCR amplicates yielded a TiLV sequence with 93%
97 homology to the published TiLV sequence (Fig. 2.). We did not isolate the virus.

98

99 **Discussion**

100 Clearly ‘summer mortalities’ of tilapia are having a significant impact on Egyptian
101 aquaculture. Anecdotal evidence suggests that the incidence is increasing year by year
102 although this survey is the first to provide credible figures on incidence and mortality
103 rates. In the current study, larger farms, higher stocking densities and co-cultivation of
104 tilapia with mullet appear to be contributory factors in the observed incidence of
105 ‘summer mortalities’ in Egyptian fish farms but further studies are required.

106 Now that a virus with high homology to TiLV has been identified in Egypt in tilapia
107 affected by ‘summer mortality’, further research including epidemiological and
108 experimental infection studies are needed to determine if it is the primary cause. If so,
109 rapid action can be taken to control the spread of the disease including immediate
110 improvements to aquaculture biosecurity practices and in the longer term, vaccines
111 and breeding TiLV resilient strains of tilapia.

112

113 **Acknowledgment**

114 This work was supported by the Swiss Agency for Development and Cooperation, the
115 Livestock and Fish CGIAR Research Program and the Marine Alliance for Science

116 and Technology in Scotland

117

Table 1

Farm management factors affecting incidence of summer mortalities

Management factor	Affected farms	Un-affected farms	P-value
Mean stocking density (1000 fish.ha ⁻¹)	51.45	33.98	0.050
Average farm size (ha)	13.44	7.00	0.043
Farms practicing tilapia monoculture (no)	3	35	0.0001
Farms practicing tilapia-mullet polyculture (no)	22	8	0.015

Stocking densities and average farm sizes were higher in ‘summer mortality’ affected farms than in un-affected farms while farms practicing polyculture were more likely to be affected than farms practicing monoculture of tilapia

Figures

Fig. 1. Histological findings in PCR positive tilapia (HE staining, × 200). Upper panel: Focally extensive chronic encephalitis – brain stem. Middle panel: Chronic perivascular hepatitis centred around vasculitis. Lower panel: Renal interstitial haemorrhage. All: Haematoxylin and eosin stain, magnification x200.







Fig. 2. Weighted dendrogram of a 255 nucleotide PCR fragment of TiV segment 3, aligned with ClustalW, modelled with RaxML. ISAV = Infectious salmon anaemia virus, FLUAV = Influenza A virus.



■ 0.1

References

- Bacharach, E., Mishra, N., Briese, T., Zody, M.C., Tsofack, J.E.K., Zamostiano, R., Berkowitz, A., Ng, J., Nitido, A., Corvelo, A., Toussaint, N.C., Nielsen, S.C.A., Hornig, M., Del Pozo, J., Bloom, T., Ferguson, H., Eldar, A., Lipkin, W.I., 2016. Characterization of a Novel Orthomyxo-like Virus Causing Mass Die-Offs of Tilapia. *Mbio*. 7.
- Eissa, I.A.M., El-lamei, M., Sherif, M., Desuky, M., Zaki, M.S., Bakry, M., 2015. *Aeromonas veronii* biovar *sobria* a Causative Agent of Mass Mortalities in Cultured Nile Tilapia in El-Sharkia governorate Egypt. *Life Science Journal* 12, 90-97.
- Eyngor, M., Zamostiano, R., Tsofack, J.E.K., Berkowitz, A., Bercovier, H., Tinman, S., Lev, M., Hurvitz, A., Galeotti, M., Bacharach, E., Eldar, A., 2014. Identification of a Novel RNA Virus Lethal to Tilapia. *J Clin Microbiol*. 52, 4137-4146.
- FAO. 2016. The State of World Fisheries and Aquaculture 2016. Contributing to food security and nutrition for all. Rome. 200 pp.
- Fitzsimmons, K., 2016. Supply and demand in global tilapia markets, 2015. Aquaculture 2016, Las Vegas, Nevada, 23-26 Feb 2016. Accessed at https://www.was.org/documents/MeetingPresentations/AQ2016/AQ2016_0005.pdf
- GAFRD, 2016. Fish Statistics Yearbook 2014. Ministry of Agriculture and Land Reclamation, Egypt. General Authority for Fishery Resources Development. Ministry of Agriculture and Land Reclamation, Egypt.
- Tsofack, J.E.K., Zamostiano, R., Wattedb, S., Berkowitzb, A., Rosenbluth, E., Mishra, N., Briese, T., Lipkin W.I., Kabuusud, R.M., Ferguson H., del Pozoe, J., Eldarb, A., Bacharach, E. 2016. Detection of Tilapia Lake Virus (TiLV) in Clinical Samples by Culturing and Nested RT-PCR. *J Clin Microbiol*. 55, 759-767.