

Baseline Studies of Biodiversity:

The Fish Resources of Western Indonesia

Edited by

D. Pauly and P. Martosubroto



Directorate General
of Fisheries, Indonesia



German Agency
for Technical Cooperation



International Center for Living Aquatic
Resources Management





**Baseline Studies of Biodiversity:
The Fish Resources of Western Indonesia**

ENTERED IN MAGA

Edited by

**D. Pauly
and
P. Martosubroto**

1996

DIRECTORATE GENERAL OF FISHERIES
Jakarta, Indonesia

GERMAN AGENCY FOR TECHNICAL COOPERATION
Eschborn, Germany

INTERNATIONAL CENTER FOR LIVING AQUATIC RESOURCES MANAGEMENT
Manila, Philippines

NOV 08 1996

**Baseline Studies of Biodiversity:
The Fish Resources of Western Indonesia**

Edited by

D. PAULY
and
P. MARTOSUBROTO

1996

Printed in Manila, Philippines

Published by the International Center for Living Aquatic Resources Management,
MCPO Box 2631, 0718 Makati City, Philippines with financial assistance
from the German Agency for Technical Cooperation (GTZ)

Pauly, D. and P. Martosubroto, Editors. 1996. Baseline studies of biodiversity:
the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 321 p.

Cover design by Robbie Cada, Alan Esquillon and D. Pauly

ISSN 0115-4389
ISBN 971-8709-48-7

ICLARM Contribution No. 1309

Contents^a

Preface • <i>D. Pauly</i> and <i>P. Martosubroto</i>	vii
Forewords	
DGF Foreword • Rear Admiral F.X. Murdjijo	ix
GTZ Foreword • Dr. Martin Bilio	x
ICLARM Foreword • Dr. Meryl J. Williams	xii
 Physical historical and conceptual background of the Western Indonesian fish resources surveys (1974-1981)	
Biodiversity and the retrospective analysis of demersal trawl surveys: a programmatic approach • <i>D. Pauly</i>	1
Oceanography of the Indonesian Archipelago and adjacent areas • <i>G.D. Sharp</i>	7
Variability of sea surface features in the Western Indonesian Archipelago: inferences from the COADS dataset • <i>C. Roy</i>	15
The marine fisheries of the Western Archipelago: towards an economic history, 1850 to the 1960s • <i>J. Butcher</i>	24
 Fish communities surveyed by <i>R/V Mutiara 4</i>	
The mid-1970s demersal resources in the Indonesian side of the Malacca Strait • <i>P. Martosubroto</i> , <i>T. Sujastani</i> and <i>D. Pauly</i>	40
The <i>Mutiara 4</i> surveys in the Java and southern South China Seas, November 1974 to July 1976 • <i>D. Pauly</i> , <i>P. Martosubroto</i> and <i>J. Saeger</i>	47
Demersal assemblages of the Java Sea: a study based on the trawl surveys of the <i>R/V Mutiara 4</i> • <i>G. Bianchi</i> , <i>M. Badrudin</i> and <i>S. Budihardjo</i>	55
Structure and dynamics of the demersal resources of the Java Sea, 1975-1979 • <i>P. Martosubroto</i>	62
 Fish communities surveyed by <i>R/V Jurong</i>	
Narrative and major results of the Indonesian-German Module (II) of the JETINDOFISH Project, August 1979 to July 1981 • <i>U. Lohmeyer</i>	77
Marine bottomfish communities from the Indian Ocean coast of Bali to mid-Sumatra • <i>J. McManus</i>	91
 Fish communities surveyed by <i>R/V Lemuru</i>, <i>R/V Dr. Fridtjof Nansen</i> and <i>R/V Bawal Putih 2</i>	
Results of surveys for pelagic resources in Indonesian waters with the <i>R/V Lemuru</i> , December 1972 to May 1976 • <i>S.C. Venema</i>	102
Box 3. Development and fleet structure of the Java Sea pelagic fisheries (1980-1992) • <i>J. Widodo</i> , <i>M. Potier</i> and <i>B. Sadhotomo</i>	120
Box 4. Catches, effort and catch per effort of the Java Sea pelagic fishery (1979-1992) • <i>J. Widodo</i> , <i>M. Potier</i> and <i>B. Sadhotomo</i>	121

^aIncluding authored "boxes" in formal chapters.

Demersal fish assemblages of trawlable grounds off northwest Sumatra • <i>G. Bianchi</i>	123
Narrative and major results of the Indonesian Module (I) of the JETINDOFISH Project, November 1980 to October 1981 • <i>P. Martosubroto</i>	131
Box 1. Small-scale Fisheries Development Project in Lombok, Nusa Tenggara Barat, Indonesia • <i>R. Hermes</i>	133

Regional syntheses

Fishery biology of 40 trawl-caught teleosts of Western Indonesia • <i>D. Pauly</i> , <i>A. Cabanban</i> and <i>F.S.B. Torres, Jr.</i>	135
Box 1. Estimating the parameters of length-weight relationships from length-frequency samples and their weights • <i>D. Pauly</i> and <i>F.C. Gayanilo, Jr.</i>	136
Box 2. The Bali Strait <i>Iemuru</i> fishery • <i>A. Ghofar</i> and <i>C.P. Mathews</i>	146
Box 3. Reasons for studying the Leiognathidae • <i>D. Pauly</i>	177
Box 4. A case study of <i>Nemipterus thosaporni</i> a.k.a. <i>N. marginatus</i> • <i>D. Pauly</i>	199
Checklist of marine fishes of Indonesia, compiled from published literature • <i>R. Froese</i> , <i>S.M. Luna</i> and <i>E.C. Capuli</i>	217
Box 1. Nomenclatural changes in <i>Trawled fishes of Southern Indonesia</i> and <i>Northwestern Australia</i> • <i>R. Froese</i>	218
Box 2. A computerized procedure for identifying misspellings and synonyms in checklists of fishes • <i>R. Froese</i>	219
Using NAN-SIS and FiSAT to create a trawl survey database for Western Indonesia • <i>F.S.B. Torres, Jr.</i> , <i>A. Cabanban</i> , <i>S.L. Bienvenida</i> , <i>J.W. McManus</i> , <i>M. Prein</i> and <i>D. Pauly</i>	276
Box 1. JETINDOFISH data processing problem • <i>B.G. Thompson</i>	277
Box 2. Importing files to NAN-SIS • <i>T. Strømme</i>	280

Appendix I.

FishBase references cited in Pauly et al. (p. 135-216) and Froese et al. (p. 217-275), by number	284
---	-----

Appendix II.

Author/Name Index.....	297
Geographic Index	303
Species Index.....	306

Preface

This book is long overdue: the trawl and acoustic surveys documented here have been conducted some 20 years ago, and several of the fish communities described in the various chapters of this book have been, in the meantime, fished strongly enough to be barely recognizable.

However, no major surveys have been conducted in Western Indonesia since the period covered here (1975-1981). During that period, a convergence of interest had led to a flurry of bilateral and international fisheries development projects along the Indian Ocean coast of Indonesia, in the Java Sea and adjacent waters, funded by German (GTZ)^a, Australian (AIDAB)^b and Norwegian (NORAD)^c aid agencies, and by FAO^d.

Twenty years ago (May 1975) is also when the two editors of this book first met; we had both just acquired our MS degrees, and were eager to apply what we had learnt. We became friends, despite our vastly different cultural backgrounds, perhaps aided therein by a car accident that occurred at Pemanukan along the coast of West Java, and which could easily have killed us both.

Although soon separated, and working in very different institutions, we both felt that the surveys we had jointly worked on, and the surveys done shortly thereafter should have been better documented than through internal reports and theses.

One of us used the opportunity provided by a book review to criticize this state of affairs^e; the other pushed from within the Directorate General of Fisheries (DGF, Jakarta). The result of our joint effort was an official request sent by DGF to GTZ to support the production of a book in which the surveys would be documented and analyzed, and which would complement the excellent volume published jointly by GTZ, DGF and AIDAB documenting the taxonomy of the demersal fishes of Western Indonesia^f.

GTZ agreed, and ICLARM was invited to submit a proposal, accepted in 1991, for a project that would lead to the publication of a book that would not only complement the previous volume on the *Trawled fishes of Southern Indonesia and Western Australia*, but also be made to resemble it. This explains the choice of the format and fonts used here, which differ from those of other items in ICLARM's Studies and Reviews Series.

The book is thus the result of a long chain of events. However, the delay this implied was, we believe, turned from a liability into an asset. Thus, we present not only the results of old surveys, but a new way of interpreting them, in the biodiversity context that has become a major issue for the outgoing 20th century.

The reconceptualization of old surveys into baseline studies of biodiversity is not followed up here in all its ramifications, the papers of D. Pauly, G. Bianchi, J. McManus and R. Froese, S. Luna and E. Capuli providing only pointers. However, the survey results made available through this book (and as computer datafiles) should ensure that there is now, indeed, a sound baseline for fish biodiversity studies in Western Indonesian waters. Thus, we are conscious of having crossed a bridge, and we are confident that this book will help its reader realize, for other areas as well, the importance of "old" trawl surveys as source of biodiversity baseline data.

All that is now left for us to do is to thank all involved directly or indirectly in the creation of this book and especially the authors, not only because without them, there would be no book, but also for their agreeing to our rather specific editorial guidelines, and for delivering their papers in a timely fashion - well, almost!

We would like to thank Ms. Eny A. Buchary, Fisheries Centre, UBC, for checking the Indonesian translations.

Also we would like to thank our former colleagues and ship crew for their collaboration during the surveys in which we participated and our present colleagues at ICLARM and FAO for their encouragement of our work on this book.

Finally, we would like to thank GTZ for its support, notably Dr. M. Bilio, and the DGF, especially Mr. Soewito, former Director of Resources Management, for enabling us to make use of the data collected by *R/V Jurong*.

D. Pauly
Manila/Vancouver

P. Martosubroto
Rome/Jakarta

^a Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ), GmbH, Eschborn, Germany.

^b Australian International Development Assistance Bureau, Canberra, Australia (now AUSAID).

^c Norwegian Agency for International Development, Oslo, Norway.

^d Food and Agriculture Organization of the United Nations, Rome, Italy.

^e Pauly, D. 1986. On identifying fish species rather than assessing fish stocks: a review of two bodies on the taxonomy of the neritic fishes of the Western Indian Ocean. *Naga*, ICLARM Q. 9(3):21.

^f Gloerfelt-Tarp, T. and P. Kailola. 1984. *Trawled fishes of Southern Indonesia and Northwestern Australia*. AIDAB/DGF/GTZ. 406 p.

Kata Pendahuluan

Buku ini terbit sangat terlambat: survei-survei trawl dan akustik yang didokumentasikan disini dilaksanakan kurang lebih 20 tahun yang lalu, sementara itu beberapa komunitas ikan yang digambarkan dalam beberapa bab di dalam buku ini, telah mengalami tekanan penangkapan yang sangat tinggi sehingga hampir-hampir mereka tidak terlihat lagi.

Namun demikian, tidak ada lagi upaya survei yang besar yang dilaksanakan di perairan Indonesia bagian barat sejak periode survei diatas (1975-1981). Selama periode tersebut, berbagai interest telah mendorong terbentuknya proyek-proyek perikanan bilateral maupun internasional di sepanjang Samudra Hindia, di laut Jawa dan sekitarnya, dibiayai oleh lembaga bantuan dari Jerman (GTZ), Australia (AIDAB) dan Norwegia (NORAD) serta FAO.

Dua puluh tahun yang lalu (Mei 1975) kedua editor buku ini pertama kali bertemu; kami berdua baru saja mengantongi gelar MS, dan sangat berkeinginan mengaplikasikan ilmu yang baru kami peroleh. Kami menjadi teman akrab, walaupun mempunyai latar belakang budaya yang berbeda, barangkali juga dipererat dengan musibah kecelakaan mobil di Pemanukan di pantai Utara Jawa Barat yang hampir-hampir merenggut nyawa kami berdua.

Walaupun kemudian kami berpisah, dan bekerja di institusi yang sangat berlainan, kami merasa bahwa survei-survei dimana kami bekerja bersama, dan survei-survei yang dilaksanakan kemudian akan lebih baik dipublikasikan daripada hanya merupakan laporan survei dan thesis saja.

Salah satu diantara kami memanfaatkan kesempatan dalam suatu resensi buku dengan menyampaikan kritikan akan masalah seperti ini; sedangkan yang lain berusaha meyakinkan dan menyamakan pandangan di Direktorat Jendral Perikanan (DGF, Jakarta). Sebagai hasilnya suatu permintaan resmi dikirim ke GTZ untuk membantu membiayai pembuatan buku ini dimana hasil-hasil survei akan didokumentasikan dan dianalisis, sehingga merupakan pasangan dari buku taksonomi ikan-ikan demersal Indonesia bagian barat hasil publikasi bersama antara GTZ, DGF dan AIDAB.

GTZ menyetujui usulan diatas dan ICLARM diminta untuk membuat suatu usulan proyek yang mana diterima pada tahun 1991, proyek tersebut bermuara pada penyusunan suatu buku yang tidak hanya merupakan pasangan buku "Trawled fishes of Southern Indonesia and Western Australia", tetapi juga dirancang agar menyerupainya. Oleh karena itu, bentuk dan format buku ini berbeda dengan bentuk dan format dari buku-buku edisi ICLARM's Studies and Reviews Series.

Dengan kata lain, penyusunan buku ini telah melalui proses yang panjang. Namun demikian, kami percaya bahwa keterlambatan keluarnya buku ini justru memberikan suatu hikmah. Yaitu, kami tidak hanya menyajikan hasil-hasil survei, tetapi juga suatu metode baru dalam menginterpretasikan hasil-hasil tersebut dari segi keanekaragaman hayati, dimana hal terakhir ini menjadi topik besar dalam era meninggalkan abad ke 20.

Konsepsualisasi kembali terhadap hasil survei untuk menjadi studi dasar dari keanekaragaman hayati tidak dibahas secara keseluruhan, tulisan-tulisan D. Pauly, G. Bianchi, J. McManus serta tulisan R. Froese, S. Luna, dan E. Capuli hanya memberikan beberapa petunjuk pokok saja. Tetapi, hasil survei yang disajikan dalam buku ini (dan dalam bentuk file computer) paling tidak menjamin bahwa sekarang telah tersedia suatu data-dasar untuk studi keanekaragaman hayati bagi perairan Indonesia bagian barat. Dengan sadar kami telah meniti suatu jembatan, dan kami yakin bahwa buku ini dapat membantu para pembaca dalam memahami, tentunya termasuk untuk daerah lain juga, akan pentingnya survei trawl yang sudah lewat sebagai sumber data dasar keanekaragaman hayati.

Tidak lupa kami menyampaikan banyak terima kasih kepada semua pihak yang telah terlibat baik langsung atau tidak langsung, dalam penyusunan buku ini; khususnya kepada para penulis, karena tanpa mereka tidak akan tersusun buku ini. Terima kasih atas keterbukaan mereka mengikuti acuan edisi yang agak spesifik, dan akan kerjasama mereka memenuhi jadwal waktu - boleh dikata hampir tepat waktu.

Kami juga mengucapkan terima kasih kepada Eny A. Buchary, Fisheries Centre, UBC, atas bantuannya memeriksa terjemahan Indonesia dari buku ini.

Kami juga berterimakasih kepada semua teman-teman lama termasuk para awak kapal atas kerjasama yang terjalin dan juga kepada teman-teman di ICLARM dan FAO atas dorongan terhadap penyusunan buku ini.

Akhirnya kami menyampaikan terima kasih kepada GTZ akan bantuannya, khususnya Dr. M. Billio, dan DGF, terutama Bapak Soewito, mantan Direktur Bina Sumber Hayati, yang memungkinkan kami memanfaatkan data yang dikumpulkan kapal penelitian Jurong.

DGF Foreword

As part of the implementation of the first five-year development plan in the fisheries sector, Indonesia received technical assistance from various governments (i.e., the Federal Republic of Germany, The Netherlands, Australia) and international organizations of the United Nations (the United Nations Development Programme and the Food and Agriculture Organization) (UNDP and FAO) in the area of resource surveys conducted in many parts of our archipelagic waters. The result of the surveys has enriched our knowledge of various aspects of marine resources including their habitats, which form a useful resource base that directly or indirectly provides a contribution to the development of fisheries in the country.

The multispecies nature of our marine resources represents the complexity of our tropical marine ecology on which our fisheries are dependent. The wealth of the survey data that has been accumulated so far is a good base from which to generate more information provided that review analysis and scrutiny are undertaken. Dr. Daniel Pauly of the International Center for Living Aquatic Resources Management (ICLARM), who has been working in Indonesia during the GTZ-funded Demersal Fisheries Project in the Java Sea in the mid-1970s, and Dr. Purwito Martosubroto of FAO, Dr. Pauly's counterpart during that time and who later became Director of Resources Management at the Directorate General of Fisheries (DGF), have kindly made an effort to coordinate the present review work with many contributions from the international scientific community. This work is certainly a useful endowment to our knowledge of the resources in Western Indonesia.

This book, which represents an in-depth review of the resources and their environment, is not only useful for scientists but also for policymakers and managers - for them to understand the dynamics of fisheries resources upon which our development policy should be based.

Rear Admiral **F.X. Murdjijo**
Director General of Fisheries
Republic of Indonesia

Kata Pengantar DGF

Sebagai salah satu pelaksanaan rencana pembangunan lima tahun pertama di sektor perikanan, Indonesia memperoleh bantuan teknis dari berbagai negara (antara lain dari Pemerintah Jerman Barat, Belanda dan Australia) dan organisasi internasional yang bernaung dibawah PBB (seperti UNDP dan FAO) dalam bentuk survei di berbagai wilayah perairan kepulauan kita. Hasil survei ini menambah pengetahuan kita akan sumberdaya perikanan laut termasuk tempat hidupnya yang merupakan informasi yang bermanfaat dan langsung atau tidak langsung memberikan sumbangan terhadap pembangunan perikanan kita.

Sifat multispecies sumberdaya laut kita mencerminkan kompleksnya ekologi perairan tropis dimana perikanan kita sangat tergantung akan sumberdaya ini. Data yang telah terkumpul dari berbagai survei merupakan data dasar yang sangat bermanfaat apabila diadakan analisis yang mendalam. Dr. Daniel Pauly dari ICLARM (the International Center for Living Aquatic Resources Management) pernah bekerja di Indonesia dalam rangka proyek Perikanan Demersal Laut Jawa yang dibiayai oleh Pemerintah Jerman Barat (GTZ) pada pertengahan tahun 1970-an dan Dr. Purwito Martosubroto dari FAO yang merupakan rekan kerja Dr. Pauly pada saat itu dan kemudian pernah menjadi Direktur Bina Sumber Hayati di Direktorat Jendral Perikanan, keduanya dengan baik hati mengkoordinasikan upaya review ini, yang juga memperoleh berbagai sumbangan dari masyarakat ilmiah internasional, dan hasilnya merupakan khasanah pengetahuan yang sangat bermanfaat mengenai sumberdaya laut di perairan Indonesia bagian barat.

Buku ini yang merupakan review yang mendalam tentang sumberdaya dan lingkungan yang tidak hanya berguna bagi para peneliti tetapi juga para pembuat kebijakan dan manajer dalam rangka memahami tentang dinamika sumberdaya perikanan, yang mana yang terakhir ini merupakan informasi dasar dalam perumusan kebijaksanaan pembangunan kita.

F. X. Murdjijo
Direktur Jenderal Perikanan
Republik Indonesia

GTZ Foreword

It is with considerable satisfaction that the undersigned sees the present book published. It follows one of the major lines of collaboration between GTZ and ICLARM, i.e., to make sure that valuable data on fish resources collected at great expenditure are made available to the public in order to allow their maximum use by scientists and resource managers. Regarding the JETINDOFISH data, it was long felt that the publication of the beautiful book by Gloerfelt-Tarp and Kailola (*Trawled fishes of Southern Indonesia and Northwestern Australia*) in 1984 could not fully develop its potential without the publication of the whole biological and ecological information obtained from the trawling surveys.

Publishing such enormous amounts of data as were accumulated during the JETINDOFISH cruises was a task that could be shouldered only by a team that was familiar with the area and its fish fauna and who experienced the circumstances under which the cruises were conducted. It was also essential that such a team be backed by an institution with well-established infrastructure for publication purposes and experienced staff for the routine work. There was, therefore, every reason to wholeheartedly agree when Daniel Pauly, after many other useful proposals of this kind, approached me at GTZ with a request to support this new undertaking.

It took a while until all difficulties of administrative and other kinds were overcome; however, the team did not lose courage and continued their tedious work making and tying up loose ends in order to obtain a consistent picture of the resource situation. In doing so, they went well beyond initial expectations: not only were the data from the JETINDOFISH surveys considered but also a number of other sources of pertinent information were tapped, notably a survey of the *R/V Dr. Fridtjof Nansen*, and several other competent colleagues were asked to join the team. The publication of the present book is thus the fruit of a true international effort worth of the much commitment of an international institution such as ICLARM.

The region to which this publication refers is known, with regard to marine organisms, as the richest on Earth. In the last decade, the importance of biodiversity and its protection and conservation to the benefit of future generations has gained increased public attention. This is not simply based on simple cost-benefit considerations, but also on other values. The ensemble of organisms that surround us and of which a considerable number are essential for our survival, is not an endpoint of evolution, but only a stage of it. The capacity for further adaptation of this ensemble to the ever changing conditions of life on Earth depends on the genetic potential held by the organisms currently in existence. Every loss of species reduces this. However, protection of biodiversity can only be successful when we know what we are going to protect. An inventory of species must, therefore, form the basis of conservation and management of the resources. The attempt at reconceptualizing the elaboration of the survey data into a baseline study of biodiversity is thus a very timely effort.

Congratulations and thanks to the editors and the authors and to ICLARM for a remarkable achievement.

Dr. **Martin Bilio**

GTZ Senior Adviser (ret.) for Living
Aquatic Resources Utilization

Kata Pengantar GTZ

Saya merasa sangat berbahagia sekali dengan terbitnya buku ini yang merupakan kelanjutan kerjasama yang baik antara GTZ dan ICLARM, khususnya dalam rangka upaya memberikan jaminan akan data survei yang diperoleh dengan biaya yang cukup besar agar tersedia bagi khalayak ramai dan dapat dimanfaatkan oleh para peneliti serta pengelola sumberdaya perikanan pada umumnya. Mengenai data proyek JETINDOFISH, telah lama dirasakan bahwa terbitnya buku "Ikan-ikan yang tertangkap dengan trawl di perairan Indonesia bagian selatan dan Australia bagian barat laut" oleh Gloerfelt-Tarp dan Kailola yang terbit pada tahun 1984 tidak akan banyak memberikan manfaat tanpa terbitnya buku ini yang berisi informasi tentang berbagai segi biologi dan ekologi dari ikan-ikan yang tertangkap dalam survei tersebut.

Menerbitkan hasil analisis data yang banyak dikumpulkan dalam survei JETINDOFISH merupakan upaya yang tidak mudah, hal mana hanya dapat dilaksanakan oleh suatu tim yang memang mengetahui dan memahami daerah survei, sumberdaya dan lingkungannya. Jelas bahwa tim tersebut juga harus didukung oleh institusi yang lengkap dengan sarana publikasi serta staff yang berpengalaman. Oleh karena itu saya menyetujui usul Daniel Pauly beberapa tahun yang lalu agar GTZ dapat memberikan dukungan biaya dalam publikasi ini.

Upaya pembuatan buku ini memakan waktu cukup lama setelah masalah administrasi dapat diselesaikan, namun anggota tim tidak kehilangan semangat dan terus berusaha dengan gigih, menyempurnakan hal-hal yang masih kurang disana-sini dan akhirnya dapat merampungkannya. Hasilnya, tanpa disadari, lebih dari perkiraan awal, yaitu tidak hanya mencakup tulisan dengan sumber data JETINDOFISH saja melainkan juga dari sumber informasi lain yang terkait, seperti halnya dari survei kapal penelitian Dr. Fridtjof Nansen, dan dengan dukungan rekan-rekan yang kompeten. Publikasi buku ini merupakan hasil pencerminan kerjasama internasional yang sungguh-sungguh dari organisasi internasional seperti ICLARM.

Sebagaimana kita ketahui bahwa daerah survei meliputi Indonesia bagian barat yang merupakan daerah yang terkaya akan sumberdaya lautnya. Dalam dekade terakhir ini arti penting keanekaragaman hayati dan upaya perlindungan serta konservasi, demi generasi yang akan datang, telah banyak memperoleh dukungan masyarakat. Pandangan ini tidak hanya berdasar kepada pertimbangan untung rugi dari segi ekonomi semata tetapi juga bertumpu kepada nilai-nilai lainnya yang terkandung. Organisme di sekeliling kita sebagian besar diperlukan bagi kelangsungan hidup manusia. Kemampuan organisme tersebut untuk beradaptasi terhadap lingkungan yang terus berubah tergantung kepada potensi genetiknya. Dengan punahnya salah satu spesies berarti berkurangnya kekayaan genetika. Perlindungan terhadap keanekaragaman hayati hanya akan berhasil kalau kita memahami benar apa yang mau kita lindungi. Oleh karena itu, inventarisasi spesies merupakan dasar pokok konservasi dan pengelolaan sumberdaya. Upaya mengkonseptualisasikan elaborasi hasil survei ini sebagai dasar studi keanekaragaman hayati adalah merupakan upaya yang tepat sekali.

Selamat dan terimakasih kepada para editor dan para penulis, demikian halnya kepada ICLARM atas hasil yang tiada taranya ini.

Dr. Martin Bilio
Penasehat Senior Pemanfaatan
Sumberdaya Hayati Perairan GTZ

ICLARM Foreword

The International Center for Living Aquatic Resources Management (ICLARM), takes great pleasure in publishing this book on the fish resources of Western Indonesia.

We believe the book will be valuable because of the very geographic location it examines - containing perhaps the most biologically diverse and rich assemblages of marine life known. But it also will have relevance well beyond Indonesian waters. In tropical demersal fisheries, our present knowledge base is quite recent and quite meager compared to that for more temperate regions. For few tropical regions therefore, do we have such a comprehensive scientific analysis including oceanography, climate, ecosystem and the biology of the major fished species. An additional value in this book is the new ground it breaks in analyses of fish diversity.

In addition to the resource surveys documented in its various chapters, this book builds onto important precursors. One of these is the book entitled, "*Trawled fishes of Southern Indonesia and Northwestern Australia*" by Thomas Gloerfelt-Tarp and Patricia Kailola, published in 1984 by the Australian International Development Assistance Bureau (AIDAB - now AUSAID), Canberra, Australia, the Directorate General of Fisheries (DGF), Jakarta, Indonesia, and the Deutsche Gesellschaft für Technische Zusammenarbeit, GmbH (GTZ), Eschborn, Germany. These three organizations with assistance from the Food and Agriculture Organization of the United Nations (FAO), had conducted, from 1978 to 1981, a series of trawl surveys in Western Indonesia, and the book in question had superbly illustrated and documented the multitude of fish species caught in these surveys.

The present volume, deliberately conceived as a complement to its taxonomically-oriented predecessor, now presents detailed community and other analyses of these, and other surveys in the same region. Moreover, its second longest chapter shows how the list of the fish in Gloerfelt-Tarp and Kailola's book was complemented and updated using FishBase 96 (Froese, R. and D. Pauly, Editors. 1996 *FishBase 96: concept, design and data sources*. ICLARM, Manila, Philippines. 179 p.), the global CD-ROM encyclopedia of fishes, and thus kept "alive", despite now being out of print.

Last but not least, this book also has an antecedent the series of documents emanating from a successful collaboration between Indonesian Institutions and ICLARM, notably "*Indonesian marine fisheries*" (Bailey, C., A. Dwiponggo and F. Marahudin, 1987. ICLARM Stud. Rev. 10. 196 p.), and the atlas of the "*Growth, mortality and recruitment of commercially important fishes and penaeid shrimps in Indonesian waters*" (Dwiponggo, A., T. Hariati, S. Banon, M.L. Palomares and D. Pauly, 1996. ICLARM Tech. Rep. 17, 91 p.).

As well as these precursors, another stimulus to the book presented here was the realization by its senior editor of the usefulness of trawl surveys of unexploited grounds as baselines for biodiversity studies, for providing targets for attempt to rehabilitate depleted demersal stocks and for estimating the economic costs of not rehabilitating such stocks. Considerations of this sort are at the heart of several present ICLARM projects, and this book thus presents themes to which we shall frequently return. The work therefore represents new directions in tropical fisheries data analysis, which we propose to pursue through collaborative research in the future.

This volume is "data-rich". That is, not only the results and conclusions of a study are presented, but also the data that led to these results and conclusions, thus encouraging replication and extensions of the analyses that were undertaken, and avoiding the frequent loss of very costly field data. Today's technology for data storage and retrieval are such that we now have no excuse for the frequent and unfortunate losses of original and valuable data as was characteristic of many earlier development projects.

Here, the data in question are available on request in form of 3.5" diskettes, one with the haul-by-haul trawl data analyzed in these pages, the other with the bulk of the fish length-frequency data collected during the surveys covered by this book.

The present volume, and its precursors, resulted not only from collaboration between institutions, but also between individuals. I conclude, therefore, by commending the editors for having expanded such collaboration to include contributions from colleagues from numerous countries not initially involved in the trawl surveys reported upon here.

Dr. Meryl Williams
Director General
ICLARM

Kata Pengantar ICLARM

Pusat Lembaga Internasional Manajemen Sumberdaya Hayati Perairan (ICLARM), merasa berbahagia menerbitkan buku ini yang menyajikan informasi sumberdaya perikanan Indonesia bagian barat.

Kami percaya bahwa buku ini berharga karena berisi penelaahan suatu lokasi yang secara geografis mengandung kehidupan laut, yang kemungkinan, secara biologis memiliki sumberdaya yang paling beragam dan paling kaya akan kelompok-kelompok fauna dan flora. Selain daripada itu, buku ini juga mempunyai keterkaitan di luar perairan Indonesia. Dasar pengetahuan kita masih baru dan sangat sedikit dalam hal perikanan demersal daerah tropis, dibanding dengan pengetahuan kita akan perikanan demersal di daerah beriklim sedang. Hanya sejumlah kecil daerah tropis yang mempunyai analisis ilmiah yang komprehensif semacam ini yang meliputi bidang oseanografi, iklim, ekosistem dan biologi dari sebagian besar spesies ikan yang paling banyak tertangkap. Suatu nilai tambah dari buku ini adalah dengan diciptakannya landasan baru dalam analisis keanekaragaman ikan.

Selain menyajikan hasil berbagai survei sumberdaya yang disusun dalam berbagai bab, buku ini melengkapi beberapa penelitian penting yang telah dilakukan sebelumnya. Salah satunya adalah sebagaimana yang diterbitkan dalam buku yang berjudul "Trawled fishes of Southern Indonesia and Northwestern Australia" oleh Thomas Gloerfelt-Tarp dan Patricia Kailola, yang diterbitkan pada tahun 1984 oleh Badan Bantuan Internasional Australia (AIDAB - yang sekarang bernama AUSAID), Canberra, Australia, Direktorat Jenderal Perikanan (DGF), Jakarta, Indonesia, dan Lembaga Bantuan Teknik Jerman (GTZ), Eschborn, Jerman. Ketiga lembaga ini, dengan bantuan Organisasi Pangan Sedunia PBB (FAO), dari tahun 1978 hingga 1981 telah melaksanakan suatu rangkaian survei perikanan dengan alat trawl di perairan Indonesia bagian barat, dan buku ini telah mendokumentasikan dan membuat ilustrasi bermacam-macam jenis ikan yang tertangkap dalam survei-survei tersebut.

Buku yang diterbitkan ini, yang secara sengaja disusun sebagai pelengkap buku-buku terdahulu yang bernafaskan taksonomi, menyajikan berbagai analisis komunitas secara terperinci dan analisis lainnya terhadap kegiatan-kegiatan survei ini dan kegiatan survei lain yang dilaksanakan di daerah yang sama. Selanjutnya, bab terpanjang kedua memperlihatkan bagaimana daftar ikan yang ada dalam buku Gloerfelt-Tarp dan Kailola disajikan kembali dan diperbaharui dengan FishBase 96 (Froese, R. and D. Pauly, Editors. 1996. FishBase 96: concept, design and data sources. ICLARM, Manila, Philippines. 179 p.), yang merupakan ensiklopedia ikan dunia dalam bentuk CD-ROM, dan ini berarti membuat buku diatas tetap hidup, walaupun sekarang sudah tidak terbit lagi.

Selanjutnya, buku ini juga merupakan kelanjutan dari beberapa seri buku yang diterbitkan sebelumnya yang merupakan hasil kerjasama ICLARM dengan beberapa lembaga di Indonesia, khususnya "Indonesian marine fisheries" (Bailey, C., A. Dwiponggo and F. Marahudin, 1987. ICLARM Stud. Rev. 10, 196 p.), dan atlas "Growth, mortality and recruitment of commercially important fishes and penaeid shrimps in Indonesian waters" (Dwiponggo, A., T. Hartati, S. Banon, M.L. Palomares and D. Pauly, 1996. ICLARM Tech. Rep. 17, 91 p.).

Seperti halnya buku-buku pendahulunya, salah satu motivasi terhadap pembuatan buku ini adalah kesadaran dari editor senior akan manfaat berbagai survei trawl di lokasi perairan dimana belum ada kegiatan eksploitasi, sebagai dasar untuk studi keanekaragaman hayati dalam rangka memberikan target sebagai upaya untuk merehabilitasi stok ikan demersal yang sudah menurun maupun untuk membuat estimasi biaya ekonomi bilamana rehabilitasi tersebut tidak dilakukan. Pertimbangan-pertimbangan semacam ini merupakan jiwa dari berbagai proyek ICLARM, dan buku ini menyajikan wahana acuan yang kerap kali dapat kita tengok kembali. Oleh karena itu, upaya ini merupakan arah baru bagi analisis data perikanan tropis dan merupakan hal yang kami usulkan untuk terus dilaksanakan melalui kerjasama penelitian di masa yang akan datang.

Buku ini "kaya akan data". Yang mana ini berarti bahwa buku ini tidak hanya menyetengahkan hasil dan kesimpulan dari beberapa studi, tetapi juga data yang dipakai dalam analisis. Dengan demikian mendorong untuk dilakukannya upaya replikasi dan perluasan analisis, serta selanjutnya menghindari sering hilangnya data lapangan - yang mana data tersebut sangat mahal harganya. Teknologi masa kini untuk penyimpanan data dan penyajiannya kembali sudah sedemikian majunya sehingga tidak ada alasan untuk sering kehilangan data asli yang berharga, hal mana sering terjadi dalam kegiatan proyek pembangunan terdahulu.

Data yang digunakan dalam buku ini tersedia sesuai dengan permintaan dan disajikan dalam dua buah diskette ukuran 3.5"; satu diskette berisi data setiap tarikan trawl sebagaimana dianalisis dalam buku ini, dan diskette kedua berisi data frekuensi-panjang yang dikumpulkan selama survei yang terliput dalam buku ini.

Buku ini, dan buku-buku sebelumnya, tidak hanya merupakan hasil kerjasama antar lembaga tetapi juga antar individu. Oleh karena itu, sebagai penutup, saya menyampaikan penghargaan kepada para editor yang telah mengembangkan kerjasama untuk mengikutsertakan kontribusi para kolega dari berbagai negara, walaupun mereka tidak ikut dalam kegiatan awal survei trawl yang dilaporkan disini.

Dr. Meryl Williams
Direktur Jenderal
ICLARM

Biodiversity and the Retrospective Analysis of Demersal Trawl Surveys: A Programmatic Approach^a

DANIEL PAULY^b

International Center for Living
Aquatic Resources Management
MCPO Box 2631, 0718 Makati City
Philippines

PAULY, D. 1996. Biodiversity and the retrospective analysis of demersal trawl surveys: a programmatic approach [*Keanekaragaman hayati dan telaah ke belakang survei trawl ikan-ikan demersal: suatu pendekatan programatik*], p. 1-6. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

Demersal trawl surveys of tropical shelves were conducted with increasing frequency and sophistication since the 1920s. Designed originally for the purposes of resources and national development, these surveys are now seen to be ideally suited for reinterpretation as biodiversity baseline studies. The conceptual and practical steps required for this are outlined and illustrated with sample surveys conducted from the mid-1970s to the early 1980s in Western Indonesia.

Abstrak

Survei trawl ikan-ikan demersal di paparan daerah tropis semakin sering dilaksanakan dan bahkan dengan peralatan yang semakin canggih sejak tahun 1920-an. Pada awalnya survei tersebut dirancang dalam rangka pengembangan sumberdaya nasional, namun kemudian survei semacam ini dianggap ideal untuk penelaahan ke belakang sebagai studi dasar keanekaragaman hayati. Langkah-langkah praktis dan terencana yang dilaksanakan dalam survei ini dipaparkan dan digambarkan dengan beberapa survei contoh dalam periode 1970-an sampai 1980-an di perairan Indonesia bagian barat.

Introduction

About 90% of the world fisheries catches originate from shelves, i.e., from the shallow waters - down to 200 m surrounding continents and islands. Of this, the overwhelming bulk stems from softbottom, rather than rocky grounds, or reefs, i.e., from trawlable areas (Gulland 1971; Pauly and Christensen 1993).

Demersal trawl surveys certainly represent the most straightforward way of finding how much and what kind of fish occur in a given softbottom area - not least because they use a gear type (demersal trawls) - initially developed for commercial fishing.

Depending on the distributions of their "stations" (or "drags" or "hauls"), three types of surveys may be distinguished:

- opportunistic,
- systematic and
- (stratified) random.

The first of these, mainly of historical interest, consists of a research vessel (e.g., the *Beagle*, or the *Challenger*) fishing occasionally, but without aiming at representative coverage. Systematic and random surveys differ from opportunistic surveys in that both are planned to cover a given *area*, the former through series of hauls placed along parallel lines as

for the *R/V Mutiara 4* surveys (see Pauly et al., this vol. and Venema, this vol.), the latter through the statistically more powerful randomization of the position of hauls as for the *R/V Fridtjof Nansen* surveys (see Bianchi, this vol.). The statistical power of random survey can be further increased through stratification, a topic that need not concern us here (but see contributions in Doubleday and Rivard 1981; Pauly 1984).

Important here, rather, is that demersal trawl surveys, though affected by mesh selection and gear avoidance, represent an effective method for obtaining *representative* samples of a (bottom) fish community, covering a wide range of sizes, far more so than for other gears deployed to catch coral reef fishes or pelagic fishes, not to speak of gears for catching terrestrial vertebrates.

This evidently is the reason why trawl surveys are conducted to assess the potential of fisheries and to monitor them, once they have developed. Moreover, and this is the theme of this contribution, trawl surveys, conducted several decades ago to estimate demersal fish biomasses, i.e., to provide the basis for the development of demersal fisheries can be turned, through appropriate retrospective analyses, into baseline studies for coastal biodiversity, and thus to help meet a contemporary challenge.

^aICLARM Contribution No. 1314.

^bAnother contact address: Fisheries Centre, the University of British Columbia, 2204 Main Mall, Vancouver, B.C. Canada V6T 1Z4; e-mail: pauly@fisheries.com

Before presenting the elements required for this reconceptualization, I shall recall, however, the key elements of the "development" approach prevailing two to three decades ago, when several major surveys were conducted, both in the tropics, e.g., the Guinean trawling survey in West Africa (Williams 1968), the Gulf of Thailand surveys (Ritragasa 1976), the *R/V Anton Bruun* surveys along the Indian Ocean coast (Hida and Pereyra 1966), and in the cold waters of the North Pacific (Alverson and Pereyra 1972).

Demersal Surveys as Stepping Stones for Development

Demersal surveys, in the 1960s-1970s, consisted of the following major elements:

1. Conduct survey (systematic, later random) in a given area, with funds from development agencies.
2. Have on board the best available field biologists, both from the donor and the surveyed countries, complemented by experts on the taxonomy of the groups covered (and publish fish identification guides if deemed appropriate).
3. Identify and measure the entire catch of each haul, and collect additional data on length composition, maturity stages, stomach contents, etc.
4. Estimate total unexploited biomass (B_o) and estimate "potential yield" (P_y) using the so-called "Gulland equation" now known to generate overestimates (Beddington and Cooke 1983), i.e.,

$$P_y = 0.5 \cdot M \cdot B_o \quad \dots 1)$$
 where M is the natural mortality (Gulland 1971);
5. Write a few papers on the biology of some of the fish, based on the data in (3) and (4);
6. Leave one of the data sets in a laboratory of the surveyed country, and bring back the other set(s) to a laboratory of the donor country.
7. Disband the staff and forget all about (6).

These steps may not have occurred in all surveys, but all contained several of the elements in (1)-(7). Together, they illustrate all that was good, and that was wrong with these surveys (see Pauly 1986 for a critique).

Thus, two to three decades ago, tropical fisheries development schemes were straightforward: fisheries scientists were to locate the resources, and estimate the amounts that could be extracted from them: hence the emphasis on potential yields. Then a development bank, devoted to industrializing the fisheries, would pick up that number, divide the annual catch of a typical commercial trawler into it, and out came the number of trawlers which construction was to be funded.

Fig. 1 documents this line of thought; it is based on a graph I did as a graduate student to illustrate my version of the then dominant thinking (or lack thereof) about the transition from small-scale to large-scale fishing in tropical developing countries, then perceived as both beneficial and unavoidable.

What this approach brought us is now well known: massive overcapitalization, collapsed stocks, impoverished

small-scale fisheries and a need to reconceptualize the entire area of fisheries research and development (Pauly 1979; Pope 1979; Beddington and May 1982; Pauly et al. 1989a; Christy 1993; Pauly 1994, in press; Garcia and Newton, 1995).

The above-described trawl surveys did have positive aspects, however, mainly derived from the nature of trawl hauls as representative samples of demersal fish communities. Thus, the efforts that went into items (2) and (3) led to these surveys providing extremely detailed samples of then largely unfished communities - just what is now needed to serve as baseline for biodiversity studies.

Biodiversity Studies and Their Shifting Baseline

Numerous studies attest to the tremendous impact of direct resource uses, and of habitat destruction on biodiversity - an impact that has accelerated recently, but which has occurred everywhere humans were numerous enough, and had the tools and/or the time to modify those parts of nature surrounding them.

Assessing this impact requires comparisons with some baseline - usually, and almost by definition the oldest available survey. This choice is justified by the fact that, given a continued impact, the more recent the baseline survey one uses, the more it will have shifted toward the present, more impacted situation, and the more human impacts will be underestimated (Pauly 1995).

Conversely, the older the study or survey one uses as baseline, the less the impacts will be underestimated, with all that this implies for attributing value to conservation measures, rehabilitation programs, etc.

Therein lies the worth of old trawl surveys, if their age does not imply a loss of reliability: the theme of the next section.

Retrospective Analyses of Trawl Data: the Issue of Quality

The results of demersal trawl surveys - contrary to the results of other types of biological field studies - are fairly standard in form, always consisting of the following elements:

- i. a general description of the survey, involving details on boat, gear, mesh sizes, trawling speed, etc. and applying to all parts (stations) of the survey;
- ii. trawling "stations", each defined, as for oceanographic stations, by a place (usually defined by a location on a map, i.e., a longitude and a latitude), a depth (that of the sea bottom), and a time (hour, day, month, year);
- iii. a list of the fish and invertebrates caught, often by species, sometimes by higher groups (e.g., genera or families);
- iv. the catch taken of each taxon in (iii).

Or in other words: trawl data are typically described by "tables" that are easy-to-computerize (see Alverson and Pereyra 1972 for an early approach to computerization, and Stromme 1992 for a recent one), a theme to which we shall return below.




Fig. 1. Historic sequence of European fishing (and other) boat developments, from ca. 1750 to ca. 1970 (based in part on Muus and Dahlström 1973; assumed to provide a model for the industrialization of fisheries in developing countries. The problem with this "model" is the fact - now obvious - that small-scale fishers cannot be recycled as crew on trawlers nor will simply disappear, once an industrial fleet has been built (adapted from an overhead prepared in 1973 by the author, for presentation at a graduate seminar).

[Gambar 1. Perubahan sejarah perkembangan kapal ikan (dan kapal lainnya) di Eropa dari kira-kira tahun 1750 hingga tahun 1970 (sebagai berdasarkan tulisan Muus dan Dahlström 1973) diasumsikan sebagai model proses industrialisasi perikanan di negara berkembang. Kesulitan dalam model ini adalah kenyataan - yang sekarang jelas bahwa para nelayan perikanan skala kecil tidak dengan sendirinya menjadi awak kapal trawler ataupun berhenti profesi dengan dibangunnya armada perikanan industri (diambil dari materi presentasi transparansi penulis yang disampaikan dalam seminar mahasiswa tingkat pasca sarjana tahun 1973 dengan sedikit perubahan).]

To ensure the quality of trawl data one must therefore:

- have access to the general description in (i) and be able to verify its integrity;
- be able to duplicate the positions given in (ii) through a reconstruction of the sailing tracks and vice versa, and to check their conformity with modern bathymetric and bottom structure maps;
- be able to assign current names to the taxa in (iii), which may have outdated names.

Item (c) is the only one that poses real problems; in fact the difficulties involved here would be insuperable were it not for the existence of FishBase, the electronic encyclopedia of fish, which has special routines for the identification of valid (new) names, given (old) synonyms and countries of occurrence (see Froese, this vol.).

Fish generally contribute over 90% of the catch of demersal trawls, and the invertebrates caught along with fish

are commonly grouped into larger categories (e.g., "squids"; "sponges", etc., for which translation from "old" to "new" is no problem).

Thus, because of the "tabular nature" of the results of demersal trawl data (see Table 1 for an example), a great amount of trawl survey data can be straightforwardly used, once the table legend (i.e., the general aspects of a survey) and the key column (i.e., the fish names) are verified.

The problem with old demersal surveys, it turns out, is thus not related to quality, but to quantity: it is expensive to have old data computerized (this cannot be reliably done by scanning the original data sheets, as many think). However, modern database techniques can help here, e.g., through preprogrammed entries (e.g., of the species names, which need not be reentered, but can be chosen from a choice list, as in the system developed by Vakily 1992).

Table 1. A typical trawler catch (45-minute haul) from Java Sea (06° 12'S, 108° 26'E, 34-35 m depth) made on 5 September 1976 by *R/V Mutiara 4* in the Java Sea. (Asterisks refer to weight and number raised from a sorted sample of one out of five boxes. Invertebrates not included; see Pauly et al., this vol., for details on the gear used).

[Tabel 1. Hasil tangkapan khas dari trawl (45 menit tarikan) yang dioperasikan di Laut Jawa (06°12'LS, 108°26'BT, pada kedalaman 34-35 m) tanggal 5 September 1976 oleh kapal penelitian Mutiara 4. Tanda bintang menandakan satuan berat dan angka, yang diperoleh dari proses pemilihan contoh (1 dari 5 boks). Tidak termasuk hewan avertebrata; untuk perincian alat yang dipergunakan lihat Pauly et al., vol. ini.]

No.	Family	Species	W(kg)	N
1	Ariidae	<i>Osteogeniosus militaris</i>	3.4	17
2	Balistidae	<i>Abalistes stellaris</i>	0.5	1
3	Carangidae	<i>Seriolina nigrofasciata</i>	0.32	1
4	Carangidae	<i>Scomberoides</i> sp.	0.15	5
5	Carangidae	<i>Alepes kalla</i>	5.0*	90*
6	Carangidae	<i>Alepes djedaba</i>	7.5*	290*
7	Carangidae	<i>Megalaspis cordyla</i>	8.5*	170*
8	Carangidae	<i>Selaroides leptolepis</i>	0.25*	10*
9	Carangidae	<i>Carangoides</i> spp.	6.1*	145*
10	Carangidae	<i>Atropus atropus</i>	1.75*	30*
11	Chirocentridae	<i>Chirocentrus dorab</i>	0.8*	5*
12	Clupeidae	<i>Anadontostoma chacunda</i>	0.15*	5*
13	Clupeidae	<i>Opisthopterus valenciennensis</i>	1.1*	15*
14	Clupeidae	<i>Dussumieria acuta</i>	1.7*	50*
15	Clupeidae	<i>Ilisha</i> sp.	5.6*	65*
16	Clupeidae	<i>Sardinella gibbosa</i>	0.3*	10*
17	Dasyatidae	not identified	2.65	1
18	Drepanidae	<i>Drepane longimana</i>	0.35*	5*
19	Engraulidae	<i>Stolephorus</i> spp.	21.0*	4,175*
20	Gerreidae	<i>Pentaprion longimanus</i>	15.25*	1,165*
21	Fistulariidae	not identified	0.15*	10*
22	Formionidae	<i>Formio niger</i>	0.2	1
23	Lagocephalidae	not identified	4.0	95
24	Leiognathidae	<i>Leiognathus splendens</i>	10.0*	720*
25	Leiognathidae	<i>Leiognathus leuciscus</i>	4.2*	780*
26	Leiognathidae	<i>Leiognathus bindus</i>	1.2*	340*
27	Leiognathidae	<i>Secutor ruconius</i>	1.2*	380*
28	Leiognathidae	<i>Secutor insidiator</i>	2.8*	560*
29	Lutjanidae	<i>Lutjanus sanguineus</i>	4.0	1
30	Lutjanidae	<i>Lutjanus johni</i>	5.0*	10*
31	Lutjanidae	<i>Lutjanus lineolatus</i>	0.2*	10*
32	Lutjanidae	<i>Caesio erythrogaster</i>	0.1*	5*
33	Mullidae	<i>Upeneus sulphureus</i>	75.0*	6,075*
34	Nemipteridae	<i>Nemipterus japonicus</i>	3.0*	15*
35	Nemipteridae	<i>Nemipterus bathybius</i>	0.4*	15*
36	Pentapodidae	<i>Pentapodus setosus</i> (?)	0.25*	5*
37	Platycephalidae	not identified	0.25*	5*
38	Plectorhynchidae	<i>Plectorhynchus pictus</i>	0.4*	15*
39	Pomadasyidae	<i>Pomadasyus maculatus</i>	0.25*	5*
40	Pomadasyidae	<i>Pomadasyus</i> sp.	0.5*	35*
41	Priacanthidae	<i>Priacanthus macracanthus</i>	3.1*	80*
42	Scombridae	<i>Scomberomorus guttatus</i>	7.2*	65*
43	Scombridae	<i>Scomberomorus commerson</i>	2.6	14
44	Scombridae	<i>Rastrelliger brachysoma</i>	3.0*	50*
45	Stromateidae	<i>Pampus chinensis</i>	0.75	1
46	Stromateidae	<i>Pampus argenteus</i>	6.3*	30*
47	Synodontidae	<i>Saurida tumbil</i>	0.35	1
48	Synodontidae	<i>Saurida elongata</i>	3.75*	45*
49	Synodontidae	<i>Saurida longimana</i>	0.9*	105*
50	Sphyrinaeidae	<i>Sphyrina obtusata</i>	0.6*	10*
51	Sciaenidae	not identified	0.25*	5*
52	Terapontidae	<i>Therapon</i> sp.	3.75	100
53	Triacanthidae	not identified	1.0*	25*
54	Trichiuridae	<i>Trichiurus lepturus</i>	1.0*	55*
55	Trichiuridae	<i>Lepturacanthus savala</i>	2.0*	25*
Σ	29 families	43 genera and over 55 spp.	232.02	15,948

Performing Retrospective Analyses: Technical Aspects

Performing analyses of the tabular demersal trawl data presented above may involve:

- reproducing the result of the old survey (see, e.g., contribution by Martosubroto et al., this vol.);
- performing new multivariate (community) analyses (see, e.g., Bianchi, this vol.; McManus, this vol.), related, where possible, to oceanographic and other features of the environments in question (see Sharp, this vol. and Roy, this vol.);
- using the biological and size composition data collected along with the catch data to describe the biology and estimate vital statistics of various species (see, e.g., Pauly et al., this vol.), then to use these and ancillary data (including catches) to construct one or several trophic models of the ecosystem in question (see contributions in Pauly and Christensen 1993 and in Christensen and Pauly 1993.);
- mapping distributions onto phylogenetic "trees" to separate taxa that evolved locally from those that immigrated after they had evolved elsewhere (Brook and McLennan 1991);
- performing other analyses, e.g., relating fish weights and numbers through ecological stress indicators (Warwick 1986; McManus and Pauly 1990), or assembling standardized datasets allowing global, simultaneous comparisons of community analyses from a number of surveys, spanning the intertropical belt.

The hardware and software tools exist for analyses that could not have been performed before, and thus for understanding features of tropical fish communities not before apprehended. Indeed, I expect that the biodiversity baseline data emerging by computerizing "old" trawl survey will lead to true discoveries, or at least rigorous test of earlier theories, such as, e.g., A.R. Longhurst's perception of roughly similar communities on both sides of the tropical Atlantic, and of their (distorted) mirror images in the Indo-Pacific (see Longhurst and Pauly 1987).

At the level of individual species, the methods now available for the analysis of length-frequency data (Pauly and Morgan 1987; Gayanilo et al. 1996) enable routine estimation of at least some of the vital statistics of the major species covered by a trawl survey. Moreover, electronic access to the literature on each of these species, through the Aquatic Science and Fisheries Abstracts CD-ROM, and to summaries of biological information on these same species, through the FishBase CD-ROM (Froese and Pauly 1996) will allow quick identification of knowledge gaps, comparisons of results among species, and verification of species lists (Froese et al., this vol.).

Box 1. Uses of boxes.

[Boks 1. Penggunaan boks.]

Most contributions in this volume include "boxes," presenting materials relevant to, but not part of their main narrative.

The use of boxes to present such material has enabled exploring the antecedents (and/or follow-up) of some important issues presented in this book by the authors themselves or by invited contributors. Notably, the development of several fisheries initiated after - and sometimes as a result of - the surveys described here was briefly followed up, e.g., by J. Widodo et al. (see Boxes 3 and 4 in Venema, this vol.) and by A. Ghofar and C. Mathews (see Pauly et al., this vol.). The latter indeed presented a surprisingly close fit between the fluctuation of the important stock of *Sardinella lemuru* in the Bali Straits and the occurrence of El Niño/Southern Oscillation (ENSO) events, a result made possible by their joint analysis of two time series that had hitherto remained disconnected.

We hope that the data assembled for this book and which are available in digital form (see Torres et al., this vol.) will encourage further discoveries of this sort.

Pauly et al. (this vol.) illustrated this new integrated and systematic approach, meant to replace the scattered analyses, covering a few species at best, that have traditionally followed trawl surveys.

Retrospective Analyses: the Institutional Aspects

Although computerization does reduce the workload, few fisheries institutions, or universities can perform all analyses of the data emanating from a set of surveys.

Rather, the scheme used for this book may be recommended: a large number of colleagues, belonging to several institutions were identified and convinced to contribute analyses, each dealing with a subarea, or a period covered by a (set of) survey(s) or even with a "side" aspect of a survey or its consequences (Box 1). This obviously will work only when a project leader, or a small group agree beforehand to help the authors standardize their approach, a *laissez faire* attitude being here completely inappropriate.

Standardization includes - among other things - agreeing on:

- an area (here Western Indonesia: note the consistency between the base maps presented in the various contributions in this book);
- a period (here 1974 to 1981);
- a file format (see Torres et al., this vol.).

(The approach proposed here is evidently the same as that used for previous studies of the anchoveta *Engraulis ringens* and its upwelling ecosystem, covering, on a monthly basis, from 1953 to 1982, the area off Central and Northern Peru, see contributions in Pauly and Tsukayama 1987 and Pauly et al. 1989b).

The large number of authors, from both developing- and developed-country institutions that will have to come together for relevant products to emerge also allows dealing with the thorny issue of scientific credit: all participants can author, or at least co-author a part of the whole story (the issue of credit is further discussed in Pauly 1986, 1993).

Moreover, the standardized databases that emerge from an exercise such as proposed here will become available - in a way that the older, original data were not - to the scientists of the countries in which the surveys were conducted and to the international scientific community. The volume describing the database will thus be a "data-rich book" *sensu* Pauly (1993, 1994), and contribute to solving the data loss problem addressed, e.g., in Janzen 1986 or Mathews 1993, as well as in other disciplines (Levitus et al. 1994).

Readers interested in the application of the approach outlined here to other tropical areas are welcome to contact ICLARM and/or the author.

Acknowledgements

The programmatic approach presented here was inspired by discussions with and the work of Dr. Gabriella Bianchi, Dr. *Fridtjof Nansen* Project, Bergen, Norway, and a seminar by Dr. Meryl Williams, in late 1993 in which she pointed out the need for ICLARM staff to reinterpret their work in the context of biodiversity.

References

- Alverson, D.L. and W.T. Pereyra. 1972. Demersal fish exploration in the northeastern Pacific Ocean - an evaluation of exploratory fishing methods and analytical approaches to stock size and yield forecasts, p. 224-254. *In* Proceedings of the Symposium on Demersal Fisheries, Brisbane, Australia. Proc. Indo-Pac. Fish. Council. 13(3).
- Beddington, J.R. and J.G. Cooke. 1983. The potential yield of fish stocks. FAO Fish. Tech. Pap. 242, 47 p.
- Beddington, J.R. and R.M. May. 1982. The harvesting of interacting species in a natural system. *Sci. Am.* 247(5):42-49.
- Brook, D.R. and D.A. McLennan. 1991. Phylogeny, ecology and behavior: a research program in comparative biology. University of Chicago Press, Chicago. 434 p.
- Christensen, V. and D. Pauly. 1992. ECOPATH II - a software for balancing steady-state ecosystem models and calculating network characteristics. *Ecol. Modelling* 61:169-185.
- Christensen, V. and D. Pauly, Editors. 1993. Trophic models of aquatic ecosystems. ICLARM Conf. Proc. 26, 390 p.
- Christy, F.T., Jr. 1993. Back to school. The world's fishery managers should revise their basic economic textbooks: open access is a catastrophe. *CERES, FAO Rev.* 26(142):32-36.
- Doubleday, W.G. and D. Rivard, Editors. 1981. Bottom trawl surveys. *Can. Spec. Publ. Fish. Aquat. Sci.* 58, 273 p.
- Froese, R. and D. Pauly, Editors. 1996. FishBase 96: concepts, design and data sources. ICLARM, Manila, Philippines. 179 p.
- Garcia, S.M. and C. Newton. 1995. Current situation trends and prospects in world capture fisheries. *In* Proceedings of the Conference on Fisheries Management: Global Trends, 14-16 June 1994, Seattle, USA. 63 p.
- Gayanilo, F.C., Jr., P. Sparre and D. Pauly. 1996. The FAO-ICLARM Stock Assessment Tools (FISAT) user's manual. FAO Comput. Info. Ser. (Fish.) 8. FAO, Rome. 126 p.
- Gulland, J.A., Editor. 1971. The fish resources of the ocean. Fishing News Books, West Byfleet, England.
- Hida, T. and W. Pereyra. 1966. Results of bottom trawling in Indian Seas by RV "Anton Bruun" in 1963. Proc. Indo-Pac. Fish. Council. 11:156-171.
- Janzen, D. 1986. Science is forever. *Oikos* 46: 281-283.
- Levitus, S.R., D. Gelfeld, T. Boyer and D. Johnson. 1994. Results of the NODC and IOC Oceanographic Data Archeology and Rescue Project, Rep. 1 NOAA, Key Oceanogr. Rec. Doc. No. 19, 73 p.
- Longhurst, A.R. and D. Pauly. 1987. Ecology of tropical oceans. Academic Press, San Diego, California. 407 p.
- Mathews, C.P. 1993. On preservation of data. *Naga, ICLARM Q.* 16(2/3):39-41.
- McManus, J. and D. Pauly. 1990. Measuring ecological stress - variations on a theme by R.M. Warwick. *Mar. Biol.* 106(2):305-308.
- Muus, B.J. and P. Dahlström. 1973. Meeresfische. BLV Verlagsgesellschaft, München. 244 p.
- Pauly, D. 1979. Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries. *ICLARM Stud. and Rev.* 1, 35 p.
- Pauly, D. 1984. Methods for assessing the marine stocks of Burma, with emphasis on the demersal species. BUR/77/003, FAO Field Doc. 6, 22 p. FAO, Rome.
- Pauly, D. 1986. On identifying fish rather than assessing fish stocks: review of two books on the taxonomy of the neritic fishes of the Western Indian Ocean. *Naga, ICLARM Q.* 9(3):21.
- Pauly, D. 1986. On using other people's data. *Naga, ICLARM Q.* 11(1):6-7.
- Pauly, D. 1993. Data-rich books. *Bioscience* 43(3):167-168.
- Pauly, D. 1994. On the sex of fish and the gender of scientists: a collection of essays in fisheries science. Chapman and Hall, London. 250 p.
- Pauly, D. 1995a. Small-scale fisheries in the tropics: marginality, marginalization and some implications for fisheries management. *In* Proceedings of the Conference on Fisheries Management: Global Trends, 14-16 June 1994, Seattle, USA. 63 p.
- Pauly, D. 1995b. Anecdotes and the shifting baseline syndrome of fisheries. *Trends Ecol. Evol.* 10(10): 430.
- Pauly, D. and G.R. Morgan, Editors. 1987. Length-based methods in fisheries research. ICLARM Conf. Proc. 13, 468 p.
- Pauly, D. and I. Tsukayama, Editors. 1987. The Peruvian anchoveta and its upwelling ecosystem: three decades of change. ICLARM Stud. Rev. 15, 351 p.
- Pauly, D. and V. Christensen. 1993. Stratified models of large marine ecosystems: a general approach and an application to the South China Sea, p. 148-174. *In* K. Sherman, L.M. Alexander and B.D. Gold (eds.) Large marine ecosystems: stress, mitigation and sustainability. AAAS Press, Washington, D.C.
- Pauly, D., G. Silvestre and I.R. Smith. 1989a. On development, fisheries and dynamite: a brief review of tropical fisheries management. *Nat. Resour. Modelling* 3(3):307-329.
- Pauly, D., P. Muck, J. Mendo and I. Tsukayama, Editors. 1989b. The Peruvian upwelling ecosystem: dynamics and interactions. ICLARM Conf. Proc. 18, 438 p.
- Pope, J.G. 1979. Stock assessment in multispecies fisheries, with special reference to the trawl fishery in the Gulf of Thailand. *SCS/DEV/79/19*, 106 p.
- Ritragasa, S. 1976. Results of the studies on the status of demersal fish resources in the Gulf of Thailand from trawling surveys, 1963-1972, p. 198-223. *In* K. Tiews (ed.) Fisheries resources management in Southeast Asia. German Foundation for International Development, Berlin.
- Strømme, T. 1992. NAN-SIS: software for fishery survey data logging and analysis. User's manual. FAO Comput. Info. Ser. (Fish.) No. 4, 103 p.
- Vakily, J.M. 1992. Assessing and managing the marine fish resources of Sierra Leone, West Africa. *Naga, ICLARM Q.* 15(1):31-35.
- Warwick, R.M. 1986. A new method for detecting pollution effects on marine macrobenthic communities. *Mar. Biol.* 92: 557-562.
- Williams, F. 1968. Report on the Guinean trawling survey. Vols. I (601 p.), II (529 p.) and III (551 p.). Publ. Sci. Tech. Res. Comm., Organ. Afr. Unity 99.

Oceanography of the Indonesian Archipelago and Adjacent Areas

GARY D. SHARP
Cooperative Institute for Research
in the Integrated Ocean Sciences
Monterey, California, USA

Oceanography
Geomorphology
Climatology
ISEW, Indonesia
Q1567

SHARP, G.D. 1996. Oceanography of the Indonesian Archipelago and adjacent areas [Oseanografi kepulauan]. p. 7-14. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

Descriptions of the geomorphology and climatology of Southeast Asia are presented as basis for the main oceanographic features of the Indonesian Archipelago. Emphasis is on longer-term global changes and their impact on fisheries resources.

Abstrak

Gambaran geomorfologi dan klimatologi Asia Tenggara disajikan sebagai dasar untuk mengetahui sifat-sifat oseanografi kepulauan Indonesia. Pembahasan ditekankan pada perubahan-perubahan umum jangka panjang dan pengaruhnya terhadap sumberdaya perikanan.

Introduction

Wyrki (1961) reviewed the geomorphological history and oceanography of the Southeast Asian region, bounded by the Bay of Bengal and Andaman Islands to the northwest, the Strait of Formosa to the northeast, North Australia and the Torres Strait to the southeast and the Indian Ocean to the west-southwest. Appropriately, he gave particular emphasis to the regional consequences of the double monsoon; also, he described the dynamics of sea levels as a result of Pacific Ocean climate changes, and of Pacific to Indian Ocean throughflows. The region clearly comprises some of the more complex geomorphological structures on earth, with high mountains in the north, the deepest known ocean trench (off the east coast of Mindanao, Philippines), numerous volcanic islands along the western boundary of the so-called "rim of fire" (Katili 1989), and a vast assortment of coral islands, shallow shelves and deep channels (Fig. 1). Also, the rivers in the region generate most of the silt deposited on continental shelves anywhere on earth (Milliman and Meade 1983; Duinker 1989; Wright 1989).

Topography and Circulation

There are only a few deep channels allowing confluence between the western Pacific and the Indian Ocean. The major ones are the San Bernardino and Surigao Straits within the

Philippine Islands chain, where the deep ocean currents are directed into the eastern portion of the archipelago, to be blocked by the Sunda shelf in the west, and the Arafura shelf in the east. There are, further, three relatively narrow passages that connect the two oceans: one at Lombok, and two on either extreme of Timor, herein referred to as Timor north and Timor south.

Several shallow basins form the Gulf of Thailand, the South China Sea and the Java Sea. To the east the deep Pacific Ocean reaches from the Celebes Sea southward to Lombok (116°E, 9°S) via the Makassar Strait. Similarly, the Banda Sea opens to the Timor Sea through the Barat Daya and Tanimbar Islands, along 5°S, from 126-135°E. The deep Ombai Strait connects the Banda Sea with the Sawu Sea along Timor north. The surface winds sweep warm surface waters southward throughout the northwest monsoon, from November to February, and the southwest monsoon pushes surface waters northeastward, back through the shallow sills and strait

Pacific Ocean deep water can only pass through the deep channels such as the Makassar Trough, which is only 2,540 m at its deepest, with sills at about 2,300 m depths, while the Timor south channel between Timor and Babar Island (the Timor Trench) reaches 3,310 m. To the north, the narrow Ombai Strait (≈ 5 km) between Timor and the Ator and Barat Daya Islands reaches down to only 1,600 m, connecting with the Sawu Basin which reaches 3,470 m. The deep channels that connect the Flores and Banda Seas with the Indian Ocean have sills between 1,900 and 2,100 m (Postma and Mook 1988; van Aken et al. 1988).

It is presently not well understood what the decadal scale changes in relative forcing might be, but one should assume that they are significant. The Asian continent warming in northern summer and that of Australia in austral summer both act as engines for monsoons. Interannual and decadal climatic patterns and trends reflect the even larger geographic scale of the events affecting the Indonesian archipelago, as consequences of a quasi-biennial oscillation, the El Niño-Southern Oscillation (ENSO), and of longer and global scale climatic and oceanographic processes. Fig. 2 shows some of the global features involved here.

Regional Climate

The region's double rainy season exhibits two peaks, at the apex of each monsoon period. The double rainfall peak and relatively large terrestrial runoff strengthen the thermal stratification of the upper ocean throughout the region. The shoal warm water that derives from the northern and western areas of the archipelago is therefore relatively lower in salt content than is that deriving from the Pacific Ocean. The north monsoon, which starts in October and peaks in January,

Fig. 2. The extent of the warm tropical (bright) ocean into the global ocean as measured during August (austral winter) when the least tropical heating is expected. Note that at 30 m, the surface ocean is contiguous in the tropical regions, except along West Africa and South America, where the cool Benguela and Humboldt currents dominate; note also that there are no tropical temperatures at 90 m off either of these continents. The deeper tropical influence in the southern Indian Ocean is evidenced by the continuity between the northwest Australian region and Madagascar, even to 125 m. At that depth, only the far western Pacific Ocean, and the region from the Caribbean eastward, and south along Brazil remain strongly tropical. The only complete bridging of any ocean basin at these depths is in the southern Indian Ocean.

[Gambar 2. Luasnya pengaruh laut tropis yang hangat (warna terang) terhadap perairan laut secara keseluruhan selama bulan Agustus (musim dingin di Australia) dimana pengaruh panas tropis sangat kecil. Pada kedalaman 30 m, permukaan laut di daerah tropis seragam, kecuali pada sepanjang pantai Afrika Barat dan Amerika Selatan, dikarenakan adanya pengaruh dominansi arus dingin Benguela dan Humboldt; perlu disimak bahwa tidak ada suhu tropis pada kedalaman 90 m di dekat kedua benua ini. Pengaruh tropis di perairan yang dalam di Samudra Hindia bagian selatan terlihat dari kontinuitas suhu antara wilayah perairan Australia Barat Laut dan Madagaskar, bahkan sampai kedalaman 125 m. Pada kedalaman ini, hanya Samudra Pasifik paling barat, dan wilayah perairan laut dari Karibia ke arah timur serta sepanjang pantai Brazil yang tetap bersifat tropis. Perairan di laut dalam yang saling berhubungan pada kedalaman ini hanyalah di bagian selatan Samudra Hindia.]





Fig. 1. Relief map of the Southeast Asian region, showing the deepest ocean trenches, some of the world's highest mountains, immense lowlands and shallow archipelagos. Combined with the topographic extremes, climatic forcing from both the Indian and Pacific Oceans, and the continental effects of Asia and Australia, these features generate a dynamic environment whose complexity is reflected by that of the biota.

[Gambar 1. Atlas relief wilayah Asia Tenggara yang menunjukkan palung laut terdalam, sebagian gunung-gunung yang tertinggi di dunia, dataran rendah yang sangat luas dan laut di sekitar kepulauan yang dangkal. Keadaan topografi yang ekstrim, yang dikombinasikan dengan kekuatan energi klimatis dari Samudra Hindia dan Pasifik serta pengaruh benua Asia dan Australia, menciptakan kondisi lingkungan yang dinamis yang mana kompleksitasnya tercermin dari sumber daya hayatinya.]



pushes the surface currents from north to south until reaching the equator, when they are deflected southeastward.

Rates of throughflow depend on both sea level and surface wind speed and direction. The south monsoon peaks in July-August, when southward transport dominates throughout the water column. This is because surface winds affect primarily the surface currents, and there is always a substantial sea level difference that favors movements of water from the Pacific to the Indian Ocean. These currents from the Pacific Ocean, through the Mindanao Strait into the Celebes Sea can exceed $100 \text{ cm}\cdot\text{s}^{-1}$ (Postma and Mook 1988). They are then channeled southward through the deep Makassar Strait.

During the south monsoon, the winds force the surface waters westward from the Makassar Strait into the Java Sea (Martosubroto, this vol.). Conversely, the north monsoon forces the surface waters southward, into the Flores Sea, and on through the Timor north and south channels, into the Indian Ocean.

Due to the near continuous monsoon wind forcing, both convergent and divergent, the eastern archipelago is characterized by a continuous 40-50 m mixed layer. However, there is some seasonal upwelling in the Banda and Arafura

Sea regions during the south monsoon, leading to shoal oxygen minima, and consequent ecological responses. Because of the very strong north monsoon winds, mixed layer depths can reach 100 m during peak periods. Oxygen is depleted rapidly due to biological activity below the mixed layer; the oxygen minimum ranges from 1.8 to $2.0 \text{ ml}\cdot\text{l}^{-1}$ at depths greater than 350 m, with local seasonal minima reaching lower values. Wyrki (1962) also addressed the upwelling (divergence) between Java and Australia, of which a faint signal might be seen in Fig. 3 (April).

Sea surface temperature (SST) varies across the north-south extent of the archipelago, with the greatest variations occurring in the northeast. For example, in the Strait of Formosa, seasonal SST ranges from under 13°C to over 26°C . To the south, within the archipelago and into the northwest Australian region, temperatures range from about 24°C to over 29°C seasonally; this warm, lower salinity surface water is locally responsive to surface forcing. Where topography permits, higher salinity water from the western Pacific Ocean passes over deep sills from east to west in response to the combined dynamics of sea level differences, surface wind forcing and regional barometric pressure, varying on seasonal and longer-time scales.




Fig. 3. One degree latitude-longitude mean temperatures at 90 m from the MOODS files (Bauer 1985; Bauer and Robinson 1985). The north monsoon (October to March) forces north-south surface currents, which are reflected in the January warming along the north coast of Australia. In April, the south monsoon begins, and forces surface transport from the Indian Ocean northward into the archipelago. Note that upwelling is indicated by the green coding, correlated with low oxygen levels.

[Gambar 3. Suhu rata-rata pada kedalaman 90 m pada tiap derajat lintang dan bujur sesuai arsip MOODS (Bauer 1985; Bauer dan Robinson 1985). Musim angin utara (Oktober sampai Maret) mendorong arus permukaan utara-selatan sebagaimana tercermin dari pemanasan bulan Januari di sepanjang pantai utara Australia. Pada bulan April, mulailah musim angin selatan yang mendorong arus permukaan dari Samudra Hindia ke utara ke wilayah kepulauan Indonesia. Proses upwelling digambarkan dengan warna hijau yang berkaitan dengan konsentrasi kandungan oksigen yang rendah.]

Recent Interests in Southeast Asia's Environment

The Indonesian Archipelago and the surrounding oceanic and terrestrial systems form a unique set of challenges to ocean modelers and would-be climate forecasters. The principal problem is that the confluence of the western Pacific "warm pool" and the Indian Ocean is severely restricted within the archipelago.

In addition to topography, there are several climatic factors that affect the flow rates between the two ocean basins. The tropical monsoons directly affect the direction of the surface water motion. At ENSO time scales, the changes in regional sea level during the various phases of the ENSO cycle affect the relative strengths and directions of both surface and subsurface flows, a problem first addressed by Wyrki (1961). He found that during the southwest monsoon (April-September) water exchange occurs at the 125-300 m level while during the northwest monsoon (October through February), the flow is greatest at about 1,000 m.

More recently, Allan and Pariwono (1990), who reviewed the works of Hastenrath and Lamb (1979a, 1979b), Hastenrath and Wu (1982), Nichols (1984a, 1984b), and Hackert and Hastenrath (1986), with emphasis on the relations among SST, radiative effects (clouds) and regional wind forcing suggested that, along with these other important variables, "SST advection and ocean mixing must also be included when considering ocean-atmosphere interactions in eastern Indonesian waters."

Allan and Pariwono (1990) also suggested that ocean mixing is more likely to be important during winter when wind strengths are greater and near-surface flows to the north overlie southward deep throughflows, creating pronounced vertical shear. In the summer, both surface currents and deeper flows are directed southward, and the ocean mixing layer is shallow. During ENSO warm phases, stronger westerlies result in stronger advection. This, coupled with southerly-directed throughflows at depth, results in stronger and more penetrative upper ocean mixing. During cool ("La Niña") events, the opposite is true. Allan and Pariwono (1990) concluded by stating that the sparseness of observations in the Indonesian archipelago precludes more precise estimations of throughflows from the Pacific Ocean to the Indian Ocean, and that due to the above interactions, and the poorly understood consequences of rainfall within the region, a much more complex structure of ocean dynamics will eventually be described.

Wind speed and direction, within the ENSO cycle, have dramatically different consequences on upper ocean temperatures, although the seasonal temperature variations are relatively small within the shallow shelves of the archipelagos ($\pm 1.5^\circ\text{C}$ around a mean of 28.5°C). The deep convection associated with strong early ENSO easterlies enhances the dynamics of both SST and sea level. Increases of the latter induce greater flowthrough, and greater SST leads to enhanced convective transport of upper ocean heat at onset of the pre-summer (October-November) equatorial Pacific westerlies. These trigger Kelvin waves, increased upper ocean

heating and deep convection toward the date line, thus promoting conditions conducive to SST rise into the eastern Pacific Ocean. The consequences are a decline in rainfall within the archipelago, and eastward propagation of deep convection and associated storm tracks in the central and eastern Pacific. All of these stages are prognostic of El Niño conditions along the west coast of the Americas.

ENSO-Related Dynamics of the Larger Region

Due to the seasonal dynamics of the Indian Ocean, sea level fluctuations are quite dynamic, reflecting both monsoonal processes and the longer-term ENSO atmospheric pressure fields. During the pre-Niño buildup when persistent westerlies sustain the surface flows across the Pacific Basin, and the southwest monsoon has driven across the Indian Ocean to Indonesia, sea level rises, sometimes reaching knee depth in the streets of downtown Jakarta. Also, upper ocean thermocline structures deepen. Regional drought and raging forest fires are concurrent with sea level rise. In fact, these are precursor signals that provide insights about the eminent release and progression of a Kelvin wave toward the Americas, itself a precursor of El Niño conditions.

Thus, while heat and drought prevail in Southeast Asia and North Australia, the western coasts of the Americas see warmth and deluges. As sea level drops around and in Indonesia, it increases in the eastern Pacific Ocean, reflecting the eastward displacement of vast volumes of ocean midwater. A major consequence of these latter stages of El Niño, i.e., of an ENSO warm event, is the return of a "normal" tropical thermal regime to the equatorial latitudes of South America. It is still not sufficiently recognized that the anti-Niño phase promotes an eccentricity in the global equatorial ocean system in that 14 to 20°C surface waters dominate the coastal and eastern equatorial regions, with profound local and global consequences. This asymmetry destabilizes the entire global thermal balance, and is a major contributor to extraseasonal climate forcing.

The easternmost edge of the warm pool (defined here as some temperature between 27.5 and 28.5°C) has been used as an indicator of ENSO warm events (Fig. 4). Conversely, the depth of the 26 or 28°C isotherm in the western Pacific Ocean can be used to infer the intensity of the heat gains or losses during the ENSO cycle. These are more difficult to portray meaningfully, due to the broad regions involved, and their greater local dynamics. However, once the Kelvin wave is released, sea level falls, westward surface currents decline and the Indo-Pacific throughflow declines. The gravity wave progresses eastward, allowing a shoaling of the western Pacific thermal structure. One little known effect is that the upper ocean thermal structure shoaling promotes catches of the larger tunas normally associated with the deeper, cooler thermal structures. In fact, their appearance in west Pacific purse seine catches is symptomatic of the Kelvin wave, and of a pending decline in catches in the eastern Pacific Ocean, when thermoclines are depressed by that same Kelvin wave. From this, one can




Fig. 4. Time series of ENSO-related processes. Upper panel: SST in the El Niño region (offshore of South America). The rise of SST precedes the migration eastward of the 28.5°C isotherm and denotes local heating, not eastward surface advection of heat. There is no emergent trend, only two unusual events in the sequence since 1950. Lower panel: eastern extent of the 28.5°C surface isotherm. (Source: Environmental Research Laboratories of NOAA, Boulder, Colorado, USA). [Gambar 4. Data tahunan dari proses yang berhubungan dengan ENSO. Grafik atas: SST di daerah El Niño (perairan lepas pantai Amerika Selatan). Naiknya SST mendorong isotherm 28.5°C bergerak ke timur dan menyebabkan pemanasan lokal, bukan adveksi panas permukaan yang ke timur. Tak ada trend tertentu yang muncul, hanya dua kejadian yang lain dari biasanya yang secara berurutan terjadi sejak tahun 1950. Grafik bawah besarnya isotherm permukaan 28.5°C di bagian timur. (Sumber: Environmental Research Laboratories of NOAA, Boulder, Colorado, USA).]

appreciate that there is more to the ENSO cycle than El Niño warming off the western Americas (see also Ghofar and Mathews, in Pauly et al., this vol.).

The variability among individual ENSO cycles is great, and thus the concept of canonical ENSO cycle has been tossed away, as should also the notion that an east to west movement of warm upper ocean water ensues. Apparently the single most common misunderstanding about the ENSO warm and cool events is a presumption that there is eastward transport of warm surface waters during the initial Kelvin wave that defines the onset of an ENSO warm event. As discussed, each El Niño is triggered by weakening or reversal of tropical westerlies around the date line. The Kelvin wave, a gravity wave, propagates eastward during the latter stages of an ENSO cycle, promoting depression of the thermocline as it passes eastward, as well as local warming of the upper ocean. The Kelvin wave then rapidly passes poleward along the coast, with dramatic consequences that have been well described.

Southeast Asian Archipelago and Downstream Ecological Consequences

A quick examination of Figs. 2 and 3 will show that the temperatures of the surface flow during the southwest monsoon period are sufficient to transport a substantive amount of warm surface waters from the Pacific warm pool southeastward, across the Indian Ocean to Madagascar, and thence to the South African coast. Note also that, due to the shoal isotherm structures of the equatorial Indian Ocean, only in May, June

and July can warmer superficial waters be transported into the northeastern Indian Ocean.

There is little direct environment observation data in the literature or in accessible archives for the entire Indo-Pacific region. Much of what is known derives from relatively short-term endeavors, and much of what we would like to know more about now has to be inferred from what is known about the dynamics of the surrounding environment. The compelling data sets that I would like to add to this picture include the broader perspectives of the western and southern Indian Ocean, using monthly time series for depths of the 14°C isotherms as proxies of the decadal scale dynamics of the Warm Pool. This stimulating data set derives from the Master Ocean Observation Set (MOODS). The monthly mean 3x3 degree latitude-longitude data were compiled along the South African coast from Somalia, south through the Mozambique Channel, around into the Benguela upwelling region. Fig. 5 provides information about the recent trends in upper ocean heat content, and changes in patterns from the early 1950s to the late 1970s. The deepening (brightening) of the isotherms around the Cape of Good Hope during the late 1970s until recently was accompanied by the onset of a substantial tropical tuna fishery for yellowfin (*Thunnus albacares*), skipjack (*Katsuwonus pelamis*) and the resurgence of sardines in the coastal environment. The lavender region denotes shoaling of the 14°C isotherm (strong upwelling) within the Benguela upwelling region.

Shackleton (1987) found 20-25 year patterns of coastal pelagic fish scale abundances during examination of laminated sedimentary materials from Walvis Bay, Namibia. Eight cycles




Fig. 5. Time series of the 14°C isotherm, the Congo (equatorial Atlantic) southward through the cool Namibia-Benguela upwelling, around the Cape of Good Hope into the Indian Ocean, i.e., the Mozambique Channel and northward to Somalia's east coast.

[Gambar 5. Isotherm 14°C berdasarkan waktu dari Congo (Atlantik ekuator) ke arah selatan melalui daerah upwelling dingin Namibia-Benguela, melingkari Tanjung Harapan masuk ke Samudra Hindia, melalui selat Mosambik dan ke utara hingga pantai timur Somalia.]

were observed for the 200-year record. There were prolonged periods within these sediments when sardine scales were present (warming periods) and absent (cool, upwelling dominated periods). This suggests that perhaps the natural decadal scale waxing and waning of the Pacific Warm Pool might contribute through strengthening of the southwestward transport of warm upper ocean water during periods of maximal western Pacific Ocean sea level, and throughflow periods associated with strong convergence due to ENSO pressure patterns. Ultimately, the western Indian Ocean consequence of a strong surface throughflow appears to be a suppression of upwelling due to the overlaying of warmer surface waters along the South African coast, with significant ecological results.

Strong equatorial convergences and countercurrents preclude surface water exchanges between the northern and southern Indian Ocean. The deepening of the upper ocean along the southwestern convergence acts as a significant pathway for heat transport from the warm pool to the southwest Indian Ocean, particularly during the northeast monsoon.

The rest of the year, the northern Indian Ocean circulation patterns are dominated by the wind-driven upwelling of cooler oxygen-poor water along the equatorial ocean and along northern coastlines, excluding valuable fish species from their usual deep habitats, and locking out major surface flows from the Pacific Ocean into the northern Indian Ocean (Sharp 1979). The situation is sufficiently limiting for both the Arabian Sea and the Bay of Bengal to become - in ecological terms - nearly enclosed waterbodies. The northern portion of the Arabian Sea, although capable of large surficial exchange, is also affected by emergence of poorly oxygenated waters during substantial

periods of the year. However, this is a local phenomenon, and in spite of local oxygen minima, the shoal shelf faunas of the Indo-Pacific consist, if not of exactly the same species, at least of congeners, from Papua New Guinea through Southeast Asia, around the Bay of Bengal and Arabian Sea, and down the east coast of Africa to Madagascar (Longhurst and Pauly 1987; Sharp 1988).

Rich species arrays are found in several areas where complex habitat structures, e.g., rocky shelves and reefs, are collocated with strong, patterned seasonal environments that prevent poorly oxygenated waters from emerging at the surface, e.g., the north coast of Somalia and around the fringes of the Indonesian Archipelago. Although the oceanic species of the equatorial Indo-Pacific are nearly all shared, local physical dynamics, and fresh water dynamics, along with seasonal upwellings determine the species that may occupy each location within the vast coastal and island nursery systems. This provides unique opportunities to compare and contrast physical environments and species interactions. The adjacent regional faunas of the Indian coast along the eastern Arabian Sea and the western Bay of Bengal are very little alike (Sharp 1988), indicating the very distinctive properties of the two physical environments and subsequent ecologies. The remaining living resources are associated with estuaries, river mouths and submerged freshwater seeps, each milieu offering refuge from anoxic upwelled waters.

Much of the coastal upwelling around the northern Indian Ocean derives from Southern Ocean midwater sources. The low-oxygen waters that upwell around the west Irian Jaya coast during wind forcing suggest the occurrence of low oxygen at

depth throughout the Indonesian Archipelago, and eastward, all the way to the Pacific Ocean. As described by Bianchi (this vol.), the northwest coast of Sumatra does not exhibit as shoal an oxygen minimum layer as do many of the areas within the Indonesian Archipelago, or around the northern coasts of the Bay of Bengal. This is due to the well mixed upper ocean, and limited primary production across the shelf.

Above the strong thermocline, the temperatures we find are relatively uniformly warm SSTs, ranging between 28 and 29°C around the edges of the warm pool, and between 29 and 30°C within. The shoal warm upper ocean and sharp oxycline suggest that only opportunistic, i.e., short-lived and fecund species will thrive (Dalzell and Pauly 1989; Pauly et al., this vol.). Highly mobile species such as the large tunas and a few related larger predator groups are only abundant at small, younger stages, and for brief periods, as the larger and older stages are tied to cooler oxygenated habitats by their unique thermobiology and respiratory requirements (Sharp and Dizon 1978; Sharp 1979). The few resident scombrid species are all very small (e.g., *Rastrelliger* spp.), or exhibit clear signs of thermal stress, such as early onset of reproduction, and smaller maximum sizes compared to conspecifics from cooler, more oxygenated regions of the adjacent tropical oceans. The food web tends to be broad, but relatively short (Pauly and Christensen 1993, this vol.), due to the intense competition for resources.

Year to year climate-driven ocean variability at local and regional scales is clearly the dominant forcing that affects local fisheries. Venema (this vol.) has documented observations from which it can be inferred that the small-scale structures that oceanographers are prone to removing from their atlas descriptions (e.g., small eddies and current fields) are important. Because of the strong wind forcing, and topographic complexities of the region, eddies and countercurrents should dominate the upper ocean dynamics.

The characteristic situation within the Indonesian Archipelago is further complicated by the dominance of relatively shoal topography, dominated by either coral reef or mangrove ecosystems. Both are affected by the strong vertical stratification set up by monsoonal rainfall and large seasonal inflow of rivers from Asia and the Irian Jaya-Papua New Guinea highlands (Milliman and Meade 1983; Wright 1987).

Conclusion

The dynamic Indonesian Archipelago provides a unique challenge to science. The impact of this domain on global climate is not questioned, it is only the magnitudes of the various competing forces that are yet to be determined.

The ecological dynamics reflect the climate dynamics. What is needed are better observations of the region's ecological and environmental properties before any grand conclusions might be drawn. The region comprises a mosaic of seasonally varying production and system dynamics, all of

which interact to generate ecological enigmas. One of these is how to estimate potential yields that are sustainable, in the context provided by myriads of species, embedded in a complex, climate-driven maze of environmental processes and interactions. What little we do know derives primarily from areas in which fishing effort has been very intense for many decades. In other areas less is known, generally due to the absence of human concentrations and activities (Dalzell and Pauly 1989).

As described by Lohmeyer (this vol.), performing pelagic fish surveys using trawling gear within the shoal, warm Indo-Pacific is nearly fruitless. Although numerous schools of surface swimming skipjack (*Katsuwonus pelamis*), longtail tuna (*Thunnus tonggol*), *Scomberomorus* spp., *Rastrelliger* spp. and frigate mackerels (*Auxis* spp.) are often observed, acoustic survey technologies are somewhat insensitive to many of these species either because they lack a gas bladder, or form small schools, and make poor acoustic targets (Venema, this vol.). Even the deepwater crustaceans and other benthic species are likely limited by seasonal oxygen levels, hence cryptic, suggesting that oxygen-rich freshwater seeps or other local phenomena might provide refugia for these species.

In the deeper channels, and open ocean around and about the Indo-Pacific region, wherever oxygen values exceed 1 ml·l⁻¹, bigeye tuna (*Thunnus obesus*) and other large predators might be found, but these would certainly never turn up in pelagic trawl surveys. On the other hand, the extensive wide mud-bottomed shoals surrounding regions of seasonal production harbor significant quantities of fishes and other marine resources. However, the high ambient temperatures preclude these species from reaching large size, and compress their life histories, resulting in intensely competitive, high throughput ecosystems, with lower overall abundances and sustainable yields.

References

- Allan, R.J. and J.I. Pariwono. 1990. Ocean-atmosphere interactions in low latitude Australasia. *Int. J. Climatology* 10:145-178.
- Bauer, R. 1985. Functional Description Master Oceanographic Observation Data Set (MOODS). Report submitted to Fleet Numerical Oceanography Center, Monterey, California. 79 p.
- Bauer, R. and M. Robinson. 1985. Description of the Bauer-Robinson numerical atlas. Version VIII. Compass Systems, Inc., San Diego, California. 13 p.
- Dalzell, P. and D. Pauly. 1989. Assessment of the fish resources of Southeast Asia, with emphasis on the Banda and Arafura Seas. *Neth. J. Sea Res.* 24(4):641-650.
- Duinker, J.C. 1989. River input into ocean systems, Theme 5 of the Snellius II expedition. *Neth. J. Sea Res.* 23(4):353-357.
- Hackert, E.C. and S. Hastenrath. 1986. Mechanisms of Java rainfall anomalies. *Mon. Weather Rev.* 114: 745-757.
- Hastenrath, S. and P.J. Lamb. 1979a. Climatic atlas of the Indian Ocean Part 1. Surface circulation and climate. University of Wisconsin Press. 109 p.
- Hastenrath, S. and P.J. Lamb. 1979b. Climatic atlas of the Indian Ocean Part 2. Heat budget. University of Wisconsin Press. 104 p.
- Hastenrath, S. and M.C. Wu. 1982. Oscillations of upper-air circulation anomalies in the surface climate of the tropics. *Arch. Geophys. Bioklimatol. Ser. B.* 31:1-37.

- Katili J.A. 1989. Review of past and present geotectonic concepts of eastern Indonesia. *Neth. J. Sea Res.* 24(2/3):103-129.
- Longhurst, A.R. and D. Pauly. 1987. *Ecology of tropical oceans*. Academic Press, San Diego, California.
- Milliman, J.D. and R.H. Meade. 1983. World-wide delivery of river sediment to the oceans. *J. Geol.* 91(1):1-21.
- Nichols, N. 1984a. El Niño southern oscillation and North Australian sea surface temperature. *Trop. Ocean Atmos. Newsl.* 24:11-12.
- Nichols, N. 1984b. El Niño southern oscillation and Indonesian sea surface temperature. *Mon. Weather Rev.* 112:424-432.
- Pauly, D. and V. Christensen. 1993. Stratified models of large marine ecosystems: a general approach and an application to the South China Sea, p. 148-174. *In* K. Sherman, L.M. Alexander and B.D. Gold (eds.) *Large marine ecosystems: stress mitigation and sustainability*. AAAS Press (American Association for the Advancement of Science Publishing Division), Washington, DC.
- Postma, H. and W.G. Mook. 1988. The transport of water through the east Indonesian deep-sea basins. A comparison of Snellius-I and -II results. *Neth. J. Sea Res.* 22(4):373-381.
- Shackleton, L.Y. 1987. A comparative study of fossil fish scales from three upwelling regions. *S. Afr. J. Mar. Sci.* 5:79-84.
- Sharp, G.D. 1979. Areas of potentially successful exploitation of tunas in the Indian Ocean with emphasis on surface methods. *Tech. Rep. IOFC/DEV/79/47*, 50 p. Indian Ocean Programme, FAO, Rome.
- Sharp, G.D. 1988. Neritic systems and fisheries: their perturbations, natural and man induced, p. 155-202. *In* H. Postma and J.J. Zijlstra (eds.) *Ecosystems of the world: Part 27. Ecosystems of continental shelves*. Elsevier Scientific Publishing Company, Amsterdam-Oxford-New York.
- Sharp, G.D. and A.E. Dizon, Editors. 1978. *The physiological ecology of tunas*. Academic Press, San Francisco and New York.
- van Aken, H.M., J. Punjanan and S. Saimima. 1988. Physical aspects of the flushing of the east Indonesian basins. *Neth. J. Sea Res.* 22(4):315-339.
- Wright, L.D. 1989. Dispersal and deposition of river sediments in coastal seas: models from Asia and the tropics. *Neth. J. Sea Res.* 23(4):493-500.
- Wyrski, K. 1961. Naga report: scientific results of marine investigations of the South China Sea and the Gulf of Thailand, 1959-1961. Vol. 2. Scripps Institution of Oceanography, La Jolla, California.
- Wyrski, K. 1962. The upwelling in the region between Java and Australia during the southeast monsoon. *Aust. J. Mar. Freshwat. Res.* 13:217-225.
- Wyrski, K. 1971. *Oceanographic atlas of the International Indian Ocean Expedition*. National Science Foundation, Washington, DC. 542 p.
- Wyrski, K. 1987. Indonesian throughflow and the associated pressure gradient. *J. Geophys. Res.* 92(C12): 12941-12946.

Variability of Sea Surface Features in the Western Indonesian Archipelago: Inferences from the COADS Dataset

CLAUDE ROY

Centre ORSTOM de Brest
B.P. 70 - Plouzané, France

ROY, C. 1996. Variability of sea surface features in the Western Indonesian archipelago: inferences from the COADS dataset [*Keragaman sifat-sifat permukaan laut di kepulauan Indonesia bagian barat: diambil dari kumpulan data COADS*], p. 15-23. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

Following a brief review of their key properties (based mainly on K. Wyrki's Naga Report of 1961), sea surface features of the Western Indonesian archipelago are characterized using time series extracted from the Comprehensive Oceanographic and Atmospheric Dataset (COADS), and covering the period from 1950 to 1990.

Abstrak

Diawali dengan tinjauan singkat sifat-sifat pokok dari laut (terutama berdasarkan pengamatan Naga Report 1961 dari K. Wyrki), sifat-sifat permukaan laut kepulauan Indonesia bagian barat selanjutnya dianalisis berdasarkan data COADS (*Comprehensive Oceanographic and Atmospheric Dataset*) tahun 1950-1990.

Introduction

This account of the regional oceanography of Western Indonesia, presented here as background to the surveys documented in this volume, is meant to explain observed patterns of productivity (Fig. 1). This account explicitly builds on the comprehensive review of Wyrki (1961), from which three sections were adapted, other sources of information on the oceanography of the Southeast Asian region being scarce. The mean spatial structure and the seasonal variability of major surface climatic parameters may be found in the Indian Ocean atlas of Hastenrath and Lamb (1979a, 1979b). Sharp (this vol.) also presented an overview of important, large-scale oceanographical and meteorological patterns.

In the introduction to his report on the physical oceanography of the Southeast Asian Waters, Wyrki (1961) noted that "a considerable number of local effects and features had to be expected". Indeed, the Southeast Asian region has one of the most complex topographical structures on earth: large and small islands subdivide the region into different seas connected with each other by many passages and channels. The variety of the physical settings also generates complex biological systems where local features are important. Thus, "the region comprises a mosaic of seasonally varying

production and system dynamics, all of which interact to generate ecological enigma" (Sharp, this vol.). By comparison with other marine ecosystems (see, for example, Parrish et al. 1983 or Pauly and Tsukayama 1987), any kind of generalization remains hazardous.

After a review of some important characteristics of the atmospheric and marine climate of the area, the seasonal and interannual variability of selected surface parameters in six areas, distributed around the Indonesian archipelago, is presented.

The Atmospheric Setting: the Monsoon Regime (Modified from Wyrki 1961)

The monsoon wind regime is a tropical phenomenon: the result of the interaction between a high atmospheric pressure cell centered over the continent in the winter hemisphere and a low atmospheric pressure cell that develops in the summer hemisphere over the continent, as a prolongation of the equatorial low. Because of the relative stationarity of the pressure distribution, the winds are very steady, especially over the sea. An important characteristic of the area is the biannual signal of atmospheric forcing, related to the movement of the sun and of the equatorial low, which crosses the equator twice each year.




Fig. 1. Composite satellite-derived map of phytoplankton pigment concentrations - indicating primary production levels - for Southeast Asia (courtesy of NASA/GSFC and Dr. F. Chavez, MBARI).

[Gambar 1. Peta komposit konsentrasi phytoplankton berdasarkan beberapa pengamatan melalui satelit yang menunjukkan tingkatan produktivitas primer untuk Asia Tenggara (data NASA/GSFC dan Dr. F. Chavez, MBARI).]

The north monsoon starts in October and is fully developed in January. The monsoon flux passes the equator as a north wind; south of the equator, the wind - due to the Coriolis effect - turns eastward, where it becomes the northwest monsoon. In February, the equatorial low, previously located at 10°S, starts to move northward and comes to lie over Java and the Lesser Sunda Islands. The southeast flux extends to the north. Southwest winds dominate south of the equatorial low, between Java and Australia. North of the equator, the direction of the wind remains unchanged but its strength declines. Little change occurs in April, but a complete reorganization of the atmospheric fluxes is observed in May. The southeast atmospheric flux crosses the equator, then turns eastward. The northeast winds over the China Sea and the Philippines are replaced by the south monsoon, which prevails over the whole of Southeast Asia. The south monsoon reaches its full development between July and August: the Asian low

and the Australian high are fully developed and the north-south pressure gradient and atmospheric circulation are maximum. In September at the end of the boreal summer season, the Asian low starts to weaken and in October the equatorial low starts to move southwards. In the north, northeast winds start to dominate. In November, the equatorial low crosses the equator and the northeast monsoon intensifies. The southward reach of the northeast monsoon follows the migration of the equatorial low which attains its southernmost position in January.

This seasonal variability of the winds cause corresponding changes in surface currents. Following the direction of the monsoon flux, which changes twice a year, the currents also reverse themselves twice a year. This is perhaps the key ecological feature in the area (see e.g., Martosubroto, this vol. and Venema, this vol.).

Some Characteristics of the Surface Circulation in the Southeast Asian Waters (Adapted from Wyrтки 1961)

Some topographical features of Southeast Asia favor the development of a strong surface circulation: the area formed by the South China Sea, the straits between Sumatra and Borneo, the Java Sea, the Flores Sea and the Banda Sea which has its main axis aligned with the wind flux during both monsoons; this, along with the relative constancy of the winds favors the development of surface circulation patterns strongly connected to the wind regime. In other parts of the region, however, it is difficult to extract any large-scale and coherent circulation pattern; local effect and intermittency appear to be dominant. Water exchange with the Pacific Ocean occurs through the Molucca Strait, the Philippines, and the Sulu Sea (Sharp, this vol.). The Makassar Strait has usually a current directed to the south, from the Pacific to the Indian Ocean. However, the water exchange through the Malacca and the Sunda Straits is small, even when the currents are strong. In the Java Sea, the surface water flow is directed to the west from May to September and to the east from November to March (Martosubroto, this vol.). In April and October, when the direction of the flow changes, eddies are generated along with a shear between the eastward current off the coast of Java and the westward current off the coast of Borneo. Through the Malacca Strait and the Sunda Strait, the surface currents are generally directed towards the Indian Ocean and are strongly related to the sea level gradient through the straits. The flow through the Sunda Strait reaches its maximum in August, during the southeast monsoon and there is a second maximum in December/January. In the Malacca Strait, the period of strongest flow is from January to April, during the northeast monsoon.

Properties of the Surface Waters (Modified from Wyrтки 1961)

High sea surface temperature (SST $>25^{\circ}\text{C}$) and small seasonal amplitude ($<3^{\circ}\text{C}$) are the dominant characteristics of Southeast Asian waters; moreover, their spatial distribution is quite uniform, with small gradient over the entire region.

The high rainfall, which largely exceeds evaporation, causes an average salinity of less than 34‰. This rainfall, the river runoff it causes and the archipelagic nature of the area are responsible for an extremely variable spatial distribution of the surface salinity. The alternance of the monsoons leads to rainy and dry seasons, and thus to large environmental variations. Rivers runoff, notably into the Java Sea, rather than rainfall is the cause of the low coastal salinities, even far offshore. The largest extent of the low salinity waters occurs in April and May when, with the onset of the southeast monsoon, they are transported from the Java Sea into the southern China Sea. In June, water with a higher salinity ($>32\text{‰}$) enters from the east into the Java Sea and, thence farther north up to the

southern China Sea, reaching its maximal westward penetration in September. With the onset of the northwest monsoon, in October/November, these water masses are pushed back again towards the Java Sea, while their salinity is reduced by the start of the rain. Salinity in the Java Sea drops below 32‰, reaching its minimum in May, when river runoff from Borneo is maximal.

A steady southeast current flows from the Sunda Shelf through the Malacca Strait into the Indian Ocean. During the northeast monsoon, this current transports relatively high salinity water from the South China Sea. During the southeast monsoon, the water transported is of low salinity, due to river runoff from Central Sumatra and direct rainfall. Strong tidal currents cause a complete vertical mixing over the water column.

Inferences from the COADS Dataset

Extraction and presentation of the data extracted from the Comprehensive Oceanographic and Atmospheric Dataset (COADS) dataset (Woodruff et al. 1987), recently published as a set of five CD-ROMs through the Climate and Eastern Ocean Systems (CEOS) project (Bakun et al. 1992), were used to document the seasonal and interannual variations of SST, of scalar wind speed and the north-south and east-west component of the pseudo wind-stress in six selected areas defined on Fig. 2. The COADS database contains the surface weather observations collected by merchant ships and other platforms (buoys, weather stations, lightships, etc.) since 1854. The data distribution and density in the area are presented in Fig. 3. Data density was, in the 1950s, low all over the region, except along trade lanes such as that passing through the Malacca Strait to Singapore. Later, data density increases along the eastern part of the China Sea but remains "spotty" over the western part of Southeast Asia.

For the purpose of this study, the Java Sea is separated in two areas, eastern and western (see Fig. 2). Other areas defined in Fig. 2 are the southern part of the China Sea, the northern and central parts of the Strait of Malacca, and the Sunda Strait.

For each parameter and each area, a time-series of monthly mean values from 1950 to 1990 was built using the individual observations extracted from the COADS database. A mean annual cycle was derived from the monthly time-series. A time-series of the mean annual value was then calculated and used to characterize the interannual variability from 1950 to 1990.

Variability of the Sea Surface Temperature

SSTs are high all year round in the six areas, with minimum values (27.5°C) observed in January and February, in the southern part of the China Sea (Fig. 4). Maximum values are comprised between 29.2°C (Sunda Strait) and 29.8°C




Fig. 2. Definitions of the six areas of Western Indonesia used to structure this contribution (clockwise from the upper left corner): (i) northern Malacca Strait; (ii) central Malacca Strait; (iii) southern South China Sea (including Karimata Strait); (iv) eastern Java Sea; (v) western Java Sea; and (vi) Sunda Strait.

[Gambar 2. Pembagian enam daerah Indonesia bagian barat sebagai dasar penyusunan tulisan ini (searah jarum jam dari pojok kiri atas): (i) Selat Malaka bagian utara; (ii) Selat Malaka bagian tengah; (iii) Laut Cina Selatan bagian selatan (termasuk Selat Karimata); (iv) Laut Jawa bagian timur; (v) Laut Jawa bagian barat; dan (vi) Selat Sunda.]

(Malacca Strait). In some areas located on or south of the equator (Sunda Strait, Java Sea and southern part of the South China Sea), there is a pronounced biannual cycle with a first SST minimum in January-February and a second in August-September.

The amplitude of the SST interannual variability is less than 1.0°C in the Malacca Strait but it increases toward the south (Fig. 5). The greatest amplitude is recorded in the Sunda Strait where it almost reaches 1.5°C. The eastern and western Java Sea, the southern tip of the China Sea and the Sunda Strait all exhibit a similar interannual variability, with mean annual SST values above the average at the end of the 1950s (a feature that may be associated with the 1957-1958 El Niño Southern Oscillation [ENSO] event). SSTs below average appear to have occurred both during the mid-1960s and 1970s. The 1980s, on the other hand, has higher than average SSTs. Major ENSO events (1957-1958; 1972-1973) are associated with a relative peak in SST except for the 1983-1984 event which is nevertheless considered as the most intense of this last century.

Variability of the Wind

The monthly seasonal cycles of the scalar wind speed (i.e., the mean of the two wind




Fig. 3. Data density in the COADS (by selected years: 1950, 1960, 1970, 1980 and 1990). Note that data density is high only along commercial routes and very scarce in the open ocean.

[Gambar 3. Kepadatan data COADS (menurut tahun: 1950, 1960, 1970, 1980 dan 1990). Perlu disimak bahwa kepadatan densitas data tertinggi terdapat di sepanjang daerah pelayaran niaga dan sangat sedikit terdapat di laut bebas.]




Fig. 4. Mean monthly cycle of SST in six areas of Western Indonesia.
 [Gambar 4. Siklus bulanan rata-rata SST di enam daerah perairan Indonesia bagian barat.]




Fig. 5. Mean annual values of SST from 1950 to 1990 in six areas of Western Indonesia (COADS dataset).
 [Gambar 5. Rata-rata tahunan SST dari tahun 1950 hingga 1990 di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 6. Mean monthly cycle of the scalar wind speed in six areas of Western Indonesia (COADS dataset).
 Gambar 6. Siklus bulanan rata-rata dari kecepatan angin di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 7. Mean monthly cycle of the north-south pseudo wind stress component in six areas of Western Indonesia (COADS dataset).
 [Gambar 7. Siklus bulanan rata-rata dari komponen pengaruh angin pseudo utara-selatan di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 8. Mean monthly cycle of the east-west pseudo wind stress component in six areas of Western Indonesia (COADS dataset).
 [Gambar 8. Siklus bulanan rata-rata dari komponen pengaruh angin pseudo timur-barat di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 9. Mean annual values of the scalar wind speed from 1950 to 1990 in six areas of Western Indonesia (COADS dataset).
 [Gambar 9. Nilai rata-rata tahunan kecepatan angin dari tahun 1950 hingga 1990 di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 10. Mean annual values of the north-south pseudo wind stress component from 1950 to 1990 in six areas of Western Indonesia (COADS dataset).

[Gambar 10. Nilai rata-rata tahunan komponen pengaruh angin pseudo utara-selatan dari tahun 1950 hingga 1990 di enam daerah perairan Indonesia bagian barat (data COADS).]




Fig. 11. Mean annual values of the east-west pseudo wind stress component from 1950 to 1990 in six areas of Western Indonesia (COADS dataset).

[Gambar 11. Nilai rata-rata tahunan komponen pengaruh angin pseudo timur-barat dari tahun 1950 hingga 1990 di keenam daerah perairan Indonesia bagian barat (data COADS).]

components) and of the two components of the pseudo wind stress (i.e., squares of the north-south and east-west wind components) are presented in Figs. 6, 7 and 8, respectively. The minimum values are observed in the southern part of the Strait of Malacca (3.5 ms^{-1}); maximum values occur in the Sunda Strait. A marked biannual cycle, due to the monsoon, appears in the southern part of the China Sea, in the Java Sea and in the Sunda Strait areas. Maximum values occur in January and August, while minimum values, in April and November-December. The maximum values stay below 6 ms^{-1} except in the Sunda Strait. This suggests that biological processes may not be dominated by hydrodynamic factors related to the wind (Therriault and Platt 1981; Cury and Roy 1989). The seasonal behaviour of the two wind stress components clearly illustrates the strong alternation (and reversal) of the wind regime due the dynamics of the monsoons (Figs. 7 and 8).

The interannual variability of the wind is rather small in the northern and central Malacca Strait (0.5 ms^{-1}) but increases toward the south (Figs. 9, 10 and 11). The mean annual scalar wind speed exhibits in almost all areas a positive long-term trend. Except for this trend, no clear pattern of variability is readily identifiable: ENSO events do not appear to affect local wind variability. Also, the interannual variability of the two components of the pseudo wind stress exhibits a behavior similar to the variability of the scalar wind.

Conclusion

The previous considerations lead one to conclude that the marine habitats of the adjacent areas to the Indonesian Archipelago are quite unique in the world: the imbrication of land and sea creates complex systems where local processes may prevail over global dynamics. Also, the monsoon regime creates such a strong seasonality of the

characteristic of the environment that the alternation of the north and south winds completely reorganizes the surface circulation; this can be expected to have a strong ecological impact. Interannual variability exists, but surprisingly, it appears not to be closely associated with ENSO events - at least, no strong anomalies in either SST or wind appear in the COADS dataset that can straightforwardly be linked with ENSO events. This begs the question whether the complexity of the Southeast Asian environment may have led to some sort of homeostasis.

References

- Bakun, A., V. Christensen, C. Curtis, P. Cury, M.H. Durand, D. Husby, R. Mendelsohn, J. Mendo, R. Parrish, D. Pauly and C. Roy. 1992. The Climate and Eastern Ocean Systems project. *Naga, ICLARM Q.* 15(4): 26-30.
- Cury, P. and C. Roy. 1989. Optimal environmental window and pelagic fish recruitment success in upwelling areas. *Can. J. Fish. Aquat. Sci.* 46(4): 670-680.
- Hastenrath, S. and P.J. Lamb. 1979a. Climatic atlas of the Indian Ocean. Part 1. Surface circulation and climate. University of Wisconsin Press. 109 p.
- Hastenrath, S. and P.J. Lamb. 1979b. Climatic atlas of the Indian Ocean. Part 2. Heat budget. University of Wisconsin Press. 104 p.
- Parrish, R.H., A. Bakun, D.M. Husby and C.S. Nelson. 1983. Comparative climatology of selected environmental processes in relation to eastern boundary current pelagic fish reproduction. *In* G.D. Sharp and J. Csirke (eds.) Proceedings of the Expert Consultation to Examine Changes in Abundance and Species Composition of Neritic Fish Resources. FAO Fish. Rep. 291(3): 731-777.
- Pauly, D. and I. Tsukayama, Editors. 1987. The Peruvian anchoveta and its upwelling ecosystem: three decades of change. *ICLARM Stud. Rev.* 15, 351 p.
- Therriault, J.C. and T. Platt. 1981. Environmental control of phytoplankton patchiness. *J. Fish. Res. Board Can.* 38(6): 368-641.
- Woodruff, S.D., R.J. Slutz, R.L. Jenne and P.M. Steurer. 1987. A Comprehensive Ocean-Atmosphere Data Set. *Bull. Am. Meteorol. Soc.* 68(10): 1239-1250.
- Wyrtki, K. 1961. Naga report: scientific results of marine investigations of the South China Sea and the Gulf of Thailand, 1959-1961. Vol. 2. Scripps Institution of Oceanography, La Jolla, California.

The Marine Fisheries of the Western Archipelago: Towards an Economic History, 1850 to the 1960s

JOHN BUTCHER

Faculty of Asian and International Studies
Griffith University
Queensland 4111, Australia

BU

Q1561
Xref Q1107
Marine fisheries
Fisheries; Shrimp
Aquaculture; Fishing
operations
Historical account
Isew, Malaysia
Isew, Indonesia

the Western Archipelago: towards an economic history, 1850 to the 1960s [Perikanan laut di kawasan sejarah ekonomi, dari tahun 1850 hingga tahun 1960an], p. 24-39. In D. Pauly and P. of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

This chapter describes the changes that took place in the marine fisheries of the Western Archipelago between about 1850 and the 1960s. About 1850, most fishing took place close to shore; passive gears such as fishing stakes were predominant in many areas; a large proportion of fish was dried or salted; and fishing oriented to production for the market occurred only in areas near large concentrations of population. In the late 1800s, fishing underwent a commercial revolution, as demand for fish grew both because of the growth of cities and the rapid expansion of mining and plantation agriculture and because of improvements in transportation. Along both sides of the Straits of Malacca, Chinese, employing fishing methods long used by Malays, responded to this demand on a large scale, particularly at Bagan Si Api Api, where the holder of the salt monopoly leased from the Netherlands Indies government provided fishers with relatively cheap salt. At the same time, the fishing communities of Trengganu, making use of cheap salt from Siam, greatly increased production. In contrast, in Java, where salt provided by the government's own monopoly was far more expensive, there was little if any increase in production. Instead, Java became a large importer of fish not only from Bagan Si Api Api but also from Siam. To increase production, the colonial authorities in both Malaya and the Netherlands Indies experimented with large trawlers but with no success. In the meantime, Japanese, working with motorized fish carriers and employing a new method of capturing reef fish, came to dominate the fish markets of Batavia and Singapore, while Chinese operating purse seiners from motor vessels greatly increased production in Malaya. In the aftermath of World War II, the primary concern of all governments was to restore production quickly in order to meet severe food shortages. Motorization and the adoption of nets made of synthetic fibers took place first in Malaya and then in Indonesia. In 1965, a few fishers in Malaysia, following the lead set by Thailand, began fishing with otter trawls in the Straits of Malacca, and soon this fishing method spread to Sumatra and then to Java.

Abstrak

Tulisan ini menggambarkan perubahan-perubahan yang terjadi pada sektor perikanan laut di kawasan kepulauan bagian barat antara tahun 1850 dan 1960an. Sekitar tahun 1850, penangkapan pada umumnya dilaksanakan di daerah dekat pantai; alat-alat tangkap pasif seperti jermal mendominasi banyak daerah; sebagian besar ikan dikeringkan atau digarami; dan kegiatan penangkapan bertujuan untuk memasok pasar yang dekat dari daerah pemukiman penduduk. Pada akhir tahun 1800an, perkembangan penangkapan ikan mengalami perubahan pesat sejalan dengan situasi pasar, dimana permintaan akan ikan meningkat akibat pesatnya pertumbuhan perkotaan, perkembangan kegiatan pertambangan dan perkebunan; disamping juga karena kemajuan transportasi. Sepanjang wilayah pantai di kedua sisi Selat Malaka, para nelayan keturunan Cina dengan menggunakan cara penangkapan ikan yang sudah biasa dilakukan oleh para nelayan pribumi Malaya, berusaha memenuhi permintaan situasi pasar dalam skala besar, khususnya di daerah Bagan Si Api-Api, dimana pemegang monopoli garam dari pemerintah Hindia Belanda memberikan harga garam relatif murah kepada para nelayan. Pada saat yang bersamaan, masyarakat nelayan di Trengganu, menggunakan garam yang murah dari Siam, yang mana ini merangsang percepatan produksi ikan. Sebaliknya di pulau Jawa, dimana garam dimonopoli oleh pemerintah harganya sangat mahal, sehingga tidak merangsang kenaikan produksi. Oleh karena itu Jawa menjadi importir ikan terbesar tidak hanya dari Bagan Si Api-Api tetapi juga dari Siam. Dalam rangka meningkatkan produksi, pemerintah kolonial di Malaya dan Hindia Belanda mengadakan percobaan dengan trawl yang besar tetapi tidak berhasil. Sementara itu, orang-orang Jepang yang dilengkapi dengan kapal-kapal penangkap ikan yang menggunakan motor serta dilengkapi dengan alat tangkap baru untuk ikan karang mendominasi pasar di Batavia dan Singapura, sedangkan para nelayan Cina dengan pukot cincin (purse seine) dan kapal motornya benar-benar meningkatkan produksi ikan di Malaya. Setelah Perang Dunia II, yang menjadi perhatian utama pihak-pihak pemerintahan adalah berupaya menjaga produksi guna memenuhi kurangnya bahan pangan. Motorisasi dan pemakaian jaring dari bahan serat sintesis dimulai di Malaya dan selanjutnya berkembang di Indonesia. Pada tahun 1965, beberapa nelayan dari Malaysia, mengikuti gerak nelayan Thailand, mulai menangkap ikan dengan otter trawl di Selat Malaka yang kemudian menyebar ke seluruh perairan Sumatra dan Jawa.

Introduction

The seas of Southeast Asia have long provided coastal dwellers with a multitude of riches ranging from fish, shrimps and whales to pearl oysters, *tripang* and seaweed. People have

collected and captured marine life for food, medicine, oil, jewelry, and a great variety of other uses. The purpose of this chapter is to trace and explain the changes that took place in fishing in the Western Archipelago from about 1850 to the mid-1960s,

when trawling was rapidly adopted as one of the main fishing methods. The focus will be on the capture of finfish, but some attention will have to be devoted to that of shrimps as well, for there has often been a close relationship between the two fisheries. It is useful to begin this survey in the middle of the nineteenth century with an overview of fishing in what I shall call the "Western Archipelago" corresponding to what are now Malaysia and Western Indonesia (Fig. 1). We must keep in mind though that such an approach carries the danger of ignoring important changes such as the development of fishing techniques that had taken place over many centuries. Box 1 defines some of the local terms used here.

Economic History

Fishing about 1850

In 1850, the Western Archipelago had a population of about twenty million, concentrated in the islands of Java and Madura, the highlands of west Sumatra, the Straits Settlements, numerous small harbor towns along the coasts of Sumatra,

the Malay Peninsula, Borneo, and in isolated mines and plantations. For virtually all of these people - whether the overwhelming majority who produced at least part of their own food or those few who lived in towns or worked in mines and plantations here and there - fish was a staple food. "Fish along with rice is the main ingredient of [the Javan's] meal" (Anon. 1882). Nearly always fish was consumed after it had undergone some form of preservation, almost always by drying, salting, or some combination of the two, for there was no other way of preserving fish in the hot, humid climate. Dried fish was, observed Crawford (1820), "an article of as universal consumption among Indian islanders as flesh is in cold countries." Even fishers consumed at least part of their fish intake in this form, since for most of them there were times of the year when there were few fish or when the weather prevented them from catching them. Also important in the diet of the people of the Western Archipelago was shrimp or fish paste, called *terasi* by the Javanese and known as *belacan* to the Malay-speaking peoples of the archipelago, "the universal sauce of the Indian islanders", without which "no food is deemed palatable" (Crawford 1820).




Fig. 1. Map of the Western Archipelago, showing all locations mentioned in the text.

[Gambar 1. Peta kepulauan bagian barat yang menunjukkan beberapa lokasi penting sebagaimana disebutkan dalam tulisan.]

Box 1. Definition of some non-English terms used in the text.

[Boks 1. Penjelasan beberapa istilah lokal yang dipakai di dalam tulisan ini.]

<i>ambai</i>	- fishing stake consisting of two converging wings of poles which guide shrimp into a trap made of a fine-meshed net at the end of which is placed a pocket of coarse sacking.
<i>belacan</i>	- fish or (more usually) shrimp paste.
<i>bubu</i>	- generic term for a variety of portable fish traps set on the seabed.
<i>jermal</i>	- large fishing stake consisting of two converging wings of poles and an enclosure into which fish and shrimps are swept by the current and from which they can be extracted by lifting a large rattan screen.
<i>kongsi</i>	- a business group.
<i>muro ami</i>	- drive-in net set on reef slopes and used for capturing fusiliers (Fam. Caesionidae) and other reef fish.
<i>payang</i>	- seine net used in conjunction with a fish lure in the <i>layang</i> fishery of Java Sea.
<i>prahu mayang</i>	- offshore fishing boat, usually operating a <i>payang</i> to catch <i>layang</i> and other pelagic species.
<i>rumpon</i>	- floating fish lure used in catching <i>layang</i> .
<i>sero</i>	- fishing stake made up of a series of successively smaller chambers through which fish move until captured in the smallest chamber.
<i>si stji</i>	- fishing stake similar to <i>jermal</i> but much smaller and easily moved from one place to another.
<i>tendak</i>	- fish lure.
<i>terasi</i>	- Javanese word for <i>belacan</i> .
<i>towkay</i>	- Chinese businessman.
<i>towkay bangliau</i>	- the head of a group of (Chinese) fishers, as at Bagan Si Api Api.

The importance of fish to nutrition took many forms. For most people, fish was almost their only source of animal protein, for meat was eaten only on special occasions. It also supplied them with calcium (for tiny dried fish were eaten whole, i.e., their skeleton was consumed as well), iodine and various other essential nutrients. But most of the inhabitants of the Western Archipelago consumed fish in fairly small quantities, or at least that is the impression given by numerous references to what people ate at this time, as in this description of a meal in a Malay house in Malacca:

Cooking operations are simple, for the meal usually consists of boiled rice, small pieces of dried fish heated over the embers of the fire, and a concoction of hot red chillies that have been ground with salt into a paste. The smoking rice is put in the centre of the floor; pieces of dried fish and fiery chillies, ground up with salt are the usual relishes, and around this simple fare the family sit with their legs crossed (Rathborne 1898).

And in this more general report from the 1880s: Where meat is almost unobtainable, or if obtained is coarse and uneatable, the dried salt fish is the only article of food to be relied upon, and, so far as my experience goes, it is both palatable and nourishing. It is soaked and cut up into small dice, and fried until quite brown. *A small quantity of this mixed with boiled rice* makes a dish, which Chinese,

Malays and Europeans seem equally to relish (Tenison-Woods 1888; emphasis added).

It appears that the consumption of fish per capita was particularly low in parts of Java. There, *terasi* made an important contribution to people's diets not because it is rich in protein, for it was consumed in tiny quantities, but because its use as a condiment stimulated the consumption of unpolished rice and soya products, their main sources of protein.

The fish and shrimp people ate came from rivers, lakes, paddy fields and irrigation canals, fresh- and brackishwater ponds specially constructed for the cultivation of fish, and most important of all, the sea. It is probably safe to say that fish and shrimp captured from the sea made up a slightly smaller proportion of the total intake in the middle of the nineteenth century than they did in the early 1900s, by which time many inland waters had been polluted or silted up as a result of human activities and (as will be seen) sea fisheries had expanded greatly. Nevertheless, the sea was by far the biggest source of the fish and shrimp people consumed.

We can safely assume that the seas of the Western Archipelago were blessed with an abundance of fish at this time. Certainly that is the impression conveyed by contemporary reports. "No part of the world abounds in more fine fish", declared Crawford (1820). "The seas of the western parts of the Archipelago particularly the Strait of Malacca, and the shores of the Gulf of Siam, are the most remarkable for their abundance of edible fish." Along the west coast of the Malay Peninsula, wrote Anderson (1824), "fish of the choicest and most delicate description is extremely abundant," while along the east coast, according to Clifford (1897), "the fish crowd the shallow shoal waters, and move up and down the coast, during the whole of the open season, in great schools acres in extent." And an Englishman who sailed along the west coast of Borneo in the 1820s commented that "the coasts and rivers abound with excellent and wholesome fish in the greatest variety, and of the most delicious flavours" (Moor 1837). We have to treat these accounts with caution. European observers may have exaggerated the abundance of the marine life of the Western Archipelago just as they did the fertility of the soils. Nevertheless, reports about the abundance of fish in very specific locations certainly confirm the impression given by these accounts. At certain times in the Brouwerstraat (Selat Panjang), "the movement near the surface of a solid mass of fish, consisting almost entirely of spawners, produces a choppy rippling of the water" (Anon. 1882). Abdullah (1970) recalled that when the British arrived at Singapore in 1819 "fish were very plentiful and large ones were found close to shore." Moreover, it is worth noting that the "abundance" of fish at this time was determined mainly by the ability of people to catch them, not by the number actually present in the sea. Thus, no one could have had much of an appreciation of the abundance of demersal species, those pelagic species moving about far from shore, and those fish (notably fusilier, Fam. Caesionidae) living above reef slopes. There were, in short, many niches remaining to be explored and exploited.

Except along the south coast of Java and the west coast of Sumatra, where fishing was generally confined to sheltered bays because of rough seas and steep cliffs, people engaged in the work of catching and processing fish all along the coasts of the Western Archipelago. The extent to which they devoted their lives to fishing, however, varied greatly from one part of the archipelago to another. In some areas, fishing was the main livelihood for many people. Veth (1875) noted that all along the north coast of Java "one finds a number of kampongs whose people devote themselves almost exclusively to fishing" and that this was even more so in Madura, "for whose population fishing is a chief means of support." In communities such as these, almost all work was related in one way or another to fishing, the women often looking after processing and marketing while the men fished or repaired boats and nets. In some other areas fishing took place only during certain months of the year, as in Sarawak:

Large fishing establishments are found at the mouths of all the principal rivers during the southwest, or fine monsoon; the fishermen usually leaving them during the northeast, or boisterous monsoon, and returned to the town, where they pursue other avocations until fine weather again brings the shoals of fish to their shores (Low 1848).

In many other areas fishing was an intermittent activity, pursued when people needed fish for their own consumption or when they did not have to tend their crops, as in Sumatra, "where fishing is a separate occupation in only a very small part [of the island]" (Anon. 1882).

Here a striking pattern becomes evident, and that is that the places where fishing was people's primary livelihood tended to be quite near large concentrations of population, particularly cities and other places such as mines and plantations where there were large numbers of people who did not produce their own food. In short, as a rule, fishing villages engaged in production for nearby markets. Thus, the fishing villages of Madura and the north coast of Java produced fish mainly for the large towns and huge rural population of Java, while the Chinese fishers along the west coast of Borneo caught fish both for the nearby towns such as Pontianak and for the gold miners of the interior. This pattern was a reflection both of the nature of transportation at this time - slow and unreliable - and the fact that fish could generally be caught in sufficient quantities within a relatively short distance of markets. There were, of course, exceptions to this pattern. The most notable was the *trubuk* fishery of the Brouwerstraat, but it is worth noting here that it was the expensive roe, not the whole fish, that was carried to markets around the archipelago, just as *tripang* and sharks' fins were shipped to China and mother of pearl to China and Europe at this time.

The fishers of the Western Archipelago employed a great range of ingenious methods to capture fish. These included many different kinds of fishing stakes such as the *sero* and the *jermal*; a variety of traps, widely known as *bubu*; an enormous variety of nets operated from shore as well as from boats;

different techniques using hook and line; and many other methods such as harpooning and the use of stupeficients. Looked at from today's vantage point, it is notable that a very large proportion of the fish caught in the Western Archipelago at this time was caught by means of fishing stakes:

This way of fishing is so important that in some parts of Java it far surpasses net fishing, for example, in Meester Cornelis in the Residency of Batavia and in the part of Surabaya bordering the Madura Strait, where the amount of fish caught by any other means is in comparison completely insignificant (Anon. 1882).

The prevalence of such "passive" devices was in no way an indication that fishing methods were "undeveloped" at this time but rather that large quantities of fish could usually be caught without chasing after them.

One of the most striking features of fishing at this time was its rhythmic nature, oscillating according to the hour of the day, the phases of the moon and the seasons of the year, all of which had a bearing both on the presence of fish and on the ability of fishers to catch them. Of crucial importance for many fishers was the daily alternation of the land and sea breezes:

Nearly everyday one can see the *prahu mayang*... push off from shore at three or four in the morning in order to cast their nets. The land wind carries them quickly out of sight, but at noon they prepare to make use of the sea wind for their return trip, which brings them back to shore at two o'clock, often with a full load (Veth 1875).

Many forms of fishing were regulated by the phases of the moon, though with great variation from place to place depending both on the behavior of the fish and the method being used to catch them:

... in Tapanuli fishing at night takes place only during a dark moon, whereas, in contrast, at Tanjong Pura (East Coast of Sumatra) it takes place during the full moon (*pasang besar*) and the first quarter and last quarter (*pasang mati*), while in the Brouwerstraat fishing is considered the most favorable during the new and full moons (Anon. 1882).

Almost everywhere the rhythm of fishing had a seasonal nature, both because certain fish, notably pelagic species, were more abundant at certain times of the year and because during certain months the strong monsoon winds made fishing too dangerous. Thus, during the three or four months when the northeast monsoon made fishing impossible, the fishers of Trengganu "build and repair their boats and houses, make and mend their nets, do a little planting, and generally pass the time in performing odd jobs" (Clifford 1895).

Despite the existence of these regularities, it is essential to keep in mind that uncertainty and risk were always a part of fishing. Migratory fish did not always arrive at the time and place they normally did. If the sea breeze did not start at the

expected time the morning's catch would spoil before the *prahu* could reach shore. A day's catch could be lost if the boat overturned in the breakers. A sudden storm might capsize a boat and drown all on board. Even operating a fishing stake exposed those who worked on it to "dangers from stray sharks, sawfish and crocodiles, from the deadly sea-snakes, from many kinds of medusae, from fish with venomous fins, from stinging-rays and torpedo-fish" (Wilkinson et al. 1904). These uncertainties and dangers may explain why taboos and rituals were an important part of fishing.

Mention must be made of the role of states at this time. Most fishing probably took place without the help or hindrance of state powers. In fact, one of the dangers faced by fishers (and discouraging people from going to sea) was the absence of control over pirates and slave raiders. Nevertheless, in some places state authorities did impose taxes and monopolies that affected the livelihoods of fishers. The sultan of Siak demanded that the *trubuk* fishers hand over first part and later all of their catch at prices well below market values, while the Dutch imposed licence fees when they assumed control over his domain. And in Java the Dutch subjected fishing to various taxes as well as a government monopoly over the production and sale of salt, which may be why in Java so little use of salt was made when drying fish (Veth 1875). In light of present-day debates about regulating access to the sea it is worth referring to a report from the 1920s that "the sea next to the north coast of Java is mutually divided by the villages into districts, where each has erected its own *tendak*. Customary law [*adaf*] has thus created a right of ownership" (Schippers 1928). It is not clear to me, however, whether this was an ancient practice or one that developed as the number of fishers increased during the nineteenth and early twentieth centuries, nor is it clear what role if any state authorities had in promoting or supporting such arrangements.

An issue of central importance about which I have little information concerns the financing of fishing at this time. A survey of sea fishing in the Netherlands Indies published in 1882 makes it clear that fishing stakes, boats and nets often cost a great deal to buy or construct, but it gives little idea of how this money was raised. With respect to Java, the survey often refers to the "owners" of boats and gear, indicating that the owner almost invariably captained the fishing team, and describes the often elaborate ways the proceeds from catches were distributed among owners and crews, but it gives no hint of whether owners used their own funds (perhaps the accumulated profits of earlier fishing) to buy boats and gears or whether they borrowed money to buy them (Anon. 1882). It is only when Chinese fish traders were involved that the survey becomes more specific, as in this comment about the *trubuk* fishery of the Brouwerstraat:

... the trade is entirely in the hands of foreigners, namely, the Chinese buyers, who often have the fishers completely in their control by giving them advances, completely regulate the market price of the product, and

take the biggest profit. The increase in price that *telor trubuk* [*trubukroe*] undergoes makes it often unobtainable for the ordinary native and more of a special dish than a daily dessert (Anon. 1882).

The fact that the *trubuk* fishery produced a high cost product for long distance trade may mean that this situation was far from typical, but this needs to be explored.

The Commercial Revolution

Beginning about 1870, the production of fish products began to increase dramatically in several parts of the Western Archipelago. We must see this change in the context of the political and economic transformation taking place at this time. In the 1850s, the Dutch began incorporating (by force in the case of Aceh, by treaty in many other places) more and more of the region into the Netherlands Indies until, by the early twentieth century, it covered the areas closely corresponding to present-day Indonesia, while the British extended their control over the Malay Peninsula in stages from 1874 to 1910. This extension of colonial authority was accompanied by (and financed by) a rapid expansion in economic activity, as first Chinese and then European entrepreneurs opened up mines and plantations, particularly in the Malay Peninsula and Sumatra. In Java, there was a big expansion in sugar production, and the population of the island continued to grow. Also significant was the growth of Singapore, Penang, Batavia, Surabaya, Semarang, Medan and many smaller towns. These changes had a profound impact on fisheries in the Western Archipelago. The market for fish skyrocketed both because there was an enormous increase in the number of people who now had to buy fish (or had it bought for them, in the case of mine and plantation workers) and because steamships could now transport fish products (mainly salted fish and *belacan*) fairly cheaply over long distances. What took place is best described as a commercial revolution because of the vast increase in production for the market and because of the development of an elaborate system of financing, processing, and marketing. It certainly was not a technological revolution, since little change took place in fishing techniques. What occurred was a spectacular expansion of fishing using existing methods.

Although fishers throughout the Western Archipelago took part in this expansion, the degree to which they responded to the new opportunities varied considerably. We can demonstrate this by looking at three areas: the Strait of Malacca, Trengganu and Java.

The most rapid expansion took place along the coasts of the Straits of Malacca. Beginning about 1860, small groups of Chinese (referred to as former pirates in some sources) began settling at many spots along the straits. Along the eastern side of the straits, Chinese set up fishing villages at Kuala Kurau, Pangkor and Pulau Ketam. Along the western side, the most important area was the estuary of the Rokan River, where

Chinese founded the village of Bagan Si Api Api. All of these fishers adopted Malay fishing techniques; those fishing along the eastern side supplied fish to Penang and the rapidly growing mining population of the interior, while those at Bagan Si Api Api may have shipped what they produced across the Straits to Malacca as well as possibly up the Rokan River. At first no ruler or government exerted much authority over these villages, but by the 1880s the colonial states had exerted enough control to force them to pay taxes.

The most spectacular growth in production took place in the Rokan estuary, a place which was extraordinarily rich in marine life but which was not even mentioned in the 1882 survey of fishing in the Netherlands Indies. In 1898 (the first year for which there is a figure), Bagan Si Api Api already exported 12,700 t of dried fish; in 1904 it exported 25,900 t of dried fish and 2,700 t of *belacan* (Haga 1917). There were several reasons for this spectacular increase. To begin with, there was the extraordinary abundance of fish in the estuary. The organic matter continually being brought from the interior of Sumatra by the Rokan River, the constant mixing and oxygenation of the water taking place because of the tidal bore, and the dense mangroves all contributed to the growth of many species of finfish and shrimps.

Just as important, however, was the way fishing was organized to produce for the market. The fishers themselves were organized into small *kongsis*, each headed by a *towkay bangliau*, who provided the capital equipment needed to undertake the business and fed and gave the fishers cash advances. Up to the early 1900s, fishing was almost entirely conducted by means of *jermals*, which were placed in the estuary in such a way that fish were driven into them on the outgoing tide. The fish immediately underwent a preliminary salting on board the boat bringing the fish to shore; once on shore the fish were placed on extensive platforms to dry. The *towkay bangliau* paid the fishers a share of the proceeds from the sale of the fish to local fish traders, Chinese from Java, once they had deducted their expenses and their own share.

The linchpin for the whole business was the salt farm, which supplied the vital ingredient needed to preserve fish to be shipped long distances. Unlike in most other parts of the Netherlands Indies, where salt had to be bought at very high prices from the government's salt monopoly, the government leased out the exclusive right to sell salt at Bagan Si Api Api. This monopoly or farm was held by a syndicate made up of prominent Chinese businessmen on both sides of the straits. The important point here is that up to about 1905 this syndicate held the farm with little or no competition from rival syndicates and that therefore the rent it had to pay to the government was low. This meant that the farm was able to sell salt - cheap, good quality salt bought in Singapore, but which had been shipped from Aden across the Indian Ocean - to the fishers at a price well below that prevailing in those parts of the Netherlands Indies subjected to the government's monopoly. The farm could, of course, have raised the price of salt but apparently chose not to in the expectation that a low price would

stimulate production. Since the rent that the syndicate owning the farm had to pay the government was fixed for the term of each contract, whatever was collected over and above that rent could be kept. In order to promote production still further the salt was supplied on credit to the *towkay bangliau*. As well as receiving credit, the *towkay* received cash advances from the fish traders. Thus, there was an elaborate system of credit backed up by a powerful syndicate, that lubricated the expansion of production. As well as supplying credit, the syndicate owning the farm was connected with a shipping line that carried the fish to Singapore and Java. The result, according to Colijn (1905), the first Dutch official to report on Bagan Si Api Api, appears to have been "a good livelihood for the fishers, a great profit for the [Netherlands Indies] treasury, and, certainly, a gold mine for the farmer".

Although the main market for the salt fish produced at Bagan Si Api Api was Java, this fish had to compete with fish imported from Siam (and Cambodia) via Singapore as well as locally produced fish, though to a much less extent, for reasons to be explained shortly. The summary report of the Welfare Commission comments that the imported fish "is transported from the bigger harbors to the interior. On the south coast one finds Siam fish in the most remote *desas*" (Hasselmann 1914). In general the consumers of Java preferred "Siam fish", which consisted of *kembong* (*Rastrelliger* spp.) preserved in a great deal of salt (in fact some of this salt could be reused, which was part of the attraction of the product), to the *ikan busuk* ("rotten fish") of Bagan Si Api Api and were therefore prepared to pay a higher price for it. As a result, the price in Java of fish produced in Bagan Si Api Api was largely determined by the supply to Siam fish, which meant that when there was a big supply of Siam fish the price of fish from Bagan Si Api Api fell.

Between 1904 and 1910, exports of dried fish from Bagan Si Api Api fell from 25,900 to 18,900 t. Just why this occurred is unclear, but we should note that three things were happening simultaneously: the farm raised the price of salt, the estuary was silting up and the many years of intensive fishing were having at least some effect on stocks, as evidenced by the fact that far fewer large fish were being caught than had been a few years earlier. In their often bitter debates about the causes of the decline (*achteruitgang*) of Bagan Si Api Api, Dutch officials tended to focus on one or another of these factors. Most blamed the rising price of salt. Although they argued that, thanks to a big increase in competition for the farm from rival syndicates, it now paid a much bigger rent to the government and so had to recoup its costs by selling salt at a higher price, it is worth noting that the farm might still have made a profit by keeping prices low and thereby promoted production still further. That it chose not to do this may have been because it may have no longer been possible to keep pushing up production as in earlier years. Most officials pointed out that the silting up of the estuary - the extension of the shore of this part of Sumatra - had been going on for thousands of years, but it is possible that the process had speeded up because of the large-scale cutting of mangroves for the poles needed to construct *jermals*

as well as cutting of trees upstream for the construction of houses and drying platforms in the town. It is also possible that the cutting of mangroves, combined with prolonged, intensive fishing (Gobee [1912] reported that the *jermals* “stood close together” in the inner part of the estuary, catching “everything carried out with the ebb tide”), was having an effect on fish stocks. In short, I am suggesting that these circumstances may have given the farmer little choice but to increase the price of salt to try to meet his obligations to the government. However we might explain the “decline”, we can at least say that, as the price of salt went up, the fishers began to use less salt when preserving their catches and that as a result the reputation of fish from Bagan SiApiApi began to fall. Moreover, the rising price of salt seems to have encouraged the production of *belacan*, for which less salt is needed in relation to the value of the product. In the early 1900s, fishers began interweaving split rattan into the interstices of the rattan screen used in *jermals* and placing sacking behind the screens so that their *jermals* could catch the tiny shrimp used to make *belacan* as well as the larger ones, which were dried. Soon after that they began fishing with another Malay device, the *ambai*, which they used specifically for catching shrimp. Between 1904 and 1909, exports of *belacan* jumped from 2,700 to 10,100 t, while those of dried shrimps rose from 400 to 1,200 t. (Although the fishers were putting much more effort into catching shrimp, it is possible that the capture of huge quantities of finfish had resulted in less predation on shrimps [see Pauly 1982, 1984; Pauly and Mathews 1986] and that therefore there also were more shrimps to be caught, at least in the first few years.) After 1909, even exports of *belacan* fell somewhat, apparently because the traders who prepared it were skimping on salt, the price of which continued to rise.

The story of the fishing industry of Bagan SiApiApi in the 1910s is extremely complex. In brief, the syndicate tried to collect the huge sums it had lent the fish traders; many of the traders were unable to pay up and went bankrupt; a new syndicate took over the farm, making a profit until the government clamped down on some of their surreptitious impositions and the price of salt on the world market went up because of the shortage of shipping in World War I; and the Dutch government, sick of the debates about Bagan SiApiApi and eager to get rid of this farm and similar ones (by this time the great opium farms of Java had been replaced by a government monopoly), arranged for salt to be sold by a company made up of local traders. It is notable that after 1920 exports again rose, as the price of salt fell somewhat, a moveable trap, the *si stji*, was introduced, thus getting around the huge cost involved in building new *jermals* whenever the water in which they stood silted up, and fishing with drift nets further out in the straits became an important part of the industry. In the 1920s and 1930s, Bagan SiApiApi was still the leading fishing port in the Western Archipelago and, for that matter, one of the biggest in the world.

Another place producing for export, though on a smaller scale than Bagan SiApiApi, was Trengganu on the east coast of the Malay Peninsula. Here fishing was performed using a

big variety of nets, since fishing stakes could not be used on the east coast except where they could be protected from the northeast monsoon by placing them on the lee side of an island. Fishing and the processing of the catch were conducted entirely by the Malays of Trengganu - the men went to sea, while the women salted and dried what the men caught - but the Chinese controlled the marketing and export of the fish after processing. The key institution was the farm for the right to collect the export duty on dried fish, virtually all of which was shipped to Singapore. (In the early 1930s, it was reported that the fish produced in Trengganu and shipped to Singapore was sent on to Java [Anon. 1932]. This pattern may have been in place for a long time.) At this stage I can say nothing about the connection between the Chinese who held this farm and those who bought dried fish and shipped these to Singapore, but it is likely that in order to increase profits (as exports went up all the profits collected could be kept as long as the rent to the sultan was paid) they lent money to the traders and that they in turn advanced money to the fishers on the understanding that they would be able to buy their catch. In any case, production increased rapidly in the late nineteenth and early twentieth centuries. In 1910, the value of exports of fish - the most important export of the state - was (Straits) \$464,000, roughly equivalent to 2,500 t of dried fish; by 1914, the value of exports had climbed to \$781,000 (Shaharil 1984). An important factor in the steady increase in production was access to cheap salt produced in the salt pans of the inner part of the Gulf of Thailand. Malay traders carried *belacan*, also made in great quantities in Trengganu, to Siam and returned carrying salt.

Bagan Si Api Api and Trengganu were the two largest exporters of dried fish in the Western Archipelago. At Bagan Si ApiApi the *towkay bangliau* had nearly all fallen by the wayside by about 1914 and the fishers (who had prospered even as the farmer and the traders had gone under) owned and operated their own stakes and boats. As far as I can tell, no one owned several stakes and hired people to work them. Similarly, along the coast of Trengganu, noted Clifford (1895), “owners of boats and nets usually take an active part in the fishing operations; and the capitalist who owns many crafts and lives on the income from their hire is almost unknown.”

Fig. 2 provides a rough picture of the trade in dried fish, *belacan* and salt about 1910. A number of things are worth noting. First, an extensive trading system had developed by this time, largely in the previous thirty years. Second, much of the salt needed to sustain the growth in production in the Western Archipelago came from Arabia and Siam. And, third, although the plantation and mining districts of Sumatra and Malaya imported large quantities of fish, by far the biggest importer of fish (and *belacan*) was Java.

This last point brings us to a brief look at Java. Although a huge quantity of fish and *belacan* was being imported into Java, the case of fish imports from all sources amounted to only about 1.5 kg per person per year, which probably was a fairly small proportion of average consumption. Nevertheless, the fact remains that the Java Sea contained more than enough fish to meet the demand in Java. During the late nineteenth




Fig. 2. Trade in dried fish, belacan and salt ca 1910 (Butcher 1992).
 [Gambar 2. Perdagangan ikan asin, belacan dan garam sekitar tahun 1910 (Butcher 1992).]

and early twentieth century, production in Java appeared to have increased at no more than (perhaps slightly less than) the rate at which the population grew. That it did not increase faster to meet the rising demand was the result of several factors, of which I will mention two associated with the preservation of catches. First, the *layang* fishers did not carry salt with them when they went out to sea, for a variety of reasons. According to the government's "Welfare Commission", the fishers of Tuban did not do so because there was not enough room on board their boats for salting fish, and those of Pekalongan had a "superstition" that stopped them from taking any salt to sea, while those of Probolinggo claimed that fish avoided nets that, because of the lack of space on their boats, had come in contact with salted fish stored on board (Anon. 1905a). Whatever the reason, the fact that the fishers did not salt their catches on board greatly hindered the expansion of production:

Because fish are not salted on board, they must be sold as quickly as possible, especially where transport is poorly developed. The result is that on long trips the fish are already sold while underway, that a brief but frantic trade takes place at the landing places, and that the market is quickly oversupplied and, when there are very large catches, the fish are sometimes unsaleable (Hasselman 1914).

Even more important was the high price of salt. The Javanese had developed a variety of techniques of preserving fish by simply drying them or by using salt-laden beach sand, but these were not very effective in preparing a product that would not be consumed within a very short period. In the early 1900s the colonial authorities, caught up in the liberal spirit of the "Ethical Movement" and desirous of increasing the supply of fish to the growing population of Java, introduced a system

of salting sheds at which fishers (at least those living nearby) could preserve their catch using salt supplied by the government monopoly. Although the salt sold at these sheds was very much cheaper than that sold to ordinary consumers, the price was still much higher than it was in Siam and the fishers were prohibited from taking the salt with them to sea, which of course limited the time they could spend catching fish, though this rule appeared to have been relaxed in a very small number of places by about 1920. One of the effects of the system of salting sheds appeared to have been that the men took over work that had been carried out by the women, at least that is what is implied in a comment by one critic that "it is extremely desirable that the women be able to remain at home salting [the fish] while the men are at sea" (Schippers 1928).

Just how restrictive the government monopoly was is illustrated by the case of Sapeken in the Kangean Islands. Up to the early 1900s, there was a thriving fishing industry there, thanks largely to the availability of cheap salt imported from Makassar. The government knew of this trade but turned a blind eye to it (Anon. 1905). In 1907, however, it prohibited the use of Makassar salt and forced the people of Sapeken to buy salt from the government monopoly. Following this clampdown, "fishing came to standstill; there were shortages because there were no fish with which to pay for the necessities of life, all of which Sapeken must import; many left the island and set up elsewhere, the population of Sapeken village falling from 6800 to 3700 people" (Van der Plas, in van Kampen 1922). When the government set up salting sheds at Sapeken, fish exports rose but not to the pre-1907 level.

During the 1920s and 1930s, the process of commercialization continued and more and more fishing became oriented to production for the market. R. Firth's sketch of the economic history of a Kelantan fishing village before he did his fieldwork there in 1940 provides a unique insight into some of the forces behind this process. According to Firth (1975), the construction of a road linking Perupok with the main population center of Kelantan "altered the fishing economy considerably", as it made possible a regular bus service, which created a market for fresh fish, for which the fishers were able to get much higher prices than the fish they sold for drying. More people took up fishing; whereas fishing had remained a subsidiary activity for many people "the life of the fishermen became more completely divorced from agriculture"; and money rather than barter transactions became the rule. "The life of the community", Firth (1975) concluded, "finally merged much more completely into the general economy of the State". (Bailey 1980, 1983 described a similar impact of road building for another Malaysian village, Mangkok.)

As already suggested, capital and finance were crucial to the whole process of commercialization, for "a good deal of capital... is needed for sea fishing" (Hasselman 1914). Capital was needed to buy boats, nets and fishing stakes, and credit was often needed to cover operating and living costs, particularly because of the seasonal and often unpredictable

nature of fishing as a livelihood. There was a great variety of ways in which capital and credit were provided. Some fishers had enough money to buy a boat and the needed gear, and some bought them on time purchase. Gathering together the capital needed to mount an elaborate fishing operation, as in the case of the liftnet fishery described by Firth, was often made possible by pooling the boats and nets of several fishers on the understanding that the catch would be divided up according to how much capital as well as how much labor the fisher had contributed to the operation. Borrowing money to buy capital items as well as to cover operating and living costs was widespread. Almost everywhere fish dealers were the principal lenders, having as they did a vested interest in pushing up the quantity of fish available for them to buy and sell. The nature of these loans varied greatly. At one extreme, generally when a small amount of money was involved, fishers were able to pay off their debts by handing over a proportion of their catch to the dealer for a limited period. At the other, the dealers in fact owned the boats and gear and advanced cash to the fishers, in return for which the fisher was required to hand over all of his catch to the dealer. In general, it appeared that as time went by, as fishers became increasingly oriented to the market, there was a trend towards arrangements of the second type. Referring to the situation in 1940, Firth (1975) made the point that the fishers of Trengganu, where production had long been oriented almost entirely towards the export market, were much more tightly bound by debt to dealer-lenders than were those of Kelantan, where commercialization came later and where a much smaller portion of the catch was exported. The Welfare Commission concluded that many fishers along the north coast of Java "are gradually coming financially completely under the control" of the powerful *kongsis* of Batavia fish traders (Hasselman 1914), who, according to van Kampen (1922):

... give advances to the fishers at high interest rates but also at great risk, for if the fishing gear or *prahuis* lost the moneylender... usually loses what he has lent. The debt is settled at the auction at the fish market, which is in the hands of the same *kongsi*. Because they have easy access to credit the fishers are able to maintain their *prahus* and fishing gear properly; the moneylenders stimulate them - in the interests of both - to greater energy and the business progresses. Fishers from Bantam, Krawang, and Tegal come to Batavia in order to fish there with advances from the *kongsis*.

It should be added that many fisheries officers of the time took a much less favorable view of fish dealers. Referring to Malaya, Stead (1923) wrote that:

In nearly all cases the fisherman is practically "bound" to sell to a specific *towkay*. Not legally so, perhaps, but for all practical purposes there is no escape, as he is in the hands of this *towkay* financially. The *towkay* has probably

advanced him money for his nets or for his other gear, and has allowed the man to run into debt in other ways, with the express purpose of keeping the necessary hold over him, so that the man will regularly contribute a fish supply to the *towkay*. By getting together a number of such "clients" the *towkay* is sure of his own livelihood.

Sentiments such as these were behind a number of attempts by colonial authorities in both British Malaya and the Netherlands Indies to promote cooperatives and alternative sources of credit for fishing communities in the 1920s and 1930s.

The Beginnings of Mechanization

What Cushing (1988) called "the first industrialization" of fisheries accelerated in the 1890s with the development of the otter trawl. One of the first projects of the Fishery Station set up in Batavia in 1904 was to explore the possibility of introducing the otter trawl and other gears commonly used in the West, apparently on the assumption that fishing would become a Western capital intensive industry just as mining and the cultivation of various crops had over the previous half century. Beginning in 1907, the *Gier*, a 650-t steam vessel, experimented with the otter trawl in the Java Sea, but in 1911 the Fishery Station abandoned the project. One of the main reasons was the nature of the sea bottom, as van Kampen (1922) explained a few years later:

Where, as in the Java Sea, the sea is not too deep for this fishery, the bottom consists of soft mud, into which the trawl sinks, or of sand, studded with coral, that damages the net with its sharp edges, or the bottom is in fact covered with various lower forms of animals (sponges, sea urchins, and so forth) which by their weight endangers the captured fish and even the net itself. The otter trawl of the research vessel was more than once shredded by meter-high cup sponges.

In the mid-1920s the British conducted similar experiments with the *Tongkol* (292 GT) in the waters off the Malay Peninsula and concluded that demersal species did not exist in sufficient abundance to sustain a trawl fishery, at least at the prices of fish then prevailing. In light of later events, however, an official report pointed out that "large catches can be made by the trawl within the ten fathom line between the Dindings and Penang" and suggested that "small motor-tractors built locally and manned by Asiatics promise to be highly remunerative in this area" (Anon. 1927).

While the colonial authorities were unsuccessfully trying to industrialize fishing by introducing the otter trawl towed by large vessels, groups of Japanese fishers had introduced a revolutionary new method of fishing, *muro ami*. The movement of Japanese fishers into the Western Archipelago was part of

the diaspora of Japanese merchants, photographers, dentists and planters that took place beginning in the late nineteenth century but accelerating after World War I, but it appeared to have been prompted at least in part by increasing pressure on their fishing grounds in the Ryukyu Islands and regulations by the Japanese government closing off certain areas from fishing. In any case, Japanese fishers very quickly made a big impact in "the South Seas". As early as 1921, Maxwell (1921a) noted that the Japanese based at Singapore "catch more than the rest of the local fishermen combined". In about 1925, a group of Japanese had set up in Batavia as well; in the late 1930s, there were three *kongsis*, supplying (by value) about a third of the fish auctioned at Batavia.

A number of aspects of this new form of fishing need to be emphasized. First, the Japanese exploited a new niche, the waters above the slopes of coral reefs. Before the arrival of the Japanese, some indigenous fishers had fished on the reef slopes using traps and spears, but their catches were very small and fusiliers (Fam. Caesionidae), fish that predominated in this niche, had been sold in local markets in very small quantities. In general, nets were not effective on reef slopes - they caught on the coral and rip - but the *muro ami* fishers fixed the net to the sea bottom and frightened the fish into the net. Second, this method of fishing required tremendous strength and skill on the part of the fishers, for they had to dive to the seabed to fix the net to the bottom and then herd the fish into the net by swimming along the surface toward the mouth of the net carrying a long rope to which were attached strips of white cloth and at the end of which was a lead weight. "They are more like fish than men", marvelled a fisheries officer who spent three months watching them at work (van Pel 1938). Third, *muro ami* also required an extremely high degree of organization. This is evident not only in the way the 50 men in a *muro ami* team cooperated in catching the fish but also in the link between the team and the market. While one of the two motor boats supporting each team carried the catch on ice to the market, the other was on hand or on its way to the fishing ground to pick up more fish. A team would stay out for as long as six months at a time while the carriers shuttled back and forth. "No time is left unused and the business operates continuously" (Reuter 1940). And, fourth, this business required a great deal of capital. None of the boats in a *muro ami* team was particularly large - a typical carrier was 15 - 20 m long - but the total amount of capital tied up in the carriers, the smaller boats used during fishing and the net was considerable, as was the amount of money needed for fuel, ice, food and other supplies. However, while the amount of capital involved was great and one person might be the sole owner of one or more *kongsis*, there is evidence that at least some owners exercised little direct control over their business. According to Reuter (1940):

The fishers work for the boat owner as a team, so that he does not deal with them separately but only with one or several team leaders. These leaders enjoy a great deal of

independence, so that one often gets the impression that in essence the ship owner merely hires his vessels out for a share of the profit. This impression is particularly strong in the case of one ship owner who has a *muro ami* business and has no understanding at all of fishing and often does not know where the boats are operating.

During the 1920s and 1930s, teams of *muro ami* fishers went farther and farther afield in their search for new fishing grounds. Those based at Singapore fished in the Riau-Lingga Archipelago, near the Anambas Islands, up the east coast of the Malay Peninsula as far as the Gulf of Siam, and along the west coast from Langkawi to the Mergui Archipelago, while those based at Batavia scoured the reefs near Bawean and elsewhere in the eastern part of Java Sea, near Bangka and Belitung, and along the west coast of Sumatra. So effective was *muro ami* that even in the mid-1920s some fisheries officers wondered whether it might threaten stocks; however, none of them appeared to have been particularly alarmed. In 1926, the director of the fisheries department in Malaya pointed out the "obvious danger" of overfishing but, while calling for scientific study of the fusiliers, seemed to accept the statements of the *muro ami* fishers that they did not reach the depths where most fusiliers lived and that therefore the areas they fished were constantly being replenished (Anon. 1925). In 1932, a fisheries officer argued that although the *muro ami* teams had found it necessary to go farther and farther from Singapore "because the nearby grounds yielded comparatively only small catches", stocks were not being depleted but rather the fish were learning how to avoid the fishers (Anon. 1932). Van Pel (1938) reported that few fish were left after the *muro ami* team he accompanied had fished all the reefs on the eastern sides of the Thousand Islands, adding matter of factly that reefs were not fished again for two years after a team had been through. None of the reports I have seen said anything about the possible effects the lead weight used by the fishers had on the coral a major source of reef destruction now well documented, e.g., in the Philippines (Corpuz et al. 1983).

In the meantime an important change was taking place in the way fishing was conducted along the west coast of the Malay Peninsula. Up to the 1920s, various forms of fishing stakes caught the great bulk of fish landed along this coast. "The whole of the West Coast... was dotted by" these stakes (Yap 1976). Beginning in about 1920, however, the Federated Malay States (FMS) government introduced regulations that discouraged use of these stakes. Because the stakes were mainly built using poles taken from mangroves the great expansion in the number of stakes had seriously depleted mangrove forests in the area. This pushed up the price of the materials needed for constructing stakes, a trend accentuated when the government closed off certain sections of the forest from cutters. The government also regulated the activities of the stake fishers more closely, mainly in an attempt to protect fish stocks, which officials believed were being undermined by

the stakes, outfitted as many were with fine-meshed nets that caught large quantities of immature fish. (Small and immature fish were used either as fertilizer or as food for pigs and ducks.) In 1920 the Perak government banned *ambais* outright. As stake fishing became either less profitable or, in some forms, altogether impossible, a new form of fishing, the purse seine, was introduced at Pangkor. Whether the people who introduced or quickly took up purse seining were the same people being pushed out of the stake fisheries or whether they were relatively recent arrivals from China is unclear to me. What is clear is that these fishers introduced a net "hitherto unknown in Malaya and... never... used by local fishermen". The main target of the fishers was *kembong* (*Rastrelliger* spp.) "both abundant at times, and universally esteemed" (Anon. 1930), which were caught and then either salted and dried or boiled in brine. The main market for these products was the tin mines and rubber plantations of the interior, but some were also sent to Singapore and Penang and then transhipped to "other countries" (almost certainly the Netherlands Indies) - the incentive to produce for an export market was increased sometime during the 1930s when the FMS government abolished the export duty on salt fish.

Up to 1937 the purse seine - a net 170 fathoms long and 28 fathoms deep with a half inch mesh - was taken to sea in 32-meter, long sailing junks, each of which was accompanied by 3 small boats, 2 for operating the net and 1 for helping to locate fish and to convey supplies and the catch. The crew consisted of a captain, 2 helmsmen, 2 expert *kembong* watchers, 11 deck hands, and 1 or 2 cooks, usually wives of the captain. The time when fishing could take place was limited, mainly because the fish could be seen swimming very near the surface only on moonless nights. Catches varied enormously depending on how many fish were present, whether the *kembong* watcher had spotted a big school, and whether the net could be placed without many fish escaping. Beginning in 1937, the scale of operations increased substantially as many owners replaced their junks with boats powered by 12-hp diesel engines. In 1936, there were 11 sailing boats based at Pangkor that engaged in purse seining; by 1938, all of the sailing boats had been abandoned and the fishing was conducted by 32 motor boats (Anon. 1938). Motorization freed the boats of their dependence on the wind, enabled them to cover a much greater area in a shorter time and to return to port every day, and combined with surveys then being conducted by *R/V Kembong*, gave the fishers access to previously unknown fishing grounds. Between 1931 and 1936, landings of *kembong* at Pangkor shot up from about 800 to 5,000 t. The great problem of the business was how to handle gluts, which severely depressed prices. On a poor night a boat might catch as little as 0.3 t, but catches could reach over 30 t a night. As of 1938 little had been done to overcome this problem, but the fisheries department experimented with freezing and canning *kembong*. The purse seine fisheries of Pangkor resembled many present-day fishing businesses, albeit on a much smaller scale. The owner of the boat and net, usually a fish dealer, had overall control over the

operation but did not take part in the fishing himself. He paid for the fuel, and he paid the crew partly in the form of wages and partly in the form of a share of the proceeds from the sale of the fish (Anon. 1937).

After abandoning the possibility of a capital-intensive trawling industry, fisheries officers in the Netherlands Indies adopted a new line of thinking, one aimed at the improvement of "native" fishing, which it was believed could only take place without sudden change. In van Kampen's view, many of the gears used by Javanese fishers, most notably the *payang*, worked well as they were; what was needed were motors for the *prahu mayang*:

Because of the regular alternation of the sea and land winds the fishing prahus as a rule only need to sail before the wind and for this purpose they are excellently designed. However, variable winds, such as prevail during the transition between monsoons, and calm make their use uncertain and are therefore the reason that fish do not reach the shore fresh. Petrol motors as auxiliary power would be very helpful in this respect. At the same time these could be used for hauling in the net, so that the crew could be smaller (van Kampen 1922).

Apparently little if anything was done in this direction during the 1920s, but when he was appointed to take charge of the development of fisheries in the Netherlands Indies in 1927, C.J. Bottomanne set the motorization of *prahu mayang* as one of his primary goals. At first he experimented with gasoline engines but because of the very high tax on gasoline he then experimented with semi-diesel and diesel engines, which were heavier and more expensive. In West Java, this experiment appeared to have been more successful than in other parts of Java. There, by 1942, about 40 motorized *prahu mayang* had been sold "to domestic skippers and some middle class owners", who "made a handsome profit", for reasons Bottomanne (1959) later explained:

In West Java, where the water was not clear and fishing was possible during the entire period of daylight [in the clearer waters off East Java fishing was only possible at dawn], 20 cycles a day were often made, whereas sailing vessels only accomplished up to 6 cycles in that region. Nets, moreover, were bigger for the motor vessels.

By the outbreak of the Pacific War, however, motorization had not gone very far. During the depression the government lowered the price of salt somewhat, but because fuel prices were high and fish prices fell sharply few people had the capital or the incentive to invest in motorized vessels.

A feature of the 1920s and 1930s was the increasing scientific interest in fisheries, particularly in the Netherlands Indies. Much of the research conducted there was devoted to studying plankton, mainly because studies of the quantity of

plankton in the Java Sea and elsewhere could be used as an indirect indicator of the abundance of fish stock, an issue about which there appears to have been intense debate at the time, in sharp contrast to the nineteenth century when the abundance of fish in the archipelago was an article of faith among Europeans. A.W. Herre's comments give some idea of the nature of the debate as well as his own views on it:

There is an impression, based upon inadequate knowledge and lack of extended field experience, that Indonesian waters are poorer in fish than more northern waters. Chemical analysis of sea water, and limited collections of plankton have been held to prove that there are few fish in tropical waters. The astounding amounts of plankton that occur at times, when planktonic organisms increase to such a vast extent as to destroy most other forms of life, and the frequent recurrence of such phenomena, seem to have totally escaped the observation of those who say tropical waters are poor in fish. When one sees vast schools of sardines, as much at times as ten miles in length, shoals of mackerel... [etc.], he knows that such ideas are erroneous (Herre 1945).

At least within the Netherlands Indies the consensus among Dutch scientists appeared to have been that the seas of the archipelago were indeed rich in fish, even if they rejected the impressionistic statements of the previous century. An implication of this consensus, of course, was that there was no need for the sort of attempts being made in Europe to regulate fisheries. According to Delsman (1939), one of the leading researchers, "there is no question of exhausting the supply anywhere in the Archipelago - except perhaps at some river mouths - nor is any such exhaustion likely to occur as soon in the tropics as it might in Europe, since the high temperature in the tropical seas causes the growth and renewal of the fish supply to take place more rapidly than in colder climates." It is interesting to note, however, that one of Delsman's predecessors, van Kampen, had begun to develop an appreciation of the difficulties of regulating a multispecies fishery, pointing out that rules (such as had been tried in Malaya, where generally fisheries officers were much more concerned about the danger of overfishing) prohibiting small-meshed nets might protect the young of large species but made it impossible to catch shrimps and fully grown small fish (van Kampen 1922).

The Acceleration of Change

The Japanese occupation of the Western Archipelago from 1942 to 1945 had a devastating effect on fish production just as it did on most other economic activities. By the end of the war a great deal of equipment - ice plants as well as boats and gear - had been destroyed or badly damaged, imports of

twine, nets and other materials had been cut off, marketing systems had been disrupted, and the purchasing power of consumers had been greatly diminished. In Malaya, total fish landings in 1944 and 1945 must have been well under half of what they had been in 1940; presumably, the same applied to Indonesia, particularly in Java, one of the places most affected by the occupation. Both in Malaya, to which the British returned in 1945, and in the Netherlands Indies, now proclaimed as independent Indonesia by Sukarno and Hatta but still controlled in some areas by the Dutch, there were desperate shortages of food. In these circumstances one of the greatest concerns of governments was to promote the production of fish, particularly since alternative sources of animal protein were in even shorter supply than they had been before the war. In Jakarta and particularly in Singapore, an added problem was the fact that the principal providers of fish to these markets in the pre-war period, namely, the Japanese fishers, had been removed. (In Malaya, the government rejected the suggestion that in order to overcome shortages the Japanese be allowed to resume fishing on the grounds that this would give them a foothold in the post-war fishing industry.) In the short term, governments did what they could to restore production to pre-war levels by facilitating imports of materials, repairing the infrastructure and restoring marketing arrangements. In Malaya, these efforts had immediate results. By 1948, landings had recovered to their pre-war level. Various restrictions imposed during the Emergency brought about a slight drop in production, but thanks to the boom prompted by the Korean War it soon began to rise again. In Indonesia, the revolution and the unsteady economic conditions that followed made the recovery more difficult than in Malaya, but by 1951, production appeared to have been back to the pre-war level.

In the longer term, governments hoped to increase production by encouraging mechanization. In Malaya, the number of motor boats increased from 327 in 1949 to 7,300 in 1958. According to the annual report of the fisheries department of the newly independent Federation of Malaya, "the introduction of smaller marine diesel engines capable of being installed in moderate-sized boats has resulted in larger, stronger fishing boats being built which are capable of withstanding rough seas and which can voyage far from land in search of fish" (Anon. 1958). The same report gives some idea of the immediate impact that mechanization had:

The fishing grounds all round Malaya, with the exception of the southern part of the Malacca Straits, have been extended in the past few years as a result of mechanization of fishing craft. The extension is particularly evident on the East Coast where the area covered by local craft has been more than doubled since 1953... it is now common for the fishermen to follow the shoals of fish round the coast of Malaya, remaining away from home for lengthy periods, a practice which was virtually non-existent before the introduction of engines

made the boats largely independent of the vagaries of the wind.

In some places mechanization took place with stunning rapidity. According to Firth (1975), the fishers of Perupok:

did not adopt the less powerful and less efficient though cheaper outboard motors.... They hung back until they were convinced of the superiority of the inboard diesel-fuelled motors, to which they converted very rapidly. They were able to observe the motorized craft in areas of the south. In a remarkably short space of time, about eighteen months apparently, all the leading *juruselam* [fishing experts] of Perupok had invested in these motor boats.

In Perupok, motorization not only allowed fishers to reach more fishing grounds more quickly, but it also facilitated the adoption of a new and extremely productive form of fishing, purse seining.

The increases in production brought about by the mechanization were accentuated by other changes taking place at the same time. One was the widespread adoption of synthetic nets, which though much more expensive lasted longer, were lighter and so much more easily handled, were less visible to fish, and required little or no drying. According to the same report, "drift net catches in the Malacca Straits have doubled with the replacement of cotton by synthetic fibers resulting in increased supplies of *tenggiri* and *parang* to the west coast markets and a subsequent reduction in the retail prices of these fish." Also significant were the much greater use of ice and steady improvements in the transportation system.

In Indonesia, production doubled between 1951 and 1967, but mechanization appeared to have contributed much less to increases in production there than it did in Malaya. There was a great increase in the number of motorized boats, but a large proportion of these spent much of their time out of commission because of "shortages of spare parts, shortage of ice, and the elaborate procedures for obtaining sailing permits" (Krisnandhi 1969). Instead, the doubling of production came about mainly because of threefold increases in the number of fishers and nonmotorized "traditional" fishing craft. According to Krisnandhi (1969), the fishing industry during these years "has grown but not developed". Nevertheless, as he looked at the Indonesian fishing industry from his vantage point right at the start of the New Order, what struck him was not that production had increased so little in relation to the estimated potential but that it had expanded as much as it had:

The industry had expanded substantially since 1951, for not only has total production risen but so also has fish consumption per head. The industry has been able to do this only by making substantial investments in fishing equipment. It has done this, moreover, from its own resources during a period when the national economy was becoming more and more unstable and the climate for economic

enterprise was becoming progressively less favourable, and in the face of very considerable specific handicaps - poor communications, poor transport facilities, a "lack of modern processing and storage facilities, and a high cost marketing system."

"Under these circumstances," he added, "it is not surprising that investments by producers for the most part took traditional forms or that returns to capital and labour declined" (Krisnandhi 1969).

In both Indonesia and Malaya, fisheries officers believed that, at least in the long term, the only way the growing demand for fish could be met would be by industrializing fishing. Just like their predecessors earlier in the century, they turned their attention to the possibilities of trawling. In the 1950s, the Directorate General of Fisheries (DGF) of Indonesia "conducted experimental trawl fishing in the Madura Strait and the Java Sea. The trials were targeted at finfish and regarded as successful, but local fishers did not respond, among other reasons, due to the difficulty of obtaining engines and spare parts" at this time (Bailey et al. 1987). The premise behind a trawl survey conducted by the Colonial Office on behalf of the governments of Malaya in 1955-56 was that because "inshore 'subsistence' fisheries are now saturated," "producing as much fish as they [are] capable of," it was necessary to look to hitherto unexplored extraterritorial [meaning, at that time, beyond the three-mile limit] waters" to feed "the very rapidly growing population" and that these new fishing grounds could only be exploited by means of powerful long range vessels (Ommanney 1962). The report of this survey concluded that only a vessel of the size and power of the research ship, the 208 GT *Manihine*, could possibly engage in otter trawling, since only such a vessel could extract the net when as often happened, it got stuck in the mud and hauled up the sponges and other marine life with which the bottom was "heavily overgrown" (and which we now know are an integral part of the ecology of soft-bottom communities, "provid[ing] habitats for invertebrates and shelters for fish" [Pauly 1986]). It also concluded, however, that not even such a vessel could make a profit, because of the huge capital costs, high running expenses, the low prices of fish (no mention is made of shrimp in the report), and, not least, the fact that stocks of demersal fish were not particularly abundant. In May 1956, when the *Manihine* was still conducting its survey, a special committee set up to investigate the fishing industry concluded that "experiments have all demonstrated the unsuitability of western gears for use in Malayan waters" (Abdul Aziz 1956). The fisheries department still hoped to introduce industrial fishing, but by 1958 this hope appeared to have been focused on exploiting tuna in the Indian Ocean, for "the fish stocks of these waters offer the only realistic hope for the rapid increase in fish supplies to the Federation to meet the ever increasing demand of the growing population" (Anon. 1958).

Within the space of a few years, however, trawling became one of the most powerful fishing methods in the Western Archipelago. Considering the profound impact of

trawling, it is surprising how little has been written about its introduction, but we can piece together an outline of how this came about. According to one account, it happened in this way:

The development of trawling in Penang could be traced to the breakdown in the barter trade with Indonesia with the onset of confrontation between the two countries in 1963. The barter traders started looking for alternative use for their boats which were of 30-50 tons category. About this time a few fishermen who were interested in trawling went to Thailand to observe trawling operations there. They brought back two types of trawl nets, namely the otter trawl and the beam trawl. They found out that the unused barter traders' boats were suitable and could easily be converted for trawling (Lam and Pathansali 1977).

Other versions of this story differ slightly in detail. Yap (1976) said that the fishers of Pangkor learned about trawling from some Thai fishers who visited the island. Leaving aside such details, however, certain points are clear. First, the introduction of trawling had little to do with any attempts by the government to promote it. Second, the boats used as trawlers were very much smaller than those used in government-funded experiments. And, third, the adoption of trawling in Malaysian waters had a Thai connection. To understand the second and third points, we need to refer briefly to what happened in Thai fisheries at about this time.

Up to 1960, the marine fisheries of Thailand was (as far as finfish were concerned) devoted almost entirely to the capture of pelagic species, particularly *Rastrelliger* spp. The demersal fish of the Gulf of Thailand were virtually untouched by fishing. In 1958, Klaus Tiews, who had been conducting research in the Philippines, where trawling had been practiced in various forms for several decades, "recommended to the Government that it should be determined whether the sea fisheries could be expanded by introducing a trawl fishery" and, in 1961, work along this lines began with aid from the Technical Assistance Program of the Federal Republic of Germany (Tiews 1973). As soon as such a net had been developed, the fisheries department demonstrated its use to fishers on the fishers' own boats (an inducement was that the owners kept the catch) and trained them to tailor the nets themselves. Within just two or three years, several hundred boats had adopted the otter trawl designed by the German team.

It was, apparently, this trawl net that the fishers of the west coast of Malaya adopted beginning in 1965, converting to trawlers not only boats that had been used in the barter trade but also many purse seiners as well. In the following year, 1966, a number of fishers based at Bagan Si Api Api and fishing in the areas surrounding the Rokan estuary took up trawling. According to Bailey et al. (1987), "the profitable operations of Malaysia's trawler fleet provided the technical inspiration for this gear [otter trawls] to be adopted by Indonesia's fishermen operating in the Malacca Straits", but the net adopted appears to have been of a different type, as

suggested by Unar's statement (1973) that "this fishery is characterized by the operation of wooden sampan-like motorized vessels of 5-20 GT employing a single Gulf-type shrimp trawl of 40 feet head-rope." In any case, as this statement indicates, the prime target of the trawl fishers in the Strait of Malacca was shrimp, the price of which was rising dramatically because of increasing demand, particularly from Japan, then entering a period of great prosperity. It was, in short, the increasing value of catches on the international market (rather than the demands of the domestic market) as well as technological changes that suddenly made trawling such a profitable investment. In Indonesia, the New Order government created conditions that encouraged investment in the fishing industry.

Conclusion

From our vantage point in the mid-1990s, we can see the period from about 1955 to 1970 as a turning point in the history of fisheries in the Western Archipelago. During these years production increased dramatically, first in Malaysia and then in Indonesia, because of rapid mechanization and a number of related changes. Where once fishers tended to own and operate their own boats and nets, more and more, those who went to sea were employees of land-based entrepreneurs, intent on maximizing the return on their investments. As the intensity of fishing increased, so too did conflict between those operating different types of gear, as did pressure on what had, only a few decades earlier, been assumed to be an inexhaustible resource. The challenges that fisheries officers and scientists face today date from this period of momentous change. Nevertheless, the rapidity with which change took place at this time can only be understood in relation to the changes that had already taken place. By the 1950s, fishing was highly commercialized; the population of the Western Archipelago had grown manyfold from what it had been in the middle of the nineteenth century, thanks in part to the nutrition provided by fish; there were signs of overfishing in certain quite localized areas, such as the inner part of the Rokan estuary and the reef slopes where the Japanese *muro ami* teams fished, just as there were signs that refuse generated by human activities on land were a threat to fishing (Maxwell [1921b] stated that the *trubuk* fishery of Malaya "appears to be on the verge of extinction" because of the pollution caused by tin mining); and fisheries scientists had made progress in mapping out the location of the remaining areas that could be exploited, even if they had little idea of the potential of demersal stocks. The revolution that occurred between 1955 and 1970 grew out of the great changes that had taken place over the preceding century - even if, from our vantage point today, they appear less spectacular.

References

- Abdul Aziz bin I. 1956. Report of the Committee to Investigate the Fishing Industry. Government Press, Kuala Lumpur.
- Abdullah bin A.K. 1970. *The Hikayat Abdullah*. Oxford University Press, Kuala Lumpur. A.H. Hill, translator.
- Abdullah, M. 1970. *The Hikayat Abdullah*. Translated by A.H. Hill. Oxford University Press, Kuala Lumpur.
- Anderson, J. 1824. Political and economic considerations relative to the Malayan Peninsula, and the British settlements in the Straits of Malacca. William Cox, Prince of Wales Island. Reprinted for MBRAS by Malaysia Printers, Singapore, 1965.
- Anon. 1882. "Zeevisscherijen langs de Kusten der Eilanden van Nederlandsch-Indië", *Tijdschrift voor Nijverheid en Landbouw in Nederlandsch-Indië* 26:75-145, 157-201, 257-272.
- Anon. 1905. *Onderzoek naar de Mindere Welvaart der Inlandsche Bevolking op Java en Madoera: Overzicht van de Uitkomsten der Gewestelijke Onderzoekingen naar de Vischteelt en Visscherij en daaruit Gemaakte Gevolgtrekkingen, Eerste Deel. Tekst*. Landsdrukkerij, Batavia.
- Anon. 1925. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1927. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1930. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1932. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1937. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1938. Annual report of the Fisheries Department, Straits Settlements and Federated Malay States.
- Anon. 1958. Annual report of the Fisheries Department, Federation of Malaya.
- Bailey, C. 1980. The road to Mangkok: achieving equitable distribution of benefits from small-scale fisheries development. *ICLARM Newsl.* 3(3):10-12.
- Bailey, C. 1983. *The sociology of production in rural Malay society*. Oxford University Press, Kuala Lumpur. 226 p.
- Bailey, C., A. Dwiponggo and F. Marahudin. 1987. Indonesian marine capture fisheries. *ICLARM Stud. Rev.* 10, 196 p.
- Bottomanne, C.J. 1959. *Principles of fisheries development*. North-Holland Publishing Company, Amsterdam.
- Butcher, J. 1992. Some notes on the trade in dried fish, *belacan*, and salt c. 1910. Paper presented at the Conference on Island Southeast Asia and the World Economy, Economic History of Southeast Asia Project, November 1992. Australian National University, Canberra.
- Clifford, H. 1895. Expedition to Trengganu and Kelantan. *J. Malay. B. R. Asiat. Soc.* 34, 1(1961). Originally printed for Colonial Office use in 1895.
- Clifford, H. 1897. *In court and kampong*. First ed. Second ed. Published in 1927 by Richards, London.
- Colijn, H. 1905. Advies betreffende de Zoutpacht in het gewest Oostkust van Sumatra, 3 August 1905, in Verbaal 15 June 1907, no. 25. Archives of the Ministry of Colonies, Algemeen Rijksarchief, The Hague.
- Corpuz, V.T., P. Castañeda and J.C. Sy. 1983. Traditional *muro-ami*, an effective but destructive coral reef fishing gear. *ICLARM Newsl.* 8(1):12-13.
- Crawford, J. 1820. *History of the Indian Archipelago*. 3 vols. Archibald Constable and Co., Edinburgh.
- Cushing, D.H. 1988. *The provident sea*. Cambridge University Press, Cambridge.
- Delsman, H.C. 1939. Fishing and fish-culture in the Netherlands Indies. *Bull. Col. Inst. Amsterdam* 2(2): 92-105.
- Firth, R. 1975. *Malay fishermen: their peasant economy*. W.W. Norton, New York.
- Gobée, E. 1912. De Oorzaken van den Achteruitgang van de Vischindustrie te Bagan Si Api Api. *Mededeelingen van het Visscherij-Station te Batavia* 7: 3-18.

- Haga, B.J. 1917. De Beteekenis der Visscherij Industrie van Bagan Si Api Api en hare Toekomst. *Economist*, p. 237-62.
- Hasselmann. 1914. Algemeen overzicht van de uitkomsten van het welvaart-onderzoek gehouden op Java en Madoera in 1904-1905. Martinus Nijhoff, s-Gøbenhavn.
- Herre, A.W.C.T. 1945. Research on fish and fisheries in the Indo-Australian Archipelago, p. 167-175. *In* P. Honig and F. Verdoorn (eds.) *Science and scientists in the Netherlands Indies*. Board for the Netherlands Indies, Surinam and Curaçao, New York.
- Kampen, P.N. van. 1922. *Visscherij en Vischteelt in Nederlandsch-Indië*. H.D. Tjeenk Willink and Zoon, Haarlem.
- Krisnandhi, S. 1969. The economic development of Indonesia's sea fishing industry. *Bull. Indones. Econ. Stud.* 5(1): 49-71.
- Lam, W.C. and D. Pathansali. 1977. An analysis of Penang trawl fisheries to determine the maximum sustainable yield. *Fish. Bull.* No. 16. Ministry of Agriculture, Kuala Lumpur, Malaysia.
- Low, H. 1848. *Sarawak: its inhabitants and productions*. Frank Cass, London, reprinted 1968.
- Maxwell, C.N. 1921a. Preliminary report on the economic position of the fishing industry of the S.S. and F.M.S. Singapore.
- Maxwell, C.N. 1921b. Malayan fishes. *J. Straits Br. R. Asiat. Soc.* No. 84: 181-280.
- Moor, J.H., Editor. 1837. *Notices of the Indian Archipelago (Singapore)*. Frank Cass, London, reprinted 1968.
- Ommanney, F.D. 1962. Malayan offshore trawling grounds. *Colonial Office Fish. Publ.* No. 18, 1961. HMSO, London.
- Pauly, D. 1982. A method to estimate the stock-recruitment relationship of shrimps. *Trans. Am. Fish. Soc.* 111(1):13-20.
- * Pauly, D. 1984. Reply to comments on prerecruit mortality in Gulf of Thailand shrimps stocks. *Trans. Am. Fish. Soc.* 113:404-406.
- Pauly, D. 1986. Problems of tropical inshore fisheries: fishery research on tropical soft-bottom communities and the evolution of its conceptual base, p. 29-37. *In* E. M. Borgese and N. Ginsburg (eds.) *Ocean yearbook 6*. University of Chicago Press, Chicago.
- Pauly, D. and C.P. Mathews. 1986. Kuwait's finfish catch three times more shrimps than its trawlers. *Naga, ICLARM Q.* 9(1): 11-12.
- Pel, H. van. 1938. *Beschrijving der Beoefening in de Japansche Moero Ami Visscherij*. Mededeeling van de Onderafdeeling Zeevisscherij 2B: 222-25.
- Rathborne, A.B. 1898. *Camping and tramping in Malaya: fifteen years' pioneering in the native states of the Malay Peninsula*. Swan Sonnenschein, London. Reprinted. Oxford University Press, Singapore, 1984.
- Reuter, J. 1940. Een nieuwe inheemsche Visscherij. *Koloniale Studiën* 24:199-218.
- Schippers, A.W. Jzn. 1928. De Zeevisscherijen van Nederlandsch-Indië. *Koloniaal Tijdschrift* 17: 15-37.
- Shaharil, T. 1984. *After its own image: the Trengganu experience, 1881-1941*. Oxford University Press, Singapore.
- Stead, D.G. 1923. *General report upon the fisheries of British Malaya with recommendations for future development*. Government Printer, Sydney.
- Tiews, K. 1973. Fishery development and management in Thailand. *Arch. Fischereiwiss.* 24: 271-300.
- Unar, M. 1973. A review of the Indonesian shrimp fishery and its present developments. IOFC/DEV/72/27. FAO, Rome.
- Veth, P.J. 1875. *Java, Geographisch, Ethnologisch, Historisch*. Erven F. Bohn, Haarlem: Eerste Deel.
- Wilkinson, R.J., H. Berkeley and H.C. Robinson. 1904. *Report on the fishing industry of the Straits Settlements and Federated Malay States on the West Coast of the Peninsula*. Federal Government Printing Office, Kuala Lumpur.
- Yap, C.L. 1976. Fishery policies and development with special reference to the West Coast of Peninsular Malaysia from the early 1900s. *Kajian Ekon. Malays.* 13(1/2): 7-15.

The Mid-1970s Demersal Resources in the Indonesian Side of the Malacca Strait^a

P. MARTOSUBROTO

Marine Resources Service
Fishery Resources Division
FAO of the United Nations
Viale delle Terme di Caracalla
00100 Rome, Italy

T. SUJASTANI

Deceased, formerly with the
Marine Fisheries Research Institute
Jakarta, Indonesia

D. PAULY^b

International Center for Living
Aquatic Resources Management
MCPO Box 2631, 0718 Makati City
Philippines

MARTOSUBROTO, P., T. SUJASTANI and D. PAULY. 1996. The mid-1970s demersal resources in the Indonesian side of the Malacca Strait [*Sumberdaya ikan demersal di Selat Malaka bagian Indonesia pada pertengahan tahun 1970an*], p. 40-46. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

The results of demersal surveys in the Indonesian half of the Malacca Strait area by the research trawlers *Mutiara 1, 2 and 4* conducted in 1973 and 1975 gave an estimate of the standing stock of 73,000 t. Comparing with other areas, and considering the state of exploitation of the resources in the mid-1970s, the unexploited stock was estimated to have been around 150,000 or 2.66 t·km⁻². Catch rates and catch composition by depth are given.

The commercial trawl fishery in the Indonesian Malacca Strait prevailing in the 1970s was analyzed on the basis of the statistics in provincial reports and of field interviews. Sustainable yields for two distinct fisheries, Aceh and North Sumatra-Riau, were estimated to be 8,000 t·year⁻¹ and 77,000 t·year⁻¹, respectively. The stocks were, at the time, beginning to be overfished.

Abstrak

Hasil survei sumberdaya demersal di Selat Malaka dengan kapal trawl *Mutiara 1, 2, dan 4* yang dilaksanakan pada tahun 1974 hingga 1975 memberikan estimasi kelimpahan ikan sebesar 73.000 t. Dibandingkan dengan daerah lain, dan mengingat tingkat eksploitasi pada pertengahan tahun 1970-an, stok ikan pada tingkat awal diperkirakan 150.000 t atau 2,66 t per km². Hasil tangkapan rata-rata dan komposisi hasil tangkapan disajikan dalam tulisan ini.

Keadaan perikanan trawl di Selat Malaka di sekitar tahun 1970-an dianalisis berdasarkan data statistik propinsi dan kabupaten serta wawancara di lapangan. Potensi lestari perikanan di sekitar dua daerah, Aceh dan Sumatra Utara-Riau, diperkirakan masing-masing sebesar 8.000 t dan 77.000 t per tahun. Saat itu sudah terlihat adanya gejala lebih tangkap.

Introduction

The Malacca Strait (Fig. 1), being one of the world's main shipping routes, is rather well documented in the geological and maritime literature (see Emery 1971; Valencia 1979 and Box 1).

The fisheries of the Malacca Strait are also well documented (SCSP 1976a, 1976b, 1978), although the small pelagics (especially mackerels and scads) have received far more attention than the demersal fishes (c.f. Sujastani 1975, 1976; Anon. 1976a, 1976c, 1976d, 1987 for pelagics; Mansor Mat Isa 1987; Sivasubramaniam 1987; Tampubolon and Sedana Merta 1987; Tampubolon 1988; Soriano et al. 1988 vs Menasveta 1970; Anon. 1976b, 1976e; and Mahyam Binti Mohd. Isa 1988).

^aICLARM Contribution No. 1047.

^bAnother contact address: Fisheries Centre, the University of British Columbia, 2204 Main Mall, Vancouver, B.C. Canada V6T 1Z4; e-mail: pauly@fisheries.com

Q1604; Q1603;
Demersal fisheries
Trawling (Catch/effort
Fisheries (Catch/impact)
ISEW, Malacca Strait
ISEW, Indonesia




Fig. 1. The Indonesian part of the Malacca Strait (shaded), covering an area of about 55,000 km².
 [Gambar 1. Selat Malaka bagian Indonesia (diarsir), meliputi luas sekitar 55.000 km².]

Table 1. Specifications of the research vessels and gear used for trawl surveys on the Indonesian side of the Malacca Strait, 1973 and 1975.
 [Tabel 1. Spesifikasi kapal dan alat yang digunakan dalam survei di Selat Malaka, 1973-1975.]

Name of boat	Gross tonnage	h.p.	Length (m)	Gear	Length of head rope (m)	Mesh size of codend (mm)
<i>Mutiara 1</i>	124	365	23.6	double rig ^a	23.4 (each)	30
<i>Mutiara 2</i>	53.4	165	18.9	double rig ^a	19.0 (each)	30
<i>Mutiara 4</i>	110	316	24.0	Thailand trawl	36.0	40

^aShrimp trawl.

Menasveta (1970) reviewed the bottom trawl surveys conducted in the Malacca Strait, notably the results of trawling by the *R/V Tongkol* (Birtwistle 1928), the *R/V Manihine* (Ommanney 1961), and the *Selayang* (Pathansali et al. 1966). Later surveys were conducted with *K.M. Jenahak* (Latiff 1973). Most of these surveys were conducted in what are now

Box 1. Sediments and benthos of the Malacca Strait.

[Boks 1. Sedimen dan benthos di Selat Malaka.]

Emery (1971), based on Keller and Richards (1967) and other sources, describes the sediments of the Malacca Strait as consisting mainly of "sand (detrital or calcareous grains 0.05 to 2 mm in diameter and having a hard smooth to rippled surface), sand-and-mud (fine sand and silt having a firm to soft smooth surface), mud (silt and clay having a soft surface), gravel (pebbles and cobbles of broken rock, locally containing many calcareous shells), rock (outcrops of bedrock and boulders near outcrops), and coral (large areas or reefs of massive calcareous algae and coral)".

The benthos of the Malacca Strait appears to be sparse, at least offshore, far from the mouth of rivers. Neiman (1973) reports of three benthos stations (see Fig. 2 for their locations) from west to east which yielded densities of 4.4 g⁻² (shallow water) 1.0 g⁻² in (deep water) and 2.4 g⁻² (shallow water). He also mentions that far from the mouths of rivers, otter trawls often catch sponges and soft corals, and sea urchins.

Malaysian territorial waters; less information is available on the Indonesian side of the Malacca Strait.

A first report on catch/effort and catch composition data obtained by two Indonesian fisheries research vessels, *R/V Mutiara 1* and *R/V Mutiara 2*, on the Indonesian stocks of the Malacca Strait, was published in Indonesian by Martosubroto (1973). The station grid of *R/V Mutiara 1* and *2* was not truly random, i.e., fishing was directed. Nevertheless, their catch rates are incorporated in the present paper, itself an updated version of Sujastani et al. (1976).

Anon. (1976e) reported on the composition of 12 trawl hauls taken on the Indonesian side of the Malacca Strait in January 1975 by *R/V Lemuru* (see Venema, this vol., for details on *R/V Lemuru*).

Further data for the present contribution originate from the survey conducted by *R/V Mutiara 4* in the Malacca Strait, in early 1975 (see Pauly et al., this vol. for the context of this survey and the sampling methods used, and Torres et al. (this vol.) for a description of a database with details on these and *R/V Lemuru's* stations).

The fourth source of material used here is the landing and effort statistics

of the Provincial Fisheries Offices of the Provinces of Aceh, North Sumatra and Riau. Additionally, interviews with skippers of commercial trawlers operating in the area were conducted towards the end of 1975 by the second author. Fig. 1 shows the area covered here.

Materials and Methods

Trawl Survey Data

The specifications of *R/V Mutiara 1, 2, and 4* and the gears used for their surveys are summarized in Table 1.

The survey by *Mutiara 1* and *2* in the Malacca Strait lasted from July to September 1973. A total of 148 hauls were made, 61 by *Mutiara 1* and 87 by *Mutiara 2*. The survey of *Mutiara 4* lasted from 27 January to 14 March 1975 and yielded 40 valid hauls (Fig. 2, Table 2). The surface area of each depth horizon is also given in Table 2. These data were used to estimate standing stock size using the swept area method as described in Pauly et al. (this vol.) and using the same assumptions as to the catchability of the gear. Particularly, the same escapement factor was assumed to apply to *Mutiara 1, 2* and *4*.




Fig. 2. Location in the Malacca Strait of trawl hauls performed by *FRV Mutiara 1* and *2* in 1973, by *FRV Mutiara 4* in January/February 1975, by *FRV Lemuru* in January 1975. [Also included are three benthos grab hauls made by *R/V Akademik Knipovich* in 1966]. [Gambar 2. Lokasi stasiun trawl di Selat Malaka oleh kapal penelitian *Mutiara 1* dan *2* pada tahun 1973, kapal *Mutiara 4* pada Januari/Februari 1975, kapal *Lemuru* pada Januari 1975. (Termasuk juga tiga stasiun pengambilan contoh benthos oleh kapal penelitian *Akademik Knipovich* pada tahun 1966).]

Table 2. Haul numbers of *Mutiara 1, 2* and *4* in the Malacca Strait, 1973 and 1975. [Tabel 2. Jumlah hauls/tarikan *Mutiara 1, 2, dan 4* di Selat Malaka, 1973-1975.]

Depth (m)	Surface area (km ²)	No. of hauls by <i>Mutiara</i> :		
		1	2	4
0-9	7,354	2	5	-
10-19	6,860	31	44	-
20-29	4,710	16	38	1
30-39	5,378	12	-	7
40-49	4,719	-	-	11
50-59	7,792	-	-	14
60-69	8,561	-	-	4
70+	10,097	-	-	3
Total	54,931	61	87	40

Box 2. Standardization of trawling effort.

[Boks 2. Pembakuan upaya penangkapan trawl.]

Standardizing the effort of trawlers can be done by relating their catch per effort to their tonnage (as index of "power", since large boats require large engines). The table below documents how the large, medium, and small trawlers were standardized such as to obtain a measure of fishing effort (days fished per year) comparable between provinces.

Vessel characteristic	Large trawlers	Medium trawlers	Small trawlers
Boat class (GT)	> 20	10-20	10
Mean gross tonnage	30	20	10
Average catch (kg-day ⁻¹)	276	218	160 ^a
Power factor	1.27	1	0.74
Nominal effort ^b (by province)			
Aceh	264	-	-
North Sumatra	288	264	240
Riau	280	-	-
Effective effort ^c (by province)			
Aceh	335	(264)	-
North Sumatra	336	264	178
Riau	356	(280)	-

^aAs obtained by linear extrapolation, down to 10 GT, of ratio of catch:tonnage.

^bTrawler days at sea.

^cDays at sea of a standard trawler.

The *FRV Lemuru* data were not used for standing stock estimates owing to the small number of hauls (Fig. 2).

Commercial Catch Data

The fisheries catch data originate from the Provincial Fisheries Offices of the Provinces of Aceh, North Sumatra, and Riau and go back as far as reports were available (to 1969). The second author also conducted field interviews in the three

abovementioned provinces from November to December 1975. In Aceh Province, four large landing places were visited with about three interviews of trawler skippers at each landing site. In North Sumatra Province, landing sites were visited with the same rate of interviews. In Riau Province, only three landing places were visited with four interviews each. Catch per unit of effort data and effort data (actual fishing days) for large and medium trawlers in the three provinces were derived from these interviews.

The provincial reports did not assign the landed fish and shellfish to the gear that caught them. It was thus necessary to identify, from the catch data, those groups of fish and shellfish which could be assumed to originate from trawl catches. This was done on the basis of Table 3 by assuming that fish contributing more than 2% to the survey catches of *Mutiara 1, 2* or *4* were mainly demersal. Those fishes were then selected from the reports' lists and their catches combined to represent the trawler's catches.

When presenting the data, care was taken to avoid double accounting, as occurred between the Province of North Sumatra, where many trawlers from Riau landed their catch, and the latter province. As no figure was available on the amount of this double accounting, a figure of 50% was assumed; accordingly, the catches reported from Riau Province were cut by half.

Annual catch was plotted against the corresponding levels of effort for the Malacca Strait as a whole and for Aceh Province and North Sumatra-Riau Provinces.

Table 3. Demersal fish catch composition (% of weight) in the Malacca Strait by research vessel and depth range.
[Tabel 3. Komposisi hasil tangkapan sumber daya demersal (berat dalam %) di Selat Malaka berdasarkan kapal dan kedalaman.]

Depth range (m) Group or species/ <i>Mutiara</i>	0-9		10-19		20-29			30	-39	40-49	50-59	60-69	70+
	1	2	1	2	1	2	4	1	4	4	4	4	4
Sharks and rays	0.2	0.4	0.7	1.7	3.1	0.3	3.3	1.1	0.5	15.7	1.4	1.0	0.9
Engraulidae	0.2	0.1	1.4	2.6	1.4	1.4	-	0.8	1.7	1.2	0.1	8.0	-
Clupeidae	1.7	4.9	1.1	4.1	0.8	2.5	41.9	0.1	3.9	4.1	3.9	12.8	0.7
<i>Chirocentrus dorab</i>	0.6	1.0	4.1	1.6	2.9	1.6	-	1.8	0.1	0.4	0.1	0.1	0.8
<i>Anadontostoma chacunda</i>	-	-	0.3	0.1	0.1	-	-	0.1	-	-	-	-	-
Synodontidae	0.4	1.5	2.3	5.3	1.8	3.9	-	9.8	2.0	0.5	2.6	2.4	2.8
Ariidae	-	0.7	0.9	0.4	2.6	0.5	-	2.9	0.7	4.3	1.8	7.0	0.2
<i>Sphyaena</i> spp.	0.3	0.3	0.6	0.5	1.2	0.6	-	0.7	6.9	1.5	0.2	-	-
Serranidae	-	0.1	0.5	0.7	1.5	0.1	-	1.9	0.4	0.1	2.1	-	0.6
Terapontidae	0.9	6.4	2.2	3.8	10.4	1.9	0.9	9.0	0.9	0.3	1.1	2.6	0.3
Priacanthidae	-	-	0.1	0.1	1.4	0.1	-	0.9	0.5	1.3	2.0	0.3	0.3
<i>Rachycentron canadus</i>	-	-	0.1	0.5	-	-	-	-	-	0.4	0.1	-	-
Carangidae ^a	1.9	8.3	4.3	6.6	3.5	4.6	4.4	3.2	16.5	11.5	6.2	7.3	15.5
<i>Formio niger</i>	-	0.2	0.3	0.3	0.4	0.1	-	0.1	0.5	0.2	0.1	0.1	0.9
Lutjanidae (incl. <i>Caesio</i> spp.)	-	-	1.0	0.5	0.9	0.8	-	2.4	1.3	5.4	14.4	8.4	.9
Nemipteridae ^b	-	-	2.1	2.8	1.8	3.5	-	2.8	7.5	5.6	6.4	6.8	3.5
Leiognathidae	0.5	7.5	26.8	5.1	20.7	12.6	17.7	8.8	2.9	8.4	7.8	0.3	0.3
Gerreidae	0.9	1.6	1.3	0.6	0.8	0.6	-	1.1	-	2.4	9.5	0.6	17.6
Pomadasyidae ^c	10.9	6.9	5.8	1.7	5.9	1.6	-	4.6	3.9	7.5	4.3	19.4	4.9
Sciaenidae	15.7	11.2	4.4	10.0	3.8	8.8	0.9	4.5	-	4.2	0.9	7.7	-
Mullidae	-	0.9	3.3	2.4	4.5	5.1	-	8.9	6.4	10.7	10.0	1.3	22.9
<i>Siganus</i> spp.	3.4	-	3.9	-	1.9	-	-	1.0	2.1	1.1	0.1	-	0.1
Trichiuridae	1.4	0.9	4.0	5.4	2.5	3.0	8.8	6.1	2.2	3.3	8.8	6.5	1.4
<i>Rastrelliger</i> spp.	1.4	-	0.2	0.6	0.1	-	-	-	-	0.2	1.3	-	-
<i>Scomberomorus</i> spp.	0.3	0.8	0.9	0.4	1.0	0.7	-	1.3	-	-	-	-	-
<i>Pampus argenteus</i>	-	3.1	0.5	0.9	0.2	0.8	9.9	0.1	2.0	0.2	-	0.2	-
Platycephalidae	0.5	2.3	0.5	0.4	1.5	-	-	2.7	0.1	0.1	0.1	-	-
<i>Psettodes erumei</i>	-	0.3	2.1	0.9	0.8	0.8	-	1.3	0.3	0.1	0.9	0.6	0.3
Other flatfishes	13.2	8.9	1.7	4.1	2.5	1.9	0.4	1.9	-	0.1	0.1	0.2	-
Other fishes	45.4	21.9	19.7	26.3	17.3	28.9	4.4	18.4	24.5	7.4	6.5	4.1	3.7
Loligoidae	-	1.3	-	1.8	1.3	2.5	0.4	0.3	10.1	3.6	5.3	2.1	3.7
Sepiolidae	-	1.3	0.9	2.2	1.3	0.9	-	0.1	0.8	0.5	0.8	0.3	0.2
Penaeidae	1.0	1.0	0.2	5.0	0.4	7.8	4.0	0.2	0.1	0.1	-	0.6	-
<i>Thenus orientalis</i>	-	-	0.3	0.1	0.7	-	-	1.1	0.1	0.1	0.2	0.1	-
Other crustaceans	-	1.7	1.0	1.0	0.3	0.6	2.6	-	0.1	0.1	-	0.1	-

^aExcluding *Decapterus* spp.

^bIncluding *Scolopsis* spp.

^cIncluding *Plectorhynchus* spp.

Commercial Effort Data

The official reports on the number of boats did not state whether these were trawlers or seiners. It was assumed that the large, medium and small motorized boats were trawlers, and that none of the sailing boats were engaged in trawling. The small motorized boats in Riau Province were mainly gillnetters, and were thus not included; neither were some unregistered motorized boats in North Sumatra operating as purse seiners.

On the basis of the field data, it appears that the large trawlers had an average gross tonnage of 30 t, the medium trawlers of 20 t, and the small trawlers of 10 t (Box 2). The medium-sized trawlers were selected as standard, and the other types converted to this by means of their catch per unit effort (Box 2).

Nominal fishing days for each type of trawler in each province were then multiplied by the appropriate power factor,

which led to a measure of effective effort ("corrected fishing days").

Combined with the abovementioned catch data, these estimates of effort allowed the parametrization, via plots of catch/effort vs effort, of surplus production models, subject to the caveats in Martosubroto (this vol.).

Results and Discussion

The area covered the surveys of *Mutiara 1* and *2*, and the trawl haul stations of *Mutiara 4* and *Lemuru* are shown in Fig. 2. The composition of the catch of *Mutiara 1* and *2* is given in Table 3 together with that of *Mutiara 4*. To facilitate comparison, all figures were converted into percentages of total catch. For the same reason, those groups of fishes recorded on only one or two of the boats have been added to "other fishes".

Table 4. Mean fish stock densities by depth in the Indonesian sector of the Malacca Strait.
 [Tabel 4. Rata-rata densitas stok ikan pada berbagai daerah di paparan Sunda.]

Depth (m)	Mean catch/effort (kg·hour ⁻¹) of <i>Mutiara</i>			Biomass (t·km ⁻²) based on <i>Mutiara</i> :			Mean
	1	2	4 ^a	1	2	4 ^a	
0-9	36.7	65.3	-	0.5	1.4	-	0.9
10-19	77.8	45.4	-	1.1	1.0	-	1.0
20-29	77.7	41.0	45.3	1.1	0.9	0.7	0.9
30-39	68.4	-	63.9	1.0	-	1.0	1.0
40-49	-	-	158.2	-	-	2.5	2.5
50-59	-	-	100.1	-	-	1.6	1.6
60-69	-	-	84.1	-	-	1.3	1.3
70+	-	-	94.4	-	-	1.5	1.5

^aMean values for the surveys documented in Pauly et al. (this vol.).

A feature discussed by Menasveta (1970), the increase with depth of the percentage of more valuable fish groups was confirmed for the Lutjanidae, Serranidae, Carangidae, Pomadasyidae, and others. Less valuable groups such as the Leiognathidae, Siganidae, Sciaenidae, Platycephalidae, and "other fishes" decreased with increasing depth.

Table 4 summarizes the estimates of standing stock, by depth range. Overall, a standing stock of about 80,000 t was estimated for the Indonesian waters of the Malacca Strait, as delimited in Fig. 1., but excluding the depth range 0-9 m, i.e., for a reference area of 47,600 km². This corresponds to a mean stock density of 1.67 t·km⁻². Here, however, the value for the depth range 10-19 m, where the *Mutiara 4* had not been fishing, was assumed to correspond to the relatively high average value of the same depth range from other, mostly unexploited areas covered by *Mutiara 4* (see Pauly et al., this vol.).

In the specific case of the Malacca Strait, this assumption was probably too optimistic, as heavy trawling was already taking place in the shallow waters. Thus, to correct this figure, the stock densities obtained from the catch rates of *Mutiara 1* and 2 were also calculated (Table 4). These estimates were then used for the depth ranges missing in the *Mutiara 4* survey, while overlapping data of the three boats were averaged. The results are also presented in Table 4, from which an average weighted mean stock density of 1.33 t·km⁻² was derived (including the depth range 0-9 m). The corrected stock estimate, thus, amounts to 73,000 t, over a surface area of 55,000 km².

The value of 1.33 t·km⁻² and the corresponding standing stock are very low when compared with the data from other areas of the Sunda Shelf (see Pauly et al., this vol.), where the average stock density was 2.66 t·km⁻², exactly double the Malacca Strait figure. The low density in the Malacca Strait is




Fig. 3. Trawler catch/effort vs effort on the Indonesian side of the Malacca Strait, 1969-1975; note decline, for both subareas considered here (Aceh and North Sumatra-Riau), and consequently, for the area as a whole (see also Table 5).

[Gambar 3. Hubungan antara hasil tangkapan per upaya penangkapan dan upaya penangkapan di Selat Malaka bagian Indonesia, 1969-1975; perhatikan grafik penurunan untuk kedua daerah (Aceh dan Sumatra Utara-Riau) dan selanjutnya untuk seluruh daerah (lihat juga Tabel 5).]

most probably the result of the existing trawl fishery, which is backed by the fact that it is mainly the shallow waters (down to 39 m) of the Malacca Strait (where almost all, if not all of the trawling takes place) that the stock densities differ from those of other areas. It seems, therefore legitimate to assume, as a first approach, that the unexploited standing stock, before the onset of the trawl fishery, was also double the stock observed in the mid-1970s, or about 150,000 t.

Based on the plots of catch/effort vs effort (Fig. 3), maximum sustainable yield (MSY) was estimated at about 8,000 t·year⁻¹ for Aceh Province, and 77,000 t·year⁻¹ for North Sumatra and Riau combined. Overall, an MSY yield of 85,000 t·year⁻¹ was estimated for the Indonesian waters of the Malacca Strait (Fig. 4).

The above figure of 85,000 t·year⁻¹ is compatible with an unexploited stock estimate (B_0) of 150,000 t (see above), i.e., implying that $MSY \approx M \cdot \frac{1}{2} B_0$ (Gulland 1971), with M set at 1




Fig. 4. Trawler catch vs effort on the Indonesian side of the Malacca Strait, 1969-1975. Note that effort was near optimal in the mid-1970s for the area as a whole, and for North Sumatra-Riau, but not for Aceh province, where effort was then already excessive.

[Gambar 4. Hubungan antara hasil tangkapan dan upaya penangkapan di Selat Malaka bagian Indonesia, 1969-1975. Perhatikan upaya yang mendekati optimal pada pertengahan tahun 1970an untuk seluruh daerah, dan juga untuk Sumatra Utara-Riau, tetapi tidak untuk propinsi Aceh, karena upaya penangkapan sudah melebihi optimal.]

year⁻¹, a value commonly used for the multispecies demersal stocks of Southeast Asia (see Pauly, this vol.). Further, the standing stock estimate of 73,000 t \approx 1/2 B₀, as should occur when fishing effort is at the level generating MSY (Gulland 1971; Ricker 1975; Pauly 1984). Thus, it can be concluded that, overall, the level of trawling effort applied in the mid-1970s to the demersal stocks of the Indonesian side of the Malacca Strait was about right - except in very shallow waters, where the small shrimps attracted an excessive amount of effort. Sujastani et al. (1976), based on these findings, presented a set of practical recommendations for managing the fisheries. [These recommendations, not recalled here, became irrelevant when trawling was banned in Western Indonesia (Sardjono 1980, and see Martosubroto, this vol.); a move which also placed constraints on field sampling with trawls by MFRI scientists, due to strong objections by small-scale fishers].

Rather, we point at the compatibility of the results presented here, which as in the case of the Java Sea (Martosubroto, this vol.) integrate fisheries and survey data into a coherent whole.

Acknowledgements

The kind assistance of the fisheries officers of the Provincial Fisheries Offices in Aceh, North Sumatra, and Riau, is here thankfully acknowledged. Thanks are also due to the

late Mr. M. Unar, Director of the MFRI and to Mr. Budihardjo, who assisted during the field work, and with the compiling of the catch and effort data.

References

- Anon. 1976a. History and development of the purse seine fishery of Peninsular Malaysia, p. 64-66. *In* Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March to 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Anon. 1976b. History and development of fish trawling on the west coast of Peninsular Malaysia, p. 61-63. *In* Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March to 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Anon. 1976c. The Thai fisheries along the Indian Ocean coast of Thailand, p. 67-75. *In* Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March to 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Anon. 1976d. Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March to 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Anon. 1976e. *R/V Lemuru* Cruise 7501 (Jakarta - Belawan, 8-27 Jan. 1975). Data and observations. Presented at the Workshop on the Fisheries Resources of Malacca Strait, 29 March - 2 April 1976, Jakarta, Indonesia. WFMS/76/6/WP. South China Sea Fisheries Development and Coordinating Programme, Manila.

- Anon. 1987. Investigations on the mackerel and scad resources of the Malacca Straits. BOBP/REP/39, 149 p. Bay of Bengal Programme, Colombo, Sri Lanka.
- Birtwistle, W. 1928. Report of the working of the *S.T. Tongkol* for the year 1927. Government Printing Office, Singapore.
- Emery, K.D. 1971. Bottom sediment map of Malacca Strait. UN ECAFE CCOP Tech. Bull. (3):149-152.
- Gulland, J.A. Editor. 1971. The fish resources of the ocean. FAO/Fishing News Books, Ltd., Surrey, England.
- Keller, G.H. and A.F. Richards. 1967. Sediments of the Malacca Strait, Southeast Asia. *J. Sedimentary Petrology* 37:102-127.
- Latiff, S.S. 1973. Resource survey and problems of fishery resource management. Annual Meeting of Fisheries Officers, March 1973. 17 p. (mimeo).
- Mahyam Binti Mohd. Isa. 1988. Population dynamics of *Nemipterus japonicus* (Pisces: Nemipteridae) off Kedah state, Malaysia, p. 126-140. *In* S.C. Venema, J. Moeller-Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO Fish. Rep. 389, 519 p.
 - Mansor Mat Isa. 1987. On the status of the *Rastrelliger* and *Decapterus* fisheries of the west coast of Peninsular Malaysia in 1984-1985, p. 81-100. *In* Investigations on the mackerel and scad resources of the Malacca Straits. BOBP/REP/39. Bay of Bengal Programme, Colombo, Sri Lanka.
 - Martosubroto, P. 1973. Studies on demersal fish stocks in the Malacca Strait and Jambi Bay. *Mar. Fish. Res. Rep.* 1, 33 p. (In Indonesian).
 - Menasveta, D. 1970. Potential demersal fish resources of the Sunda Shelf, p. 525-559. *In* J. Marr (ed.) The Kuroshio: a symposium on the Japan current. East West Center Press, Honolulu.
 - Neiman, A.A. 1973. Shelf benthos in the northern part of the Indian Ocean, p. 52-61. *In* A.S. Bogdanov (ed.) Soviet fisheries investigations in the Indian Ocean (Second cruise of the *R/V Akademik Knipovich*). Israel Program of Scientific Translation, Jerusalem.
 - Ommanney, F.D. 1961. Malayan offshore trawling grounds. The experimental and exploratory fishing cruises of the *F.R.V. Manihine* in Malayan and Borneo waters, 1955-56 with a note on temperature and salinities in the Singapore Strait. *Colon. Office Fish. Publ.* 18:1-95.
 - Pathansali, D., K.S. Ong, S.S. Latiff and J.L. Carvalho. 1966. Preliminary results of trawling investigations off Penang. *Proc. Indo-Pac. Fish. Counc.* 12, Tech. Pap. 7, 22 p.
 - Pauly, D. 1984. Fish population dynamics in tropical waters: a manual for use with tropical waters. *ICLARM Stud. Rev.* 8, 325 p.
 - Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.* 191, 382 p.
 - Saeger, J., P. Martosubroto, and D. Pauly. 1976. First report of the demersal fisheries project (Results of a trawl survey in the Sunda Shelf area). *Mar. Fish. Res. Rep./Contrib. Demersal Fish. Proj.* No. 1, 75 p.
 - Sardjono, I. 1980. Trawlers banned in Indonesia. *ICLARM Newsl.* 3(4):3.
 - SCSP. 1976a. Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March - 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
 - SCSP. 1976b. Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March - 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/6, 113 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
 - SCSP. 1978. Report of the workshop on management of resources of the Sunda shelf, Malacca Strait and related areas. SCS/GEN/78/16, 13 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
 - Sivasubramaniam, K. 1987. Scads and chub mackerels (*Decapterus* spp. and *Rastrelliger* spp.) in the Bay of Bengal, p. 117-145. *In* Investigations on the mackerel and scad resources of the Malacca Straits. BOBP/REP/39. Bay of Bengal Programme, Colombo, Sri Lanka.
 - Soriano, M.L., G. Tampubolon and J. Widodo. 1988. Discriminant analysis of morphometrics of Indian mackerel (*Rastrelliger kanagurta*) in the Malacca Strait and scad (*Decapterus russelli*) in the Java Sea, Indonesia, p. 411-415. *In* S.C. Venema, J. Moeller-Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO Fish. Rep. 389, 519 p.
 - Sujastani, T. 1975. Pelagic fisheries in the Malacca Strait of Indonesia. *Mar. Fish. Res. Rep.* 1:102-117.
 - Sujastani, T. 1976. Pelagic fisheries of Indonesia in the Malacca Strait, p. 48-60. *In* Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March to 2 April 1976, Jakarta, Indonesia. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
 - Sujastani, T., P. Martosubroto and D. Pauly. 1976. A review of the demersal fishery of Indonesia in the Malacca Strait, based on recent surveys and catch and effort data, p. 23-47. *In* Report of the Workshop on the Fisheries Resources of the Malacca Strait, 29 March - 2 April 1976, Jakarta. Part I. SCS/GEN/76/2, 85 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
 - Tampubolon, G.H. and I. Gede Sedana Merta. 1987. Mackerel fisheries in the Malacca Straits, p. 101-116. *In* Investigations on the mackerel and scad resources of the Malacca Straits. BOBP/REP/39. Bay of Bengal Programme, Colombo, Sri Lanka.
 - Tampubolon, G.H. 1988. Growth and mortality estimation of Indian mackerel (*Rastrelliger kanagurta*) in the Malacca Strait, Indonesia, p. 372-384. *In* S.C. Venema, J.M. Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO Fish. Rep. 389, 519 p.
 - Valencia, M.J. 1979. South China Sea: present and potential coastal area resource use conflicts. *Ocean Manage.* 5:1-38.

The *Mutiara 4* Surveys in the Java and Southern South China Seas, November 1974 to July 1976^a

DANIEL PAULY^b

*International Center for Living
Aquatic Resources Management
MCPO Box 2631, 0718 Makati City
Philippines*

PURWITO MARTOSUBROTO

*Marine Resources Service
Fishery Resources Division
FAO of the United Nations
Viale delle Terme di Caracalla
00100 Rome, Italy*

JÜRGEN SAEGER

*#79 Buencamino Street, BF Homes
Parañaque, Metro Manila, Philippines*

Q 1604;
& 1603
Fishery surveys
trawling
Demersal fisheries
catch composition / catch/effort
ISEW, Java Sea

PAULY, D., P. MARTOSUBROTO and J. SAEGER. 1996. The *Mutiara 4* surveys in the Java and southern South China Seas, November 1974 to July 1976 [*Survei Mutiara 4 di Laut Jawa dan bagian selatan Laut Cina Selatan, November 1974 hingga Juli 1976*], p. 47-54. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

This account, which consolidates data from two earlier reports, presents the methodology and major results of a series of Indonesian-German trawl survey cruises conducted in the mid-1970s with *FRV Mutiara 4* in the Java Sea, and the southernmost tip of the South China Sea, to serve as background for detailed analyses of the demersal communities thus sampled.

Abstrak

Makalah ini, yang menggabungkan data dari dua laporan terdahulu, mempresentasikan metode dan hasil utama beberapa kali survei trawl Indonesia-Jerman dalam pertengahan tahun 1970-an dengan kapal penelitian *Mutiara 4* di perairan Laut Jawa dan bagian selatan Laut Cina Selatan, sebagai dasar analisis rinci komunitas sumberdaya ikan demersal.

Introduction

The Indonesian-German Demersal Fisheries Project reported upon here was one of the outcomes of the *Seminar on the Possibilities and Problems of Fisheries Development in Southeast Asia*, which took place in 1968, in Berlin, under the sponsorship of the German Foundation for Developing Countries and FAO (Tiews 1969). In 1972, the Indonesian Government and that of the Federal Republic of Germany agreed to cooperate in carrying out a survey of the bottom fishes in the Java Sea and adjacent areas, to provide the basis for a development of the trawl fisheries similar to the one that

had occurred earlier in the Gulf of Thailand.

This account covers the first operational phase of the project, from late 1984 to mid-1986 (see below for details on the second phase), and is based on Saeger et al. (1976) and Martosubroto and Pauly (1976).

Materials and Methods

Institutional Arrangements and Personnel

The executing agencies were, on the Indonesian side, the Directorate General of Fisheries (DGF), with Mr. V. Suzantc as counterpart Project Manager. On the German side, the executing agencies were "GAWI" and "BFE", both predecessors

^aICLARM Contribution No. 1050.

^b Another contact address: Fisheries Centre, the University of British Columbia, 2204 Main Mall, Vancouver, B.C. Canada V6T 1Z4; e-mail: pauly@fisheries.com

of the German Agency for Technical Cooperation (GTZ), which fielded Dr. J. Saeger as Project Manager.

The project started with the construction of a trawler, the *Mutiara 4* (Box 1, Fig. 1).

The Marine Fisheries Research Institute (MFRI) in Jakarta, where the project was based, provided as biologists Messrs. Purwito Martosubroto, Johannes Widodo, Badrudin, Ir. Achmad Sudradjat, Achmad Basyori, Ir. Isom Hadisubroto, and Toman Panggabean. Mr. Wasilun and Mr. Widji Santosa also participated in the activities of the project.

The counterpart scientists functioned under the general supervision of the late Mr. M. Unar (MFRI Director), Daniel Pauly was assigned by GTZ to the project from May 1975 to

Box 1. The *Mutiara 4*.

[Boks 1. *Mutiara 4*.]

The *Mutiara 4* ("Pearl" in Indonesian) is a wooden stern trawler. She is equipped with a 316-hp main engine (nominal rating), auxiliary engines and echosounders. The fishing winch allows trawling at depths down to 100 m. The cooled fish hold has a capacity of 12 tonnes. A two-drum oceanographic winch allows studies of water properties as well as benthos and plankton sampling. The principal dimensions of *FRV Mutiara 4* are:

Length overall	24.15 m
Length designed WL (=DWL)	23.00 m
Length between perpendicular	21.00 m
Breadth moulded	5.80 m
Breadth overall	5.92 m
Depth moulded a midship	2.87 m
Draft moulded (DWL)	2.23 m
Draft max. aft (DWL)	3.02 m
Drag of keel	1.00 m
Tonnage of keel	111 GRT ^a
Fish hold capacity	45 m ³ approx.
Fuel oil capacity	12.2 t in two tanks
Freshwater capacity	8.92 t
Crew	13 persons (1 captain, 4 officers, and 8 other crew)
Main engine output (t) (tropical conditions)	286 hp at 1,650 Rpm (Deutz B/ 12M716)

The trawler was built at the IPPA Shipyard in Semarang, Central Java, under the supervision of GTZ which provided the abovementioned equipment while the DGF funded the construction of the hull.

^a1 GRT = 100 cubic feet = 2.83 m³.

December 1976. Captain P. Jarchau replaced Captain W. Spiering during his home leaves in 1974 and 1975, and permanently from 1976 on.




Fig. 1. The fisheries research trawler *Mutiara 4* (see also Box 1).
[Gambar 1. Kapal penelitian trawl *Mutiara 4* (lihat juga Box 1).]




Fig. 2. The area surveyed by *Mutiara 4* from November 1974 to July 1976 (excluding the Malacca Strait, treated by Martosubroto et al., this vol.).
[Gambar 2. Daerah survei *Mutiara 4* dari November 1974 hingga Juli 1976 (tidak termasuk Selat Malaka yang dibahas dalam Martosubroto et al., pada volume ini).]

The Survey Area

The survey area covered much of the Sunda Shelf, about 772,500 km², or 238,400 nm², and was divided in six areas as shown on Fig. 2. Area 6, the Malacca Strait, is not reported upon here (but see Martosubroto et al., this vol.).

The distribution of the various depth ranges in the six areas was estimated by drawing isobaths on maps as large as possible, cutting out the figures thus obtained and weighing the paper pieces. The percentage weight of the various depth horizons was then related to the total area. Mangrove and other areas of intertidal vegetation and coralline areas were all included in the 0-9 m depth range. Areas deeper than 70 m were all put in the "70+" category. These results are shown in Table 1 and were used for all subsequent analyses. To avoid the fixed gears of small-scale fishers, no trawling occurred at depths of 0-9 m;

Table 1. Surface of the Sunda Shelf in km² surveyed by *Mutiara 4* from November 1974 to July 1976^a.

[Tabel 1. Luas paparan Sunda dalam km² yang disurvei oleh *Mutiara 4* dari November 1974 hingga Juli 1976^a.]

Depth (m)	Area 1	Area 2	Area 3	Area 4	Area 5	Total
0-9	3,740	6,787	14,161	24,403	10,341	59,432
10-19	2,063	8,118	17,838	28,880	22,960	79,859
20-29	5,932	23,688	24,782	28,880	35,054	118,336
30-39	6,706	14,239	27,097	55,016	54,333	157,391
40-49	13,025	50,836	27,914	7,220	36,105	135,100
50-59	16,378	24,886	15,523	-	13,671	70,458
60-69	42,042	3,593	5,447	-	2,103	53,185
70+	39,076	931	3,268	-	526	43,801
Total	128,962	133,078	136,030	144,399	175,093	717,562

^aExcluding Area 6 (Malacca Strait, treated in Martosubroto et al., this vol).

consequently, this depth range was excluded from all standing stock estimates, as were the stations used for selection experiments (Box 2) for inferences of fish behavior (see Box 5 below), or for obtaining live fish for tagging and aquarium experiments, not reported here.

Trawl Sampling and the Swept Area Method

Throughout the whole survey, a trawl has been used which was similar to the one developed in Thailand at the onset of this country's expanding trawl fisheries expansion, and commonly known as the "Thailand trawl" (Fig. 3). This was

Box 2. Mesh selection experiments.

[Boks 2. Percobaan seleksi mata jaring.]

A selection experiment was conducted on 18 November 1975 close to the harbor of Semarang at depths ranging from 20 to 30 m. For this, the codend of the trawl, with a mesh size of 40 cm (stretched mesh), was covered with a codend of 12 cm (stretched mesh). The results were as follows:

Genus/Species	Number in codend		Mean selection length (TL, cm) ^a
	Inner	Outer	
<i>Rastrelliger kanagurta</i>	318	327	10
<i>Arothron leopardus</i>	119	38	6 ^b
<i>Upeneus</i> spp.	300	806	10.2
<i>Gerres</i> spp.			
<i>Pentapirion longimanus</i>	62	123	8.8
<i>Dussumieria hasselti</i>	136	102	9.5
<i>Saurida undosquamis</i>	63	445	13 ^b
<i>Apogon ceramensis</i>	47	106	8

^aMean length at first capture, i.e., L_c or L₅₀.

^bRough estimates.

Aoyama (1973) published mean selection lengths for 15 species for codend of different mesh sizes. His values and ours can be compared only for *Apogon*, the only fish covered both by his and our experiments. Here, the results correspond nicely, as he gives 8.2 cm mean retention length for a mesh size of 3.8 cm, while we found 8 cm for a mesh size of 4.0 cm.

slightly modified in June 1976 (see Martosubroto, this vol.) presumably without marked impact on catch rates.

All fishing took place between 0500 and 1900 hours, the stations generally following each other on a "station line" of 4-7 hauls (our few oceanographic stations were placed at the




Fig. 3. The "Thailand trawl" and the otter boards used by *Mutiara 4*.

[Gambar 3. Diagram "Thailand trawl" dan otter board yang dipakai *Mutiara 4*.]

Box 3. Species identification and sorting.

[Boks 3. Identifikasi dan sortasi jenis ikan.]

The practice of sorting scientific trawl catches to the family level only was inherited from the previous Thai-German Trawl Fisheries Development Project (see e.g., Ritragosa 1976) which served as model for the project reported upon here. Actually, sorting to species would have been possible - even for what at first appeared a bewildering number of species - given adequate (self)training of all participating biologists.

Available identification tools for shipborne identification included the book of Munro (1967), the FAO identification sheets of Fischer and Whitehead (1974) and an array of family-specific keys, e.g., that of Kühlmorgen-Hille (1968) for the Leionathidae.

In the second phase of this project, from about mid-1976 on, sorting was done to species level, which greatly increased the value of the survey results for stock assessment and biodiversity studies (see also Pauly, this vol. and Bianchi et al., this vol.).

end of such lines). Steaming between the different station lines occurred in the first night hours, after which the ship was anchored until the beginning of the next line.

The station lines were parallel to the predominant current, and similar numbers of trawl stations were carried out with and against that current.

The method for estimating standing stock size is based on the assumption that there is a direct relationship between catch per unit of effort (i.e., catch/hour trawling) and stock density. Here, the area in km² (A) swept by the net in one hour is given by

$$A = 2/3 (HR)(2.8)(1.85)(10^{-3}) \quad \dots 1)$$

where HR is the length of the net's head rope in m (here 36 m), 2.8 is the mean trawling speed of the boat in knots and 1.85 converts nautical miles to kilometers.

Assuming a catchability of 0.5 (Shindo 1973), later confirmed by Pauly (1980) the stock density (D, in t.km⁻²) was determined by

$$D = \frac{\text{kg} \cdot \text{hour}^{-1} (10^{-3})}{A(x_1)} \quad \dots 2)$$

Stations were considered successful when trawling proceeded for one hour without incidents, or when the net was only slightly damaged when retrieved. When trawling had to

Box 4. Observations on zooplankton.

[Boks 4. Pengamatan zooplankton.]

Some zooplankton sampling was done in Area 5, first through vertical tows of a Hansen (fish) larval net; this was not effective, so new collectors were built which could be towed obliquely. Our zooplankton biomass estimates were as follows:

Station ^a	38	44	50	56	62	68	75	82	88	94	100	104
Depth (m) ^b	30	40	38	26	22	16	30	20	32	46	20	20
Water filtered (m ³) ^c	179	327	251	187	200	90	46	60	103	72	64	50
Zooplankton biomass (g.m ⁻³) ^d	0.27	0.14	0.10	0.10	0.08	0.15	0.17	0.15	0.06	0.05	0.13	0.08

^aArea 5 only; see Fig. 4 for locations.

^bLowest depth of oblique tow, as estimated from simultaneous BT records.

^cBased on a calibration experiment conducted in Tanjung Pinang Harbour.

^dWet weight.

These data match closely previous biomass estimates (Anon. 1973) which give values of 0.1 g.m⁻³ for the Java Sea. We also noted large concentrations of jellyfish, when surveying Area 4.

Box 5. Observation on zoobenthos.

[Boks 5. Pengamatan zoobenthos.]

Some benthos samples were taken in Areas 4 and 5 with a van Veen grab whose open jaws covered a surface area of 0.1 m². Generally, 50 % of each bottom sample was kept and preserved in 4% formaldehyde in seawater, and later sorted after sieving over 500 micron meshes. Mollusc shells, found to be regularly empty, were discarded, leaving the polychaetes, mainly errantian, and small crustaceans. Large forms, such as echinoderms or decapod crustaceans never occurred in our grab and are therefore discussed below on the basis of their appearance as trawl by-catch. The results thus obtained were as follows:

Area/ Station ^a	Depth (m)	Biomass (wet weight; g.m ⁻²)		Total *
		Polychaetes	Crustaceans	
4/15	24	1.0	0.2	1.2
4/23	35	0.8	-	0.8
4/31	20	1.0	-	1.0
4/42	35	-	-	-
4/47	34	6.4	0.4	6.8
4/54	13	5.4	0.2	5.6
4/59	35	1.4	0.4	1.8
4/62	32	1.6	0.2	1.8
5/38	36	5.6	0.8	6.4
5/56	42	3.2	-	3.2
5/62	44	2.8	1.6	4.4
5/68	27	5.6	1.0	6.6
5/82	40	1.0	1.4	2.4
5/102	52	5.0	0.4	5.4

^aSee database in Torres et al. (this vol.) for positions, times, etc.

The following observations were made on the macrobenthic bycatch:

Crustaceans: Penaeid shrimps, palinurids, *Thenus orientalis*, and other crustaceans such as brachiurans (crabs) and stomatopods (mantis shrimp) never occurred in commercially relevant concentrations. This is most probably due to the gear used, the depths, and the areas surveyed but it may also indicate an overall scarcity of these resources in the area of the survey. The following penaeid shrimp were found and identified: *Penaeus merguensis*, *P. semisulcatus*, *Metapenaeus monoceros*, *Parapenaeopsis* spp., *Solenocera* spp.

Echinoderms: except for a tendency for large sea-stars to become entangled in the net, and for ophiuroids and echinoideans to make the deck sticky and slippery, the animals of this phylum seem to represent neither a nuisance nor a resource.

Porifera: During coverage of Areas 4 and 5, large mushroom-shaped sponges (tentatively identified as *Poterion nautilus*) occurred in almost every haul; they were counted, and some of them weighed; a rough average of 40 kg per sponge was estimated.

be interrupted, but the haul was nevertheless considered successful, the catch was made comparable to an hourly catch by application of a multiplicative factor. Stations were considered unsuccessful when the net was badly damaged, torn, or when the otter boards (Fig. 2) were not properly set. The position and details on all stations, including the time (Western Indonesian time), the depth, the catch and/or remarks in the case of unsuccessful stations are given in the database documented in Torres et al. (this vol.).

Sorting of the Catch

The catch of each haul was generally sorted to families, or to genus or species in some cases (Box 3). This work was done by the biologists, who were herein kindly assisted by the ship's crew. The catch of each group was weighed with the precision of 0.1 kg. Weighings were done with an arm balance, not very sensitive to the movements of the boat.

Fish specimens were gathered throughout the survey area for the reference collection of the MFRI, and included in mid-1976, over 750 specimens distributed in 80 families, 146 genera and 231 species (Widodo 1976). The corresponding occurrence records will be part of FishBase 97 (see Froese et al., this vol.).

Length-frequency Measurement

A total of over 40,000 fish, belonging to 40 species were measured during the surveys presented here. Most of these measurements are documented in Martosubroto and Pauly (1976) and Pauly and Martosubroto (1980). Pauly et al. (this vol.) present the analyses based on these data, which are also included in the database documented in Torres et al. (this vol.).

Results and Discussion

The measurements of temperature, O₂ and transparency performed in the course of the survey are too scattered to warrant presentation here; they may be found, however, in the database documented in Torres et al. (this vol.). Box 4 documents our observations on the zooplankton in survey area 5, while Box 5 documents the zoobenthos.

Catch Rates, Stock Densities and Standing Stocks

Table 2 shows the numbers of successful hauls in each area and depth horizon (excluding the Malacca Strait, discussed in Martosubroto et al., this vol.). These numbers vary greatly between areas (see also Fig. 4), due to several breakdowns of

Box 6. Back and forth fishing experiment.

[Boks 6. Percobaan penangkapan "maju-mundur."]

During the coverage of Area 5, an experimental depletion of the fish stock by means of repeatedly fishing on the same station was conducted, based on a suggestion by Captain P. Jarchau, whose help largely contributed to its relative success. This was to provide information on:

- i) how fast an area swept by the trawl is "resetted", hence on how close two trawlers might follow each other when operating; and
- ii) the possibility of separating fish groups into "sedentary" and "motile", hence on whether the thinning out of certain fish groups in a given area will affect the nearby areas.

Nine hauls were made by trawling back and forth between two buoys anchored with a distance of approximately 2.3 nm from each other. The average trawling time was 49 minutes, and all catches were adjusted to the standard of one hour's trawling. The experiment was carried out at the geographical position of Station 110, in Area 5 (i.e., 0°47.6'N; 104°27.0'E).

Six groups, assumed to represent the main behavioral type of "sedentary" and "motile" fishes were selected. Of these, only two showed a decreasing trend over the nine hauls: the sharks, and the rays; the latter of these can safely be assumed to represent the most sedentary of the group below:

Group	Successive densities (kg-hour ⁻¹)								
	Haul number								
	1	2	3	4	5	6	7	8	9
Sharks	10.1	17.5	7.5	10.1	4.1	4.7	2.0	4.8	3.9
Rays	45.9	48.8	13.2	44.1	20.9	22.7	29.7	20.8	14.3
Clupeidae/Engraulidae	1.5	5.0	0.1	0.8	0.1	0.4	0.2	0.4	2.4
<i>Chirocentrus dorab</i>	0.5	9.1	6.0	1.1	1.4	1.4	0.9	12.0	4.5
Ariidae	0.9	2.0	1.8	2.1	0.2	0.3	0.9	0.7	1.4
Leiognathidae I	61.7	94.1	32.2	6.0	-	-	-	-	-
Leiognathidae II	-	-	-	-	45.3	249.2	50.0	73.1	26.0

The pelagics (Engraulidae/Clupeidae, *Chirocentrus dorab*) were not affected by the back and forth fishing. The Ariidae, surprisingly, also appeared to be very "motile" although they are benthic fishes.

The catch data for the Leiognathidae (*Leiognathus splendens* for more than 90%) first decreased markedly, then rapidly increased, suggesting that a large school entered the fished zone. This school was then depleted by subsequent fishing.

the ship's main engine. Still, these hauls allowed mapping the catch rates over most of Areas 1-5 (Fig. 5).

The estimates of overall stock density obtained by the swept area method are given in Table 3, by area and depth range. The highest stock density of 5.2 t·km⁻² occurred in Area 1, between 50 and 59 m, while the lowest density of 0.8 t·km⁻² occurred, as expected, in the heavily exploited shallow waters off Western Java, i.e., in Area 2 (see also Martosubroto, this vol.).

Table 2. Number of successful hauls by *Mutiara 4*, November 1974 to July 1976^a.

[Tabel 2. Jumlah tarikan trawl yang sukses dari *Mutiara 4*, November 1974 hingga Juli 1976.]

Depth (m)	Area 1	Area 2	Area 3	Area 4	Area 5
10-19	-	1	17	5	13
20-29	-	29	22	25	24
30-39	-	11	11	30	48
40-49	3	24	14	1	50
50-59	17	35	15	-	9
60-69	30	4	6	-	3
70+	-	1	-	-	-
Total	50	105	85	61	147

^aExcluding Area 6 (Malacca Strait, treated in Martosubroto et al., this vol.).




Fig. 4. The station grid of Mutiara 4, November 1974 to July 1976 (excluding the Malacca Strait, treated by Martosubroto et al., this vol.).

[Gambar 4. Stasiun pengambilan contoh kapal Mutiara 4, November 1974 hingga Juli 1976 (tidak termasuk Selat Malaka, dibahas dalam Martosubroto et al., this vol.).]




Fig. 5. Distribution of catch rates ($\text{kg}\cdot\text{hour}^{-1}$ of *Mutiara 4* November 1974 to July 1976 (excluding the Malacca Strait, treated by Martosubroto et al., 1976).
 [Gambar 5. Sebaran rata-rata hasil tangkapan per jam tarikan trawl dari *Mutiara 4*, November 1974 hingga Juli 1976 (tidak termasuk Selat Malaka, di-

Table 3. Mean density (in t·km⁻²) of demersal fish as estimated by the swept area method for the period from November 1974 to July 1976^a.

[Tabel 3. Estimasi kepadatan rata-rata (dalam km²) sumberdaya ikan demersal berdasarkan metode dalam kurun waktu November 1974 hingga Juli 1976^a.]

Depth (m)	Area 1	Area 2	Area 3	Area 4	Area 5
10-19	(3.0)	0.8	2.1	4.8	3.4
20-29	(2.5)	2.7	2.4	4.3	2.8
30-39	(1.9)	2.4	2.5	2.1	2.3
40-49	3.2	2.2	2.2	1.4	1.8
50-59	5.2	1.9	1.6	-	1.4
60-69	3.4	1.9	1.5	-	1.2
70+	(1.5)	1.5	-	-	(1.5)
Biomass ^b	362	272	259	388	389

^aExcluding Area 6 (Malacca Strait, treated in Martosubroto et al., this vol.).

^bIn t·10³.

From the density data, the standing stock (biomass) for each area and depth range was calculated, based on the surface area of the various depth ranges in Table 1. We abstain here from presenting more than the sum of these estimates, by area (Table 3, last row).

In Saeger et al. (1976), "Gulland's equation" i.e., (Potential yield = 0.5·M· unexploited biomass; Gulland 1971) was used to derive potential yield estimates from the biomasses in Table 2, and M = 1 year⁻¹. Then, the number of trawlers was estimated which would be required to harvest this yield, and rough estimates of the value of their potential catch were presented - the point being to assemble all the elements required for assessing the prospects for the development of trawl fisheries.

The demersal fisheries did develop - so fast, indeed, that trawling was banned a few years later (Sardjono 1980). Pauly (this vol.) discusses this, and the new uses to which the data we gathered (documented in Torres et al., this vol.) can be put; Pauly et al. (this vol.) discuss the biology of some of the fish species surveyed by *Mutiara 4* and other research vessels, while Martosubroto (this vol.) and Bianchi et al. (this vol.) discuss the structure of the assemblages formed by these fishes.

Acknowledgements

The Indonesian-German Demersal Fisheries Project was based on suggestions made by Prof. Dr. K. Tiews. Thanks are due to him for his guidance and encouragement during the course of the work.

The work of the project was greatly enhanced by the close cooperation with Mr. V. Susanto and Mr. Soewito and the staff of the DGF, and with Mr. M. Unar, and Mr. Mangundjojo of MFRI, to whom our thanks are extended. We also take this opportunity to thank Messrs. J. Widodo, Badrudin, Sudradjat, Hadisubroto, A. Basyori, and T. Panggabean for their enthusiasm.

The Captains, W. Spiering and P. Jarchau, and Mr. A. Adli, Co-Captain and their crew substantially contributed to the success of the various trips of the *Mutiara 4*.

References

- Anon. 1973. Report of the exploratory operations of trawl in the Gulf of Siam and Java Sea. Fish. Res. Dev. Agency, Office of Fisheries, Busan, Korea.
- Aoyama, T. 1973. The demersal fish stocks and fisheries of the South China Sea. IPFC/SCS/DEV/73/3, 80 p. Rome.
- Fischer, W. and P.J.P. Whitehead, Editors. 1974. Identification sheets for fishery purposes. Eastern Indian Ocean (Fishery Area 57) and Western Central Pacific (Fishing Area 71), Vols. I-IV, FAO, Rome.
- Gulland, J.A., Editor. 1971. The fish resources of the oceans. Fishing News (Books) Ltd., West Byfleet, England. 255 p.
- Kühlmorgan-Hille, G. 1968. An illustrated field key to the fish family Leiognathidae in the Gulf of Thailand. Contrib. Mar. Fish. Lab., Bangkok 12, 7 p.
- Martosubroto, P. and D. Pauly. 1976. *R/V Mutiara 4* survey data November 1974 to July 1976. Mar. Fish. Res. Rep. (Special Report)/Contribution of the Demersal Fisheries Project No. 2: 1-135.
- Munro, I.S.R. 1967. The fishes of New Guinea. Department of Agriculture, Stock, and Fisheries. Port Moresby, New Guinea. 651 p.
- Pauly, D. 1980. A new methodology for rapidly acquiring information on tropical fish stocks: growth, mortality and stock-recruitment relationships, p. 154-172. In P. Roedel and S. Sails (eds.) Stock assessment for tropical small-scale fisheries. International Center for Marine Resources Development, University of Rhode Island, Kingston.
- Pauly, D. and P. Martosubroto. 1980. The population dynamics of *Nemipterus marginatus* off western Kalimantan, South China Sea. J. Fish Biol. 17: 263-273.
- Ritragas, S. 1976. Results of the studies on the status of demersal fish resources in the Gulf of Thailand from trawling surveys, 1963-1972, p. 198-223. In K. Tiews (ed.) Fisheries resources and their management in Southeast Asia. German Foundation for International Development, Berlin.
- Saeger, J., P. Martosubroto and D. Pauly. 1976. First report of the Indonesian-German Demersal Fisheries Project (Report of a trawl survey in the Sunda Shelf area). Lap. Penel. Per. Laut. (Special Report)/Contribution of the Demersal Fisheries Project No. 1: 1-46.
- *Sardjono, I. 1980. Trawlers banned in Indonesia. ICLARM News. 3(4):3
- Shindo, S. 1973. General review of the trawl fishery and the demersal fish stock of the South China Sea. FAO Fish. Tech. Pap. 120, 49 p.
- Tiews, K., Editor. 1969. Possibilities and problems of fisheries development in Southeast Asia. German Foundation for International Development, Berlin.
- Widodo, J. 1976. A check-list of fishes collected by *Mutiara 4* from November 1974 to November 1975. Lap. Pen. Per. Laut. Special Report/Contribution of the Demersal Fisheries Project No. 1: 47-77.

Demersal Assemblages of the Java Sea: A Study Based on the Trawl Surveys of the *R/V Mutiara 4*

GABRIELLA BIANCHI

Institute of Marine Research
Division of International Development Programmes
P.O. Box 1870
5024-Bergen, Norway

M. BADRUDIN and SUHENDRO BUDIARDJO
Research Institute for Marine Fisheries
Agency for Agricultural Research and Development
Komplek Pelabuhan Perikanan Samudra
Jl. Muara Baru Ujung, Jakarta 14440, Indonesia

Q1604
Demersal fisheries
Abundance
Geographical distribution
Seasonal variations ✓
Fishery surveys
Trawling
ISEW, Java Sea

BIANCHI, G., M. BADRUDIN and S. BUDIARDJO. 1996. Demersal assemblages of the Java Sea: a study based on the trawl surveys of the *R/V Mutiara 4* [Kelompok sumberdaya demersal Laut Jawa: suatu studi berdasarkan survei trawl dengan kapal *Mutiara 4*], p. 55-61. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

An attempt is made to analyze the data collected through the trawl surveys conducted in the Java Sea by *R/V Mutiara 4* to detect patterns of species associations and of seasonal variation in abundance. The analyses based on the data collected through the first survey period (1974-1976), by large categories and including an undetermined number of species, could - at the family level - only confirm the homogeneity of the area surveyed. Data from the second survey period revealed differences at the species level, allowing the division of the Java Sea communities into at least three major groups: assemblages of the central and of deepest part of the basin (depth >30 m) and assemblages of the shallow coastal waters, where major differences appear to be due to the presence of estuaries. Seasonality is another important structuring factor: southeast monsoon catches were over two times higher than northwest monsoon catches, and less variable.

Abstrak

Analisis terhadap data yang terkumpul dari survei trawl di Laut Jawa oleh kapal penelitian *Mutiara 4* dilaksanakan dalam rangka melihat bentuk asosiasi spesies dan variasi musiman dari kelimpahannya. Analisis terhadap data yang dikumpulkan dalam periode survei pertama (1974-1976), berdasarkan pengelompokan kasar termasuk beberapa spesies yang belum teridentifikasi, - pada tingkatan famili - hanya mampu memperkuat adanya keseragaman jenis di daerah survei. Data dari periode kedua menunjukkan adanya perbedaan pada tingkat spesies yang mana hal tersebut dapat dipakai untuk membagi komunitas di Laut Jawa menjadi tiga kelompok besar: yakni kelompok di daerah pertengahan dan daerah yang dalam (kedalaman > 30 m), serta kelompok di daerah yang dangkal, dimana perbedaan tersebut adalah akibat pengaruh estuari. Pengaruh musim juga merupakan faktor penting: hasil tangkapan pada waktu musim angin tenggara tercatat dua kali lebih tinggi daripada waktu musim angin barat laut, dan memiliki fluktuasi hasil tangkap yang lebih kecil.]

Introduction

The present study utilizes data collected by the *R/V Mutiara 4* through the survey program started by the Indonesian-German Demersal Fisheries Development Project in 1974. The surveys aimed to evaluate the abundance and distribution of the demersal fishes of Indonesia. They took place in two survey periods: one between November 1974 and June 1976, devoted to a single coverage of Western Indonesia and another in June 1976-1979, which consisted of several coverages of the Java Sea. Various cruise reports were

produced, including descriptions of the data collected and analyses of catch composition, biomass estimates and potential yields (Martosubroto and Pauly 1976; Saeger et al. 1976; Losse and Dwiponggo 1977; Dwiponggo and Badrudin 1980). Pauly et al. (this vol.) presented an account of the methodology and main results obtained from the surveys of the *R/V Mutiara 4* in 1974-1976 while Martosubroto (this vol.) analyzed the changes that have occurred in the composition of the demersal resources of the Java Sea in the period 1975-1979. This study follows up on this with an attempt to identify the major fish assemblages of the Java Sea, and to identify some of the environmental factors shaping them.

Table 1. Survey effort by region and period (arranged chronologically). NW: northwest monsoon; SE: southeast monsoon.
 [Tabel 1. Kegiatan survei menurut daerah dan waktu (disusun secara kronologis). NW: musim angin barat-laut; SE: musim angin tenggara.]

Dates	Season	From	To	No. of stations
Survey period: November 1974 to June 1976 ^a				
Central and East Java (Leg I)	NW	17.11.74	18.12.74	56
South China Sea (Leg V)	NW	31.01.75	02.02.75	13
Malacca Strait (Leg VI)	NW	09.02.75	21.02.75	41
Central and West Java (Leg II)	SE	29.04.75	26.06.75	112
South China Sea (Leg V)	SE	09.08.75	16.09.75	147
Karimata Strait (Leg IV)	SE	07.11.75	11.12.75	62
South Kalimantan (Leg III)	NW	19.03.76	22.06.76	80
Survey period: July 1976 to December 1979 ^b				
East Java Sea (Areas H & G)	SE	15.06.76	24.06.76	25
East Java Sea (Areas H & G)	SE	21.07.76	29.07.76	23
Central Java Sea (Areas E, F, C, D)	SE	04.08.76	13.08.76	28
Central and West Java Sea (Areas A, B, E, C)	SE	03.09.76	10.09.76	30
West Java Sea	SE	17.09.76	22.09.76	19
Central and East Java	NW/SE	20.01.77	12.10.77	112
Central and East Kalimantan	SE	14.10.77	07.11.77	37
Central, East and West Java	NW	28.11.77	21.12.77	82
Central Java	NW	10.01.78	11.01.88	7
Southeast Sumatra	NW	13.01.78	19.01.78	29
Central and East Java	SE	19.05.78	24.11.78	99
East Kalimantan	NW	27.12.78	01.01.78	46
West and Central Java	NW	12.01.79	19.01.79	44
Southeast Sumatra	NW	02.02.79	07.02.79	26
Central, East and West Java	NW	07.02.79	24.05.79	108
West Kalimantan	SE	14.06.79	19.06.79	30
Southeast Sumatra	SE	21.07.79	25.07.79	27
East Kalimantan	SE	10.08.79	13.08.79	20
East and Central Java	SE	05.09.79	10.09.79	37
Central Kalimantan	SE	20.09.79	25.09.79	33
Central Java	SE	12.10.79	18.10.79	41
Southeast Sumatra	SE	25.11.79	26.11.79	6

^aIncluding stations considered unsuccessful.

^bExcluding gear tests and aimed fishing.

Materials and Methods

Table 1 presents an overview of the surveys by *R/V Mutiara 4* during the period 1974-1979, by region and season. The area covered by the project is presented in Pauly et al. (this vol.) and Martosubroto (this vol.) for the periods 1974-May 1976 and June 1976-1979, respectively. Fig. 1 presents the position of the trawl stations for the latter period. From 1974 to May 1976, the southern part of the South China Sea, and the Malacca and Karimata Straits were also surveyed besides the Java Sea, while the area of operation was later limited to the Java Sea. The sampling design and the taxonomic levels used to identify the catches varied between the two survey periods. From 1974 to 1976, identifications were carried out at family level, including about 30 families, in addition to such broad groupings as "sharks," "rays" or "trashfish". From mid-1976 on, fish identification was much improved and species categories were included, leading to a total of 117 systematic and other groups (see Table 2 and Pauly et al., this vol.).

The sampling design during the period 1974-1976 consisted of tracks, usually parallel to the coast, with sets of

stations very close to each other (cluster sample approach). The observations collected with this type of sampling design have the drawback of nonindependence and autocorrelation between the cluster of observations, a feature that is problematic when biomasses are to be estimated (Gunderson 1993). This problem was dealt with by using, for our community analyses, only a subset of all stations, obtained by extracting the most distant stations from each track, for the whole area.

The second survey period (June 1976 to 1979) also involved variable survey designs. During the first three months, June-September 1976, a systematic and complete coverage of the Java Sea (Fig. 1) was performed. Successive surveys did not follow an explicit survey design, with stations mainly in shallow coastal areas and "clumped" in some areas as off Southeast Kalimantan and Central Java (Fig. 1). The analysis of community types is thus based on the quasi-synoptic survey of the period June-September 1976. Nonvalid stations, i.e., cases when the gear was damaged, or aimed stations (fishing trials) were excluded from all analyses.

The analysis of seasonal differences in catch rates is based exclusively on stations near Tanjung Selatan, Southern Kalimantan, i.e., from the only area of the Java Sea where

Table 2. Systematic and other groups used for classification of the catches during the *R/V Mutiara 4* surveys. Numbered names indicate the 42 categories used for sorting the catches from 1974 to mid-1976; the other names are the taxa added to the initial list in mid-1976.

[Tabel 2. Kelompok hasil tangkapan (menurut sistematika dan grup lain) dari survei dengan kapal penelitian Mutiara 4. Tercatat ada 42 kategori yang dipakai dalam pengelompokan hasil tangkapan tahun 1974 hingga pertengahan 1976; nama-nama yang tidak pakai nomor adalah yang ditambahkan pada pertengahan 1976.]

1. Sharks	10. Formionidae ^a	21. Pomadasyidae
<i>Carcharhinus sealei</i>	<i>Formio niger</i> ^a	<i>Pomadasys argyreus</i>
Other sharks	11. Gerreidae	<i>Pomadasys hasta</i>
2. Rays	<i>Pentaprion longimanus</i>	Other Pomadasyidae
Dasyatidae	Other Gerreidae	22. Priacanthidae
Other rays	Heterosomata	<i>Priacanthus macracanthus</i>
3. Ariidae	12. <i>Psettodes erumei</i>	<i>Priacanthus tayenus</i>
<i>Arius coelatus</i>	13. Lactariidae	23. Rachycentridae
<i>Arius maculatus</i>	14. Leiognathidae	24. Sciaenidae
<i>Arius thalassinus</i>	<i>Gazza minuta</i>	25. Scombridae
<i>Arius venosus</i>	<i>Leiognathus brevirostris</i>	<i>Rastrelliger</i> spp.
<i>Osteogeneiosus militaris</i>	<i>Leiognathus bindus</i>	<i>Scomberomorus</i> spp.
Other Ariidae	<i>Leiognathus daura</i>	26. Serranidae
4. Balistidae	<i>Leiognathus elongatus</i>	27. Stromateidae
<i>Abalistes stellaris</i>	<i>Leiognathus equulus</i>	<i>Pampus argenteus</i>
5. Carangidae	<i>Leiognathus splendens</i>	<i>Pampus chinensis</i>
<i>Alectis</i> spp.	<i>Leiognathus smithursi</i>	28. Sphyraenidae
<i>Alepes</i> spp.	<i>Secutor insidiator</i>	<i>Sphyraena barracuda</i>
<i>Atule</i> spp.	Other Leiognathidae	<i>Sphyraena jello</i>
<i>Atropus atropus</i>	15. Lutjanidae	Other Sphyraenidae
<i>Carangoides</i> spp.	<i>Lutjanus johni</i>	29. Synodontidae
<i>Caranx</i> spp.	<i>Lutjanus sanguineus</i>	<i>Saurida longimanus</i>
<i>Decapterus</i> spp.	Other Lutjanidae	<i>Saurida micropectoralis</i>
<i>Megalaspis cordyla</i>	16. Mullidae	<i>Saurida undosquamis</i>
<i>Selar</i> spp.	<i>Upeneus bensasi</i>	Other Synodontidae
<i>Selaroides leptolepis</i>	<i>Upeneus sulphureus</i>	30. Terapontidae
<i>Seriolina nigrofasciata</i>	Other Mullidae	31. Trichiuridae
Other Carangidae	17. Muraenesocidae	32. Other foodfish
6. Chirocentridae	18. Nemipteridae	33. Trashfish
7. Clupeidae	<i>Nemipterus bathybius</i>	34. Squids
<i>Anadontostoma chacunda</i>	<i>Nemipterus hexodon</i>	35. Cuttles
<i>Dussumieria acuta</i>	<i>Nemipterus japonicus</i>	36. Shrimps
<i>Ilisha</i> spp.	<i>Nemipterus marginatus</i>	37. Crabs
<i>Sardinella</i> spp.	<i>Nemipterus mesoprion</i>	38. Lobster (<i>Thenus</i> sp.)
Other Clupeidae	<i>Nemipterus peronii</i>	39. Other invertebrates
8. Drepanidae	<i>Nemipterus tolu</i>	40. Snakes
9. Engraulidae	<i>Scolopsis</i> spp.	41. Turtles
<i>Stolephorus</i> spp.	Other Nemipteridae	42. Sponges
<i>Thryssa</i> spp.	19. Pentapodidae	
<i>Setipinna</i> spp.	20. Polynemidae	
<i>Coilia dussumieri</i>		

^aNow put within the family Carangidae (Nelson 1994).

trawl stations covering several years were available for both seasons (southeast monsoon: 1977-1979; northeast monsoon: 1978, 1980 and 1982-1984).

The methods used to identify patterns of species association include multivariate analysis techniques of ordination and classification type. They are the same as those described in Bianchi (this vol; see also McManus, this vol.).

Study Area

The study area includes the Java Sea and a part of its connection to the southern South China Sea, i.e., the southern Karimata Strait. This region is part of the triangle bounded by the Philippines, the Malay Peninsula and New Guinea, characterized by the extraordinary richness of its marine fauna. Although there are a few species unique to this region, the total number is higher than in other similar areas of the world

(Ekman 1953). The reasons for the high diversity are usually related to climatic stability at various time scales and the "effect of area," the availability of large shelf areas with favorable conditions for speciation through eustatic sea level changes (see Sharp, this vol.; McManus, this vol.).

The area studied is mostly shallower than 100 m, with rather stable conditions, although under the influence of the monsoon regime. The southeast monsoon blows from April to November and reaches its peak in June-September. The northwest monsoon blows the rest of the year, with peaks in December-February. The main changes in the oceanographic conditions entail a major westward movement in the Java Sea during the southeast monsoon. Surface salinities vary from 33 to 36 but are noticeably lower near the shores of Sumatra, Kalimantan and Java because of local runoff. Here the waters are also less transparent because of the fine sediments transported with the fluvial outflow. Temperatures vary little from 27 to 29°C (see Roy, this vol.).

During the northwest monsoon (December to February), the general transport is eastwards and low-salinity surface waters from the China Sea enter the Java Sea. The heavy rains and increased runoff from the rivers in Sumatra, Kalimantan and Java contribute to the further decrease in salinity in this season.

Unfortunately, comprehensive coverages for both monsoon seasons were not available; hence the limitation, for the interseason comparison, to a small area of the Java Sea (see above and Fig. 1).

Results

Fig. 2 shows the result of the TWINSPAN analysis based on a subset of stations from the survey period November 1974-May 1976, while Fig. 3 shows the location of the stations of Group 2 generated by this analysis.

Fig. 2 indicates a general uniformity in the composition of the major families in the trawl catches. The main difference identified by the first TWINSPAN division is between the very shallow waters (depth 10 to 25 m) in proximity of river outlets and the remaining stations (depth 25 to about 70 m). All taxa found in the deeper range were also present in shallow waters, except for the family Balistidae. Also, a few taxa appeared to be restricted to shallow areas. These include both small pelagic species (Engraulidae and Clupeidae), and demersal species of the family Sciaenidae and Drepanidae.

The set of 119 stations from the survey period June-September 1976 fully covered the area under study and were sampled within one season, during the southeast monsoon. Figs. 2 and 4 show the results from the TWINSPAN analysis and the plot of the above stations on DCA axes 1 and 2, respectively Fig. 3. The first division resulting from the TWINSPAN analysis shows the presence of two major groups: a shallower, with average depth of 27 m, and a deeper one, with average depth of 43 m. The former group has as indicator species *Dussumieria acuta*, *Ilisha*, Sciaenidae and *Leiognathus splendens*. The other group corresponds to the deeper part of the Java Sea basin. Although many taxa are common with the shallow group, the deepwater stations do not have consistent indicator species mentioned above, or their presence is less consistent.




Fig. 1. Trawl stations of the Java Sea surveys conducted by the *R/V Mutiara 4* from June 1976 to December 1979 (fishing experiments not included); (dotted line - 20 m; dashed line - 40 m; solid line - 200 m). The box around Tanjung Selatan shows the stations used for between-season comparisons of catch rates.

[Gambar 1. Stasiun pengamatan survei Mutiara 4 di Laut Jawa dari Juni 1974 hingga Desember 1979 (tidak termasuk percobaan penangkapan); (garis bertitik - 20 m; garis terputus - 40 m; garis - 200 m). Kotak di sekitar Tanjung Selatan menunjukkan stasiun pengamatan yang dipakai untuk membandingkan hasil tangkapan antara dua musim.]

	Group 1 (> 25 m)	Group 2 (10-25 m)
Balistidae	51322541124223321-111121121-1142-32133--3-2-1-----	---1---4-----1-----
Priacanthidae	--2-13-1122131131-----113221---1124-3-11314123-5--	---13411--12-11-11---1-----1-----
Gerreidae	--121-1-3--1-31--2-----111-11---4211--1231-3355554	--11--51--52432---211-33-1-----
Lutjanidae	44354554255155334231-1321332-3-2112--31151441-3--21	55-41-3-14---1-1-122--1-----
Nemipteridae	2112-54223233234111311-341321-1-21332233334455414142	211143423132--22412-111--1211-2-----
Synodontidae	-11211-112122122111--111111111-111111111-11112321-	1-11112221311-11-1--11111-1111-1----
Serranidae	223-----1-111-1-11-----11-1-----1-----34-----	21--1-1-1-----1-----1-----
Scyllariidae	--1--1--11--111-1-1111-11111-11-1111-11-111-1-1--	-----1-----11-1--1-----1-----
Unidentifed	-13-2311333232343123355455551-33532531331333--2--	221511225--3-13251-1-1--111211-25511
Pomadasyidae	5-41-4554234554-542--211-233--2233--2324444313--	542214224-1--221--2-2111-1-1-224--2
Mullidae	22411122121322231141123312311311-31315332-55224432-	2412555554424334-113311-112411-44---
Loliginidae	13111--112111-111-111121221212-1121113321121142444	1212-2232-311313212432111231111-111--
Carangidae	--2212-211112211111-11111212-2134224555245343323455	53134351232223334335434322111142113--
Ariidae	1-21114312--21354421--2222433--54233231-2--111311-	---545345-22121211--214411--13-441311
Sepiidae	--1-1-1-1111-11--111-1-1-1111111-----	1--1211--321--111111--1-11-1-----
Sharks	525--1-1--3-1-----1-42543523-----21	--1-22-11343-5-1452---12--35-212311
Sphyraenidae	--1-----1-----1-1-15131-1-11-----	124-512-1--1-----1-111111--11-----
Psettodes	-----1-----1-----4-----1-----	1-1-1111-1-1-1-----1-----1-2-----
Leiognathidae	--11-1-1111-1--1-1111-1-1-1-1311-12-112-45243-	555555525-555413-54-23123155553221114
Trichiuridae	--21-1-----1-----1-12-----11-----	1122111-1222121--1-4-22-4142-145-42-1
Terapontidae	-----1-----1-----1-----1-----	4--1312-1-1-----355121-2--1-42-11-11
Polynemidae	-----2-----1-----1-----1-----	-----1-1-----1-1-----1121-3-----
Engraulidae	-----1-----1-----1-----1-----	--221-312-2215-15-14-55232-3153555554
Clupeidae	-----1-----1-----1-----1-----	24-51451352-4411351525535335524255524
Shrimp	-----2-----1-----1-----1-----	1-111-1-----1-11-1-11--11111142211
Stromateidae	-----1-----1-----1-----1-----	13-1-----211--1-1--1-11--52113315511
Drepanidae	--1-----11-----1-----1-----	54553-111-----1-----1111-11-1-----
Lactariidae	-----1-----1-----1-----1-----	1-1-1-1-1-1-----2-2-----
Sciaenidae	-----11-----1-1-----11-----	25-255221-21-112-114-11115512-242211
Chirocentridae	--1-----1-----1-----1-----1-----	111-1-1-1-31-3--2111-122-1111-1--13--
Parastromat	-----1-----1-----2-----1-----5-----	-----2-----1-11--21--11-122-11-11-1-----
Scombridae	--2-1-----1-1-1-1-1-1-1-1-1-1-1-----1-----	12121-121121221-2144332321111-----
Crabs	-----1-----1-1-1-1-----11-----	-----1-----1-2-1-----
Rays	--55--22--22534-1-1--11-1-----1-2121-----	4415--323--1-32--213--545--55555

	Group 1 (36 m)	Group 2 (50 m)	Group 3 (26 m)	Group 4 (29 m)
<i>Lutjanus sanguineus</i>	-55-2---1195-1111--	55432333521-312-1--3-42-312231-312-42143	-----255-23-1---1-55-41-21-2-----55---	1---21-335---
<i>Pentapodon longimanus</i>	-21421-1-1-5-11234-	24555-4441-5445535-2255435524555333	11--1-4-23114541154-4--11--113---	---1355545-1-
<i>Upeneus sulphureus</i>	1-1-----1-4-14-1-1	5455534552334511255354512154515555452-55	54-45153545555135-523555512-11-33451344-1	11-111255535-35
Sponges	1-----3432-43-4333-	-----1-2-----1-----1-----	-----3-----1-----1-----	-----1-----
Pentapodidae	-23133213132552323141	-3-322232211-1212-1-1-14121-1-1-----	-----11-4-1-1-----1-----	-----11-1-1
<i>Abalistes stellatus</i>	-4-4333425355341541-	153333244223211-2-----41211-322-1114-	-----3-23-224-14--22-2-----1-----	-----11-1-1
Other mullids	1--33211131-4-423254	-1-12--1-1-1-1-1-1-1-1-11-11-----	-33313-----1-11-1-3-11-----	-----11-1111
Other lutjanids	--15312--25155-4111--	-5-333312-1-12-11-2-1-1-1-121-2-13---	-----243-----111-2-15-2--3-25-55---	-----4-----
<i>Selaroides leptolepis</i>	11-254155-525553133-5	-----112233-541-1-11-11222-111-1-----52	-3452554333-21-1-1-231411-11-1-----1---	-----1121111-11
<i>Carcharhinus sealei</i>	-532-5---33-----	-----32-3-----2-----4-----3355-2225	4-----235-2221-2-----2-----2-21-21--	-----11-41-12-----
<i>Arius thalassinus</i>	--5-5131-52-2-21--4-	--112--1-1-11--1212332--1-12-3-43342	--1-424-24-5552-42-1-13-----3-----	-----2-----
Squid	13123---13244112441	1344211111232252113-----211541222	1111211232-1121-1-111212141415422-321551--	1-----1-----
Other foodfish	-55-----1331251-15114	-332211-211-12-3-1-----1211-11421	2111223-13--141111-11-414115-51-11222-55	1-----1-----
Trashfish	122-5--322345555151	14543213125234323133-----2343422251	1253-32-23-122133531343539455424-3411-2-2	1-----1-----
Other carangids	153232-31351541113331	-333212144122121-2322221413444-2134-31245	421221544514224-5-5224444553534551444541--	313213222542121
<i>Deeyatiidae</i>	--2---11-153221-1-1	---5---13---1-1-2---112-1-1-1-----	55-131--21-2-312-1122-55-5-55--5-555--	455-11-5531--1
Other leiognathids	111-121111112--11-13	--5551-55133534--141455543125513-5432111-	1114523434511314155115554533552314541555-1	122115142433-15
Other aridids	-5-14-3-----	-23--223-1-33--2--42-443--221-----	-----5-3-12-1-2-121-533--5555555445	1--1153213-3--
<i>Rastrelliger spp.</i>	-4-12--1-1-215--11-1	-----1-112-1-1-1-----11-1-214	-412-14113-5-1-3--111111-2-----21-4--	1-----1-----3
<i>Psettodes erumei</i>	-----1-1--111--	-----11-----1-1-11122	-----1-1-122-1-11-11--1-121--	-----1-----
Lobster	-----1-11-211111-1	-----1-2-1-1-1-1-1-----1-1-1-	1323211-----1-1-21111--2--1-1-1-----	-----1-----
Other rays	52--4--5-----	4-----1-----1-----1-----12-----	-----1-1-3-44-----21-1-55-452-43-1-	-----1-----
Cuttletfish	1-1-----11-11121	---111--11--11-11-1-----1-11-1-	3422423-3-11-11-11-111424-2233-1-1-----	-----1-----
Trichiuridae	111-----1-22-----1	3131--11-1-----2-----241-1351	1-525-1311451312411334-3-1-24-512212415-	-----1-----
<i>Chirocentridae</i>	-----1-----1-----1-----	-----1-----1-----1-----	2-----1-----21-----1-1-1-1-----	-----12111-311-----
<i>Clupeidae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----	-----1-----1-----1-----5-----	-----212112-1-----
<i>Scombridae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----21-----	-----1-----1-----1-----4-----	-----1-----
<i>Formio niger</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----211-----4	-----1121--31-1-----1-1--2-1-33244--	2-----11-11--
<i>Sardinella spp.</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----4	-----1121--31-1-----1-1--2-1-33244--	2-----11-11--
Other engraulids	-----1-----1-----1-----1-----	-----1-----1-----1-----3-----	-----5-1251-14-35-5151455-115214513-	1-----11-11--
<i>Stolephorus spp.</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----2111111-1	-----1-----1-----1-----2-2551551--	1-----1-----
<i>Drepanidae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----22-----	-2112-22242-121-1-1-1123312-----42234--	-----1-1-----
Other sphyraenids	-----1-----1-----1-----1-----	-----1-----1-----1-----1-11-----	-213-2314111-1-1-1-1-4114-124-1121323-	-----1-3-----
<i>Terapontidae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----4-----	-----111-1142-----431-----	-----1-----
<i>Pomadasy argyreus</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----3	311-415154-2112-1-11-11-11-432-3-54145--	1-1113--24-21-
<i>Dussumieria acuta</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----3	-----5122-3-25-421--43515431-2-32322552	1-133-1-41--
<i>Ilisha spp.</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----311-----	22-144--1-211223111-132543-----412544433	313111111-11--
<i>Sciaenidae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----11-----	1141252555515134551555451-422-5454551-1	222-55142453-15
<i>Leiognathus splendens</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	-----55-----2-5-5--2511-2151	-----4-11--
Other clupeids	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	1-----12-----1-----115111-----2-223-1	-----3-----
<i>Polynemidae</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	14331-----1-----1-----22111--13--	-----1-----
Crabs	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	34121-----1-1-1-----13-1111-1131-11452	-----1-----
Shrimps	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	-----1-----112-1-1-211-11-----11-3-----	-----1-----
<i>Lactarius lactarius</i>	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	-----1-----1-----1-----1-----	-----1-----

Fig. 2. Dendrogram of the Java Sea communities resulting from TWINSPAN analysis of R/V *Mutiara 4* data. Above: 1974-May 1976. Below: June-September 1976 (note that only part of the stations used for the analyses are shown).
[Gambar 2. Dendrogram komunitas ikan di Laut Jawa berdasarkan analisis TWINSPAN terhadap data Mutiara 4. Gambar atas: 1974-Mei 1976. Bawah: Juni-September 1976 (perlu diketahui bahwa hanya sebagian dari stasiun pengamatan yang dipakai untuk analisis).]




Fig. 3. Division of the Java Sea fish communities identified by TWINSPAN analyses. Above: November 1974-May 1976, with the shallow areas identified as a separate group (Group 2, top of Fig. 2). Below: TWINSPAN Groups 1-4 superimposed on stations for period June-September 1976.

[Gambar 3. Pembagian komunitas ikan di Laut Jawa berdasarkan analisis TWINSPAN. Gambar atas: November 1974 - Mei 1976, dengan daerah dangkal teridentifikasi sebagai suatu kelompok terpisah (Grup 2, bagian atas Gambar 2). Gambar bawah: TWINSPAN Grup 1-4 yang ditumpang-tindihkan pada lokasi stasiun pengamatan periode Juni-September 1976.]

Fig. 4. Plot of the stations on DCA Axes 1 and 2 for the survey period June-September 1976. Numbers indicate the TWINSPAN groups defined in the text.

[Gambar 4. Plot dari stasiun-stasiun pengamatan pada sumbu DCA 1 dan 2 untuk periode survei Juni - September 1976. Angka-angka menunjukkan grup TWINSPAN sebagaimana dibahas didalam tulisan.]




Table 3. Catch rates in kg hour⁻¹ of R/V Mutiara 4 in the waters near Tanjung Selatan, South Kalimantan, by season and year.

[Tabel 3. Hasil tangkapan per jam tarikan trawl oleh Mutiara 4 di perairan sekitar Tanjung Selatan, Kalimantan Selatan, menurut musim dan tahun.]

Fish group/Year	Southeast			Northwest				
	1976	1977	1979	1978	1980	1982	1983	1984
Ariidae	38	87	64	24	31	11	12	17
Carangidae	13	14	7	4	19	1	8	14
Clupeidae	56	30	35	22	41	8	10	13
Drepanidae	12	27	45	14	2	-	4	20
Leiognathidae	77	122	82	11	53	7	39	87
Lutjanidae	16	14	23	13	8	7	3	6
Pomadasyidae	15	25	32	13	20	6	6	11
Rays	25	66	41	19	31	43	-	37
Other fish	104	90	68	29	58	23	31	40
Total	356	475	397	149	263	106	113	245

Other species, such as *Lutjanus malabaricus* (non *sanguineus*) and *Pentaprion longimanus*, are relatively more abundant in the deepwater group (Group 2). This division is confirmed by the DCA analysis; the groups, although widely overlapping, do show some identity. Sponges, characterizing Group 1, were trawled in the depth range 20-40 m, in association with sandy bottoms and particularly in the depth range 30-40 m off South Kalimantan. The separation between Groups 3 and 4 is more difficult to interpret: Group 3 represents well the rich fauna of shallow inshore waters, including areas influenced by estuarine environments, but Group 4 appears to be less simply diverse. The stations of this group are geographically close to each other which might indicate that different ecological conditions are responsible for this separation. However, given that there are no indicator species and this group is mainly identified by a lack of taxa, it might well be that this difference results from the incomplete identification of the catches.

Table 3 presents a summary of mean catch rates (total and by major taxonomic groups) by year and seasons. The grand mean southeast monsoon catch rate in the period 1977-1979 was 409 kg-hour⁻¹ (SD: 60.5 kg-hour⁻¹); the grand mean northwest monsoon catch rate in 1978-1984 was only 175 kg-hour⁻¹ (SD: 75.2 kg-hour⁻¹). Table 3 also suggests shifting patterns of dominance, with, e.g., the Leiognathidae always dominating during the southeast monsoon, but only in one out of five seasons during the northwest monsoon.

Discussion

The data from the first survey period hardly allow a community study, mainly because of the rough taxonomic categories used to identify the catches. The results obtained reflect the uniformity in the distribution at family level in the study area and the lack of environmental barriers, with the exception of the inshore waters influenced by the estuarine environment. A better resolution was obtained by the analysis of the quasi-synoptic survey in June-September 1976, which revealed a major separation between the deeper waters of the Java Sea basin and the shallow inshore waters. The main faunal transition appeared to be at 25-30 m. Furthermore, it was possible to identify further distinctions, as for example Group 1, characterized by the presence of sponges and the scarcity of Leiognathidae as compared to the other groups where this family dominates.

It was not possible, due to scarcity of comparable stations, to study seasonal changes in community structure; they are presumed to exist, given the large observed changes in biomass.

The data collected through the *Mutiara 4* resources surveys, and now stored in an easily accessible database (Torres et al., this vol.) are of great value as they constitute the evidence of the status of the fisheries resources in the Java Sea 20 years ago. This type of information is not available in many tropical regions, where important environmental, mainly

fisheries-induced, changes have taken place in the last few decades.

It is, however, important to recognize that the quality of these data limits the range of further analyses that are possible (i.e., biodiversity and community dynamics). Particularly critical is the identification of the catches. Although this improved during the survey period, the use of large categories such as "other fish" or "trashfish" gives the opportunity to the data collector to easily "dump" into a large category any species that is difficult to identify. Furthermore, this process leads to inconsistencies between different parts of a survey, as it largely depends on the collector's personal capabilities.

Many trawl surveys have suffered, and still do because of similar shortcomings. The justification usually given is that the main aim is to monitor the important commercial groups. This is obviously valid but it might be argued that given the high cost of the surveys, and the relatively small increase in cost to carry out a full identification of the catches, identification to species level should be standard in any type of survey. Furthermore, the rapid change in the environment from the one side and the possibility to easily store and access the information derived from surveys from the other, would make surveys with properly identified catches of much greater value. Through this contribution, the authors recommend that the community of marine scientists ensures, as far as possible, a thorough identification in any type of marine resources surveys. This will give the surveys a more general value in addition to fulfilling their immediate objectives. They will become a piece of history.

Acknowledgements

The authors wish to acknowledge Daniel Pauly's effort to increase our understanding of tropical fish ecology by stimulating the maximum utilization of information from existing data sets. Without doubt, he has been the *primus motor* in the realization of this project.

References

- Dwiponggo, A. and M. Badrudin. 1980. Data of the Java Sea inshore monitoring survey by *Mutiara 4*. Mar. Fish. Res. Rep. Contrib. Demersal Fisheries Project 7A, 88 p.
- Ekman, S. 1953. Zoogeography of the sea. Sidgwick and Jackson, London. 417 p.
- Gunderson, D.R. 1993. Surveys of fisheries resources. John Wiley and Sons, Inc. 248 p.
- Losse, F.G. and A. Dwiponggo. 1977. Report on the Java Sea southeast monsoon trawl survey, June-December 1976. Mar. Fish. Res. Rep. Contrib. Demersal Fisheries Project 1, 119 p.
- † Martosubroto, P. and D. Pauly. 1976. *R/V Mutiara 4* survey data, November 1974 to July 1976. Mar. Fish. Res. Rep. Contrib. Demersal Fisheries Project 2, 136 p.
- Nelson, J.S. 1994. Fishes of the world. 3rd ed. John Wiley and Sons, New York. 600 p.
- † Saeger, J., P. Martosubroto and D. Pauly. 1976. Fish report of the Indonesian-German Demersal Fisheries Project. (Results of a trawl survey in the Sunda Shelf area). Mar. Fish. Res. Rep. Contrib. Demersal Fisheries Project 1: 1-18.

Structure and Dynamics of the Demersal Resources of the Java Sea, 1975-1979

PURWITO MARTOSUBROTO

Marine Resources Service
Fishery Resources Division
FAO of the United Nations
Viale delle Terme di Caracalla
00100 Rome, Italy

Demersal fisheries
catch/effort / Trawling
Abundance
Fishery surveys
1975-1979
1984

MA

1996. Structure and dynamics of the demersal resources of the Java Sea, 1975-1979 [Struktur dan dinamika perikanan demersal di Laut Jawa, 1975-1979], p. 62-76. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

The demersal fisheries of the Java Sea and their resource base were studied using commercial catch and effort data and the results of scientific surveys. Surplus production models were applied to the catch and effort data. Model results indicate that effort levels were near optimal in the late 1970s, except for the southeastern coast of Sumatra and northern coast of Java where, due to nearshore shrimp trawling, fishing effort was well above that required to generate maximum sustainable yield.

A principal component analysis demonstrated that 76% and 65% of the changes among years of demersal resource abundance was explained by the first two principal components for the lightly exploited areas and the highly exploited areas, respectively. Among areas representing different levels of fishing pressure, no difference was detected in the number of taxonomic groups at each trophic level significantly contributing to the first two principal components. Overall, throughout the 1970s, no major changes occurred in the trophic structure of the demersal resources in the various areas of the Java Sea.

Abstrak

Sumberdaya demersal dan perikananannya di Laut Jawa dianalisis melalui data statistik perikanan dan data dari kapal penelitian. Model surplus produksi diaplikasikan terhadap hasil tangkapan dan upayanya. Hasil analisis menunjukkan bahwa tingkat upaya mendekati optimal pada tahun 1970-an, kecuali untuk daerah pantai tenggara Sumatra dan pantai utara Jawa, dimana karena banyaknya trawl udang maka tingkat penangkapan sudah melebihi titik batas untuk mendapatkan hasil tangkapan yang maksimum.

Analisis "komponen pokok" terhadap hasil tangkapan menunjukkan bahwa masing-masing perubahan antar tahun (76% dan 65%) dari kelimpahan sumberdaya demersal dapat dijelaskan oleh dua komponen pokok awal untuk daerah yang tekanan penangkapannya kurang dan yang tingkat penangkapannya sudah jenuh. Daerah-daerah dengan keragaman tingkat penangkapan yang berbeda tidak menunjukkan adanya perbedaan dalam grup taksonomi pada setiap jenjang rantai makanan yang berarti mendukung pada dua komponen pokok awal. Secara keseluruhan, selama tahun 1970-an, tidak terlihat adanya perubahan dalam struktur rantai makanan untuk sumberdaya demersal di berbagai daerah Laut Jawa.

Introduction

The Java Sea is the most southern part of the Sunda Shelf, where the latter connects the western part of Indonesia with the Asian mainland. The Java Sea itself is bordered by the southern part of Sumatra, Kalimantan and by the northern coast of Java (Fig. 1).

The Java Sea resources make an important contribution to Indonesian fisheries, being the main supplier of fish protein for the population of Java, where most of the Indonesian

population lives. Moreover, the large number of fishers based on the shore of the Java Sea - more than 120,000 in 1979, plus 90,000 part-timers (DGF 1981) - reflects the importance of its fishing grounds for the livelihood of many people.

Development of the Indonesian fisheries started in the early 1970s when the first five-year plan was launched (Zachman 1973). At the start of the 1970s, estimates of potential yield were not available, except for extrapolations from the Gulf of Thailand. Thus, Tiews (1966) estimated annual yields of demersal fish to be 3.6 t·km⁻² for the Sunda Shelf, including the Java Sea, while Gulland (1971) presented a more




Fig. 1. Map of the Java Sea showing its bathymetry, and defining the (sub)areas used for data analysis and presentation of results, and roughly matching provincial borders.

[Gambar 1. Peta Laut Jawa yang menunjukkan pola batimetriknya dan pembagian daerah (sub-daerah) yang digunakan dalam analisis, yang mana batasannya kurang lebih mendekati batas provinsi.]

conservative estimate of 1.5-2.5 t·km⁻². However, the first systematic trawl surveys in the Java Sea indicated standing stocks at 2.15 and 3.24 t·km⁻² (Saeger et al. 1976; Pauly et al., this vol.). Assuming annual yield to be about half the standing stock size leads to potential yields even lower than suggested by J. Gulland, which points both at the usefulness of surveys and the danger of extrapolating.

The rapid expansion of the Java Sea fisheries during the 1970s was driven by the deployment of bottom trawlers and purse seiners (Martosubroto 1978). However, other fishing gears also increased substantially. During this period, total catch from the Java Sea more than doubled, from 140,000 t in 1970 to 332,000 t in 1979. The demersal catch, however, increased at a slightly slower rate, from about 60,000 t to 138,000 t in 1979, or an increase of 130%.

The operation of the demersal trawlers was largely limited to waters shallower than 40 m, and most fishing effort was applied along the northern coast of Java, where the small-scale fishers were (and still are) concentrated. Latent conflicts between the small-scale fisheries and trawl operators were inevitable.

One way to address these was through studies of the fisheries and their resource base, aimed at identifying a level of demersal trawl effort compatible with the overall productivity of the resource, and the need to sustain existing small-scale fisheries (Sujastani 1978).

Some management measures for the fisheries in the Java Sea were introduced in the early 1970s, before these studies were conducted. These measures included boat size regulations (Zachman 1973), zoning of fishing activities, and mesh size limits (SCSP 1979). They did not achieve their aims, however. This led, in conjunction with the breaking out of open

conflicts between small-scale fishers and trawl operators, to the banning of trawl fishing in the waters surrounding Java and Sumatra in 1980 and 1981, respectively (Sardjono 1980).

Trawl survey data in the Java Sea became available from 1974 onwards and were collected continuously to the end of the 1970s (Martosubroto and Pauly 1976; Saeger et al. 1976; Lose and Dwiponggo 1977; Dwiponggo and Badrudin 1980; Bianchi et al., this vol., Pauly et al., this vol., and see the database described in Torres et al., this vol.). The present study evaluates the status of demersal resources of the Java Sea in the 1970s, based on these survey data, and on commercial catch and effort data for the same period, collected by staff of the Indonesian Directorate General of Fisheries (DGF).

Materials and Methods

Study Area

The Java Sea has an almost rectangular shape with a long axis of approximately 890 km parallel to Java and a short axis of 390 km. Its bottom gently slopes from the shoreline to the center and from west to east; the western part has an average depth of about 20 m while the eastern part is about 60 m deep (Fig. 1).

The bottom sediment of the Java Sea consists mostly of mud. A map compiled by Emery (1969) revealed that 69% of the total area consists of thick gray mud, 17% of mud and sand, and 12% of sand, with the remaining 2% consisting of rocks and corals mainly near Bangka and Billiton Islands, in the vicinity of the Sunda Strait and, in the east, along the edge of the continental shelf (Fig. 2).




Fig. 2. Bottom types of the Java Sea (after Emery 1969).
 [Gambar 2. Tipe dasar perairan Laut Jawa [(menurut Emery 1969)].

The hydrography of the Java Sea has been studied since the beginning of the century by Dutch scientists (Berlage 1927; van Riehl 1932) and from the 1950s by Indonesian scientists (Soeriatmadja 1956; Sjarif 1959). A comprehensive description of the Java Sea was presented by Wyrki (1961) as part of his extensive study of Southeast Asian waters.

These studies demonstrated a strong impact of the west monsoon which extends from December to February, and of the east monsoon which lasts from June to August. The rest of the year forms transition periods between the two regimes. During the west monsoon, the wind generates eastward-moving surface currents that bring low salinity water of the South China Sea into the Java Sea. The heavy rains during the west monsoon increase the runoff from rivers in Sumatra, Kalimantan and Java and further depresses the salinity in the Java Sea. It is not uncommon, in its eastern part, for the 3.0‰ isohaline to be pushed far into the open ocean. During the east monsoon, these conditions are reversed; westward winds generate surface currents that bring high salinity water from the Makassar Strait and the Flores Sea into the Java Sea and pushes its low salinity water into the southern South China Sea. The high salinity water masses reach their maximal westward penetration in September (see Fig. 2 in Venema, this vol.).

The distribution of primary productivity also varies with the monsoons (Doty et al. 1963). During the west monsoon, the

average primary productivity of the surface layer increases not only toward the coast but also from west to east. The 1 mgC·hour⁻¹·m⁻³ isoline runs approximately in the mid-part of the Java Sea, dividing the area into two parts (Fig. 3A). During




Fig. 3. Surface primary productivity in the Java Sea, in mgC·hour⁻¹·m⁻³ (after Birowo et al. 1975): A, during the west monsoon; B, during the east monsoon.
 [Gambar 3. Produktivitas primer Laut Jawa dalam mgC·jam⁻¹·m⁻³ (menurut Birowo et al. 1975): A, selama musim barat; B, selama musim timur.]

the east monsoon, this isoline is pushed westward and it eventually lies parallel to the northern coast of Java (Fig. 3B).

Statistical (Catch and Effort) Data

Commercial fishing operations exploiting demersal stocks are normally limited to depth ranges shallower than 40 m, whereas traditional small-scale gears operate inside the 20-m depth contour. For these reasons, the surplus production and principal component analyses presented below were applied to areas within the depth of 40 m. To analyze the trawl survey data, the Java Sea was arbitrarily divided into eight areas: five off the southeast coast of Sumatra and north coast of Java (A, B, C, D and H), and three off the south coast of Kalimantan (E, F, G; see Fig. 1). The data from areas G and H were not considered because they were surveyed on two occasions only. The limits of these areas roughly correspond to the provincial borders (Fig. 1), defining how catch and effort statistics were aggregated.

The publication of Indonesian fisheries statistics by DGF began only in 1972, and those published prior to 1975 were of limited use for stock assessment purposes, mainly because the information was overaggregated: the landing statistics were broken down by province without further partition within each province. Thus, those interested in assessing the stocks of, say Central Java Province, with fishing grounds in the Java Sea (a part of the Pacific Ocean), and in the Indian Ocean, were forced to visit the Fisheries Services at the *Kabupaten* (i.e., county) level to obtain data pertinent to the one, or the other ocean (see e.g., Martosubroto 1978 and Sujastani 1978). Moreover, although the statistics from the mid- to late 1970s presented more detailed information, the species composition of the catch of various gears still could not be derived from the data; hence the adjustments presented below.

Fish caught in Java Sea are landed in three provinces (West Java, Central Java and East Java) and one municipality (Jakarta) along the north coast of Java, two provinces in southeast Sumatra (Lampung and South Sumatra) and three provinces along the southern coast of Kalimantan (West Kalimantan, Central Kalimantan and South Kalimantan) (Fig. 1). As appropriate statistics were not available at the *Kabupaten* level and provincial statistics often lump catches from different areas of the Java Sea, a number of assumptions were made to disaggregate the available data:

1. the catches and number of fishers along the coast of East Java Province facing the Java Sea represent 45% of these values for the whole of Java (the Madura Strait is excluded in the analysis as it has been closed to trawling since 1975);
2. the fisheries statistics for that part of the coast of South

Sumatra Province facing the Java Sea account for 35% of the total landings of that province;

3. the fisheries statistics for that part of the coast of West Kalimantan Province facing the Java Sea represent 5% of the statistics of the whole province; and
4. the fisheries statistics for that part of the coast of South Kalimantan Province facing the Java Sea account for 75% of the statistics of the whole province.

These assumptions are based on relative coastline length within each provincial unit. Obviously, any drastic changes in the deployment of effort within these provinces could invalidate the assumptions for a particular area. Over the 1975 to 1979 period, however, such drastic changes did not occur. Since 1980, there have been major changes, most notably the above mentioned ban on trawling.

Trawl Survey Data

Pauly et al. (this vol.) and Bianchi et al. (this vol.) described the trawl survey conducted from 1974 to 1979 in the Java Sea, which generated the data analyzed here.

The *Mutiara 4* and "Thailand trawls" were used for all of these surveys (see Pauly et al., this vol. for descriptions). The rigging of the net was slightly changed in June 1976 (Table 1), prior to gathering the data underlying Fig. 4; however, as the modifications were minor, catch rates can be assumed not to have been altered.

Fig. 5 illustrates the variability of the trawl survey data used here. As might be seen, there is a linear correlation between the means and the standard deviation of the catch rates for the demersal groups, which suggested that a log-transformation of the data might be appropriate for at least some of the analyses (Steel and Torrie 1960). Retransformed means (Fig. 6) were thus obtained following Bliss (1967) using

$$y = \exp(x + s^2/2) - 1 \quad \dots 1)$$

where

y = the retransformed catch per haul;

Table 1. Specification of trawl net used during the *Mutiara 4* survey.
[Tabel 1. Spesifikasi jaring trawl Mutiara 4 yang digunakan dalam survei.]

Trawl characteristics	Prior to June 1976	Since June 1976
Circumference (lower bosom)	147 meshes, 160 mm stretched	147 meshes, 160 mm stretched
Cod-end mesh	40 mm	40 mm
Headline length	15.0 + 3.0 + 15.8 = 33.8 m	16.85 + 3.0 + 16.85 = 36.7 m
Footrope length	20.1 + 2.0 + 20.1 = 42.2 m	21.25 + 2.0 + 21.25 = 44.5 m
Floats on headline	27-30	21
Iron rings on footrope	24	20
Chains and weights	1 chain (8 kg) on bosom, 15 kg chain on wing ends, 8 kg weight on legs	1 chain (8 kg) on bosom, 15 kg chain on wing ends
Otterboard dimensions	2.0 x 1.0 x 0.05 m, one float 50 x 20 cm on each board (92 kg each)	2.0 x 1.0 x 0.05 (92 kg each)
Bridles and legs	50 m bridles + 25 m legs	50 m bridles + 18 m legs




Fig. 4. Trawl survey of *R/V Mutiara 4* in the Java Sea, from 17 June to 22 September 1976, illustrative of other surveys considered on this contribution. Each dot represents a valid 1-hour haul (see Pauly et al., this vol., for details on the survey methodology and assumptions).
 [Gambar 4. Survei trawl kapal penelitian Mutiara 4 di Laut Jawa, dari 17 Juni hingga 22 September 1976, digambarkan terhadap survey yang lain. Tiap bulatan menggambarkan 1 jam tarikan (lihat Pauly et al., dalam buku ini, dalam hal metodologi survei dan asumsinya.)




Fig. 5. Relation between the magnitude and the variability of demersal trawl survey catch rates. Left: survey cruises of 1979 in areas A and B (southeastern Sumatra and north Java coast). Right: mean annual C/f and their standard deviation in areas C and D (north coast of Central and East Java) for the years 1975 - 1979.

[Gambar 5. Hubungan antara hasil tangkapan per upaya dan variasinya dari trawl survei ikan-ikan demersal. Kiri: pelayaran survei tahun 1979 di daerah A dan B (bagian tenggara Sumatra dan pantai utara Jawa). Kanan: rata-rata hasil tangkapan per upaya (C/f) dan simpangan bakunya di daerah C dan D (pantai utara Jawa Tengah dan Timur) dalam tahun 1975-1979.]

x = the original mean catch per haul; and
 s^2 = the variance of the log-transformed data.

Further details on the catch/effort data used here may be found in the survey reports, i.e., Martosubroto and Pauly (1976) and Saeger et al. (1976) for the first phase (November 1974 to July 1976; see also Pauly et al., this vol.) and Losse and Dwiponggo (1977) and Dwiponggo and Badrudin (1980) for the second phase (August 1975 to 1979; see also Bianchi et al., this vol.)

Principal Component Analysis

Principal component analysis (PCA) as used here involves the analysis of a multidimensional cloud of sample points, as defined by the abundances of the species they contain. Each species represents an axis (dimension), and the position of each sample point is defined by as many species

(axes, dimensions) as the set of samples included. A set of samples with 100 species, for example, would be defined as a cloud of points in a space of 100 dimensions. The relationships among the points can be studied by combining sets of axes (species) into a reduced set of principal axes, each of which is made up of a simple linear combination of species. The axes are extracted in decreasing order of their eigenvalues, which represent successively smaller amounts of the variation of the cloud in the hyperspace. Often, 80% or more of the total variation of the cloud can be represented in the first few principal axes. This means that the original data cloud can be represented with very little distortion by replotting the sample points based on their coordinates on a few principal axes. The version used in this study employed an eigenanalysis of the correlation matrix, thus yielding axes based on a data cloud which has been centered and standardized with respect to the principal axes. These principal axes summarize the variations




Fig. 6. Mean trawl survey catch rates in areas A-D (above) and E-F (below), illustrating small differences between original data (i.e., arithmetic means; solid lines) and log-transformed data (i.e., geometric means; dotted lines), thus justifying the use of the log-transformation to stabilize variances.

[Gambar 6. Rata-rata hasil tangkapan per upaya survei trawl di daerah A - D (atas) dan E - F (bawah), menggambarkan perbedaan kecil antara data asli [rata-rata, garis penuh] dan data yang ditransformasikan ke dalam logaritma (rata-rata geometrik, garis putus-putus), yang menunjukkan perlunya transformasi untuk menstabilkan keragaman.]

of sets of species, which can be contrasted against variations in environmental factors or time to yield insights into casual relationships (Sokal and Rohlf 1969; Sneath and Sokal 1973; Pielou 1977).

Surplus-Production Modelling

The application of the surplus production model (Schaefer 1954, 1957; Ricker 1975) to data from a fishery implies that the dynamics of the population exploited by that fishery follows a logistic growth trajectory.

Although this assumption has not been explicitly demonstrated to apply to multispecies tropical stocks, experience has shown that the fit of surplus-production model to multispecies assemblages is generally better than that of the individual species, as indicated by the higher correlation in the regression of catch per unit of effort with effort (FAO 1978; Pauly 1979). Two likely explanations for this have been offered by Pope (1979), i.e.:

1. treating the whole species as a single unit implicitly accounts for the biological interactions between the component species; and/or
2. the fishers react to local decline of their catch/effort by shifting their effort to other components of the multispecies resources, and thus cause an evening out of the catch/effort vs. effort relationship.

However good the fit of the surplus-production model to the multispecies situation may be, the main criticism remains, however, that the model does not explicitly incorporate interaction between the different component species of stocks (Pauly 1979).

In view of these drawbacks, Pope (1979) extended Schaefer's surplus-production model to explicitly account for biological and technological interactions. In Pope's reformulation, biological interactions are incorporated through either predator-prey or competitive relationships, while technological interactions are included through fixed ratios of the fishing mortalities, affecting the different component species of the multispecies stock.

The most important insights resulting from Pope's work are:

1. the apparent maximum sustainable yield (MSY) that can be extracted from a multispecies assemblage using a gear with a fixed set of catchabilities (e.g., a trawl) will almost invariably be lower than the theoretical MSY that may be obtained from the assemblage as a whole;
2. if the species in the assemblage are linked mainly through competitive relationships, then the stronger the competition, the smaller will the overall MSY of the assemblage be; and
3. if the species in the assemblage are linked mainly through predator-prey relationships, then the overall MSY that can be extracted from the assemblage will be higher than if the species are mainly competing with each other.

Conclusion (3) certainly makes sense: higher exploitation of predators should lead to an increase in prey population (May et al. 1979).

While a useful heuristic tool, Pope's multispecies extension of the surplus-production model cannot be used for practical assessments, due to an excessive number of parameters (x), which is a quadratic function of the number of interacting species (n), i.e.,

$$x = (cn + 1)^2 - 1 \quad \dots 1)$$

However, when investigating the demersal stocks of the Gulf of Thailand, Pope (1979) noticed that while the majority of species declined under exploitation, the ratio of their relative abundances remained roughly constant. This led to his suggestion that simple surplus-production models treating an entire multispecies assemblage as if it were a single species may be appropriate, and would lead to an MSY estimate not

very different from those that would have been obtained, had an explicitly multispecies model been used.

This provides the justification for the application, in this study, of a simple Schaefer-type surplus-production model to the multispecies assemblages of the Java Sea, performed by plotting catch per effort (C/f) against effort (f) viz.

$$C/f = a + bf \quad \dots 2)$$

then multiplying both sides by f to obtain a (parabolic) surplus-production curve viz.

$$C = af + bf^2 \quad \dots 3)$$

whose maximum indicates MSY and optimum effort (f_{opt}).

To evaluate the sensitivity of the results to the quality of the input data, three different measures of C/f were used (Index 1, 2 and 3, defined in the legends of Figs. 13 and 14).

On the other hand, in view of the limited number of available data points (=years with catch and effort data), I have abstained from applying Gulland's approach for simulating equilibrium (i.e., plotting catch/effort of one year against the mean effort for that year and one or several preceding years; Gulland 1961). This should have resulted in an overestimation of MSY and optimum effort (Pauly 1984, 1987).

Results and Discussion

Fisheries of the Java Sea

NORTH COAST OF JAVA

Java is the most developed area in Indonesia and, at the same time, one of the most densely populated areas in the world. It is therefore not surprising that the island shows also the highest concentration of fulltime fishers. Also, the infrastructure (landing and auction sites, wet markets, roads) of the fishery industry on the north coast of Java is more developed than in other areas of Indonesia. As a result, the northern coast of Java accounted, in 1979, for 71.4% of the total catch from the Java Sea. However, most of this consisted of pelagic fish since 41.3% was caught by seines, 21.4% by gillnets, 15.7% by liftnets and other gears, and only 21.6% by trawlers.

Information on the number of fishers for each type of fishing gear is not available. However, based on the total number of trawlers and purse seiners operating based along the north coast of Java, and on the average number of crew on trawlers (8) and purse seiners (28) (Baum 1978; Budihardjo 1978), the number of fishers engaged in these fisheries could be estimated as approximately 24,600 in 1979, or about 25% of the total number of fishers in the area. The majority of the fishers (75%) were thus engaged in small scale or artisanal fisheries and their

productivity ($1.25 \text{ t-fisher}^{-1} \cdot \text{year}^{-1}$) was much lower than that of the crew of purse seiners and trawlers ($4.55 \text{ t-fisher}^{-1} \cdot \text{year}^{-1}$). Consequently, the number of crew on trawlers and purse seiners increased from 14,000 in 1976 to 25,000 in 1979, while the number of traditional fishers dropped in the same period from 84,000 to 75,000, many of them transferring from the artisanal to the industrial subsector.

SOUTHEAST COAST OF SUMATRA

The fisheries on the southeast coast of Sumatra were dominated in the 1970s by liftnets and gillnets which in 1979 contributed 53% and 19%, respectively, of fisheries catches, while trawlers contributed only 3%.

The total number of fulltime fishers in 1979 was 9,000; of these, about 3% worked as crew on trawlers. The rate of increase of the number of fishers ($11\% \text{ year}^{-1}$) in this area was higher than along the northern Java Sea coast and may have been related to population transmigration from Java.

Table 2. Annual demersal catches (t) by gear and year from the North Java and Southeast Sumatra coasts.

[Tabel 2. Hasil tangkapan tahunan ikan-ikan demersal (t) berdasarkan jenis alat dan tahun untuk pantai utara Jawa dan Sumatra bagian tenggara.]

Type of gear	Year				
	1975	1976	1977	1978	1979
Trawl	30,789	40,323	47,799	49,118	53,960
<i>Dogol</i> ^a	3,937	10,099	8,521	10,045	9,357
Beach seine	12,624	4,640	2,702	3,547	5,280
Bottom gillnet	1,631	13,155	12,023	12,538	14,566
Liftnet	20,070	26,190	34,325	43,370	40,102
Longline	5,051	1,794	1,565	1,339	1,470
Handline	4,041	10,143	10,143	10,839	11,741
Stow net	585	667	125	200	178
Trap	1,138	8,927	7,232	7,550	4,764
Total	79,866	115,938	124,435	138,546	141,418
Total demersal catches	69,106	90,252	95,279	103,088	111,703

^a Modified Danish seine.

Table 3. Annual demersal catches by gear and year from the coast of Southern Kalimantan.

[Tabel 3. Hasil tangkapan tahunan ikan-ikan demersal berdasarkan jenis alat dan tahun untuk pantai selatan Kalimantan.]

Type of gear	Year				
	1975	1976	1977	1978	1979
Trawl	576	1,016	641	899	1,100
Beach seine	2,411	2,748	2,604	1,803	2,451
Bottom gillnet	6	6	72	1,479	2,708
Liftnet	408	2,722	1,917	4,999	3,897
Bottom longline	320	240	679	1,618	1,836
Handline	30	2,032	350	747	513
Stow net	362	1,073	450	631	705
Traps	354	2,726	338	72	47
Total	4,467	12,563	7,051	12,248	13,257
Total demersal catches	15,401	19,572	15,780	22,718	26,321

Table 4. Structure of taxonomic principal components for the demersal resources in areas A-D (Southeast Sumatra and North Java Coast), ranked by their eigenvectors; groups with similar eigenvectors tend to co-occur.

[Tabel 4. Struktur komponen pokok berdasarkan pembagian taksonomi dari sumberdaya demersal di daerah A hingga D (Sumatra bagian tenggara dan pantai utara Jawa), diklasifikasi berdasarkan nilai eigenvector; grup dengan nilai eigenvector yang berdekatan cenderung untuk muncul bersama.]

Taxonomic group	PC-1 (40.1% of var.)		PC-2 (25.2% of var.)	
	Eigen-vector	Correlation ^a	Eigen-vector	Correlation ^a
Gerreidae	0.25	0.99**	-0.02	-0.06
Mullidae	0.24	0.95**	0.04	0.13
Carangidae	0.23	0.89**	-0.10	-0.32
Lutjanidae	0.22	0.85*	-0.13	-0.41
Sharks and rays	0.21	0.84*	0.11	0.35
<i>Thenus orientalis</i>	0.19	0.74	0.18	0.57
Nemipteridae	0.19	0.75	0.22	0.67
Serranidae	0.05	0.21	-0.04	-0.13
Squids	0.05	0.21	0.01	0.03
<i>Priacanthus</i> spp.	0.04	0.16	0.29	0.90**
Balistidae	0.03	0.10	0.25	0.79
Sphyraenidae	0.01	0.04	-0.26	-0.81*
Muraenesocidae	-0.03	-0.12	0.23	0.71
Pentapodidae	-0.05	-0.19	0.27	0.84*
Trichiuridae	-0.05	-0.20	-0.07	-0.22
Chirocentridae	-0.06	-0.22	0.05	0.15
Other food fish	-0.09	-0.35	-0.04	-0.13
Trash fish	-0.09	-0.36	-0.29	0.90**
Rachycentridae	-0.10	-0.39	-0.14	-0.44
Terapontidae	-0.11	-0.45	0.00	0.01
Synodontidae	-0.11	-0.43	0.21	0.64
Ariidae	-0.12	-0.45	0.25	0.76
<i>Formio niger</i>	-0.14	-0.53	-0.22	-0.67
Leiognathidae	-0.14	-0.56	0.02	0.05
Polynemidae	-0.15	-0.59	-0.04	-0.13
Engraulidae	-0.16	-0.64	0.16	0.51
Heterosomata	-0.16	-0.63	0.24	0.73
Pomadasyidae	-0.17	-0.66	0.23	0.71
Shrimps	-0.17	-0.65	0.19	0.60
Other invertebrates	-0.18	-0.71	-0.14	-0.44
Sciaenidae	-0.19	-0.75	0.12	0.36
Scombridae	-0.19	-0.73	-0.05	-0.16
Drepanidae	-0.20	-0.78	-0.09	-0.29
Stromateidae	-0.21	-0.83*	-0.15	-0.46
Clupeidae	-0.22	-0.89**	-0.09	-0.29
Crabs	-0.22	-0.88**	-0.13	-0.40
Cuttles	-0.23	-0.90**	0.00	0.00
<i>Lactarius lactarius</i>	-0.25	-0.98**	-0.01	-0.02

^a* significant at p <0.10; ** significant at p <0.05.




Fig. 7. Trend in abundance of some taxonomic groups of Areas A-D (Southeast Sumatra and North Java coast) showing high correlations with PC-1 and PC-2 (see also Table 4); A: positive correlations with PC-1; B: negative correlations with PC-1; C: positive correlations with PC-2; D: negative correlations with PC-2
[Gambar 7. Perubahan kelimpahan beberapa grup ikan [menurut taksonomi] di daerah A-D (tenggara Sumatra dan pantai utara Jawa) yang menunjukkan korelasi tinggi dengan PC-1 dan PC-2 [lihat Tabel 4]; A: korelasi positif dengan PC-1; B: korelasi negatif dengan PC-1; C: korelasi positif dengan PC-2; D: korelasi negatif dengan PC-2.]

Table 2 summarizes the available data on demersal catches for the north coast of Java and the southeast coast of Sumatra.

SOUTH COAST OF KALIMANTAN

In the 1970s, Kalimantan was far less developed than the two above discussed areas. Landing facilities were lacking and roads hardly existed. In 1979, the bulk of the fisheries catches (62%) originated from gillnets, followed by liftnets (13%).

In 1970, a joint venture company was established in Kotabaru to exploit the shrimp resources off the southern and eastern coasts of Kalimantan. This company started with 6 trawlers in 1970 and had 60 trawlers by 1975 (Naamin and Uktolseya 1976) and employed relatively high technology compared to the existing trawlers in this area. Because this company was only interested in shrimp, all other fish were thrown back to the sea, a practice otherwise very uncommon in Western Indonesia. Data on the quantity of fish thrown overboard are not available, although rough estimates could be made using the average ratio of 1:22 in weight of shrimp to fish. Since total shrimp landings of this company on the south coast of Kalimantan were 1,124 t in 1975 (Naamin and Uktolseya 1976), the by-catch dumped in that year could have reached 24,000 t, an amount similar to total fish landings in the same area. Whether the increase of fishing pressure on shrimp caused changes in the ratio of shrimp to by-catch is not known, as the studies reported on by Naamin and Uktolseya (1976) were discontinued.

The number of fulltime fishers along the coast of southern Kalimantan was estimated as about 11,000 in 1979.

Table 3 summarizes the available data on demersal catches from the coast of South Kalimantan.

Table 5. Product moment correlation^a of the first principal component for the demersal resources in individual areas of the Java Sea.

[Tabel 5. Korelasi product moment^a dari komponen pokok awal dari sumberdaya demersal di setiap daerah (A hingga D) di Laut Jawa.]

Taxonomic group	A-D	A	B	C	D
Gerreidae	0.99**	-0.69	0.67	0.78	0.71
Mullidae	0.95**	-0.49	0.73	0.46	0.68
Carangidae	0.89**	-0.75	0.09	0.69	0.92**
Lutjanidae	0.85**	-0.64	0.81**	0.42	0.86**
Sharks and rays	0.84**	0.32	0.56	0.85**	0.78
Nemipteridae	0.75	0.35	0.79	0.78	-0.96**
<i>Thenus orientalis</i>	0.74	-0.13	0.77	0.17	-0.99**
Other invertebrates	0.71	0.58	-0.95**	0.54	0.44
Serranidae	0.21	-0.83	0.35	0.17	0.06
Squids	0.20	-0.75	-0.74	0.83**	0.78
<i>Priacanthus</i> spp.	0.16	-0.90**	-0.59	0.71	0.49
Balistidae	0.10	0.89**	0.55	0.51	-0.99**
Sphyracidae	0.04	-0.66	-0.50	0.48	0.52
Muraenesocidae	-0.12	0.49	0.67	0.64	0.40
Pentapodidae	-0.19	-0.20	-0.87**	0.13	-0.94**
Trichiuridae	-0.20	0.85	0.04	-0.97**	0.41
Chirocentridae	-0.22	0.74	0.13	0.58	0.39
Trash fish	-0.36	0.19	-0.66	0.89**	0.42
Other food fish	-0.36	-0.36	0.71	0.36	0.99**
Rachycentridae	-0.39	0.64	-0.45	0.89**	0.33
Synodontidae	-0.43	0.70	0.05	0.79	0.81**
Terapontidae	-0.45	0.94**	0.28	-0.56	-0.98**
Ariidae	-0.45	0.23	0.17	-0.93**	0.66
<i>Formio niger</i>	-0.53	0.69	-0.97**	-0.54	0.33
Leiognathidae	-0.56	0.65	-0.52	-0.81**	0.90**
Polynemidae	-0.59	0.82	-0.17	0.15	0.99**
Heterosomata	-0.63	0.47	0.36	0.89**	-0.92**
Engraulidae	-0.64	0.80	0.56	-0.68	0.88**
Shrimps	-0.65	0.93**	-0.46	-0.55	-0.99**
Pomadasyidae	-0.66	0.99**	0.17	0.43	0.83**
Scombridae	-0.73	0.43	-0.68	0.33	0.61
Sciaenidae	-0.75	0.74	0.51	0.48	0.71
Drepanidae	-0.78	-0.69	0.62	0.34	0.53
Stromateidae	-0.83**	0.99**	-0.48	-0.19	0.70
Clupeidae	-0.89	0.90**	-0.64	0.23	0.98**
Crabs	-0.89**	0.63**	-0.71	-0.96**	0.72
Cuttles	-0.90**	0.93**	-0.44	0.75	0.85**
<i>Lactarius lactarius</i>	-0.98**	-0.69	0.22	0.90**	0.79
Years included	1976-79	1975-79	1975-79	1976-79	1975-79
Variance explained (%)	47.4	33.0	40.9	57.3	40.1

^a ** significant at $p < 0.05$.

Principal Component and Surplus-production Analyses

NORTH COAST OF JAVA/SOUTHEASTERN SUMATRA

Based on the correlation matrix of taxonomic group abundances during the five-year period from 1975 to 1979, the PCA indicated that 65.5% of the total variances of demersal resources abundance can be explained by the first two principal components (Table 4). The high positive values for the coefficients of the sharks and rays, Carangidae, Lutjanidae and Mullidae in the first principal component demonstrate their important contribution to the total variance of demersal abundance (Fig. 7A). The high negative values for Clupeidae, *Lactarius lactarius*, Stromateidae, cuttlefish and crabs reflect their increasing trends during the study period (Fig. 7B). However, they cannot compensate for the decline of the major groups, the overall result being a weak declining trend for the demersal assemblage as a whole.

The second principal component indicates the relative importance of Pentapodidae, Priacanthidae and "trash fish" (Fig. 7C). Again, the declining trend of these groups explains the variation of the total group in the second principal component, there being no strong compensation by other groups (Fig. 7D).

An analysis of the principal components in the individual areas (Table 5) shows that the various taxonomic groups contributed differently to the first principal component. However, it is not clear why there are more species groups explaining the total variation in Areas A, C and D than in area B. Still, the Gerreidae, Nemipteridae and Synodontidae show a declining trend in most areas. Biological data are not available which could be used to explain this phenomenon, but it may be that their juveniles are more susceptible to artisanal fishing gears that employ very small mesh sizes, such as liftnets and pushnets.




Fig. 8. Trends in abundance of some taxonomic groups of areas C and D (north coast of Central and East Java), showing high correlations with PC-1 and PC-2 (see also Table 6); A: positive correlations with PC-1; B: negative correlations with PC-1; C: positive correlations with PC-2; D: negative correlations with PC-2.

[Gambar 8. Perubahan kelimpahan beberapa grup ikan [menurut taksonomi] di daerah C dan D [pantai utara Jawa Tengah dan Timur] yang menunjukkan korelasi tinggi dengan PC-1 dan PC-2 [lihat Tabel 6]; A: korelasi positif dengan PC-1; B: korelasi negatif dengan PC-1; C: korelasi positif dengan PC-2; D: korelasi negatif dengan PC-2.]

Table 6. Structure of taxonomic principal components for the demersal resources in areas C-D (north coast of Central and East Java), ranked by their eigenvectors; groups with similar eigenvectors tend to co-occur.

[Tabel 6. Struktur komponen pokok berdasarkan pembagian taksonomi dari sumberdaya demersal di daerah C-D (pantai utara Jawa Tengah dan Timur) diklasifikasi berdasarkan nilai eigenvector; grup dengan nilai eigenvector yang berdekatan cenderung untuk muncul bersama.]]

Taxonomic group	PC-1 (43.0% of var.)		PC-2 (31.5% of var.)	
	Eigen-vector	Correlation ^a	Eigen-vector	Correlation ^a
Muraenesocidae				
<i>Lactarius lactarius</i>	0.23	0.94***	-0.06	-0.20
Heterosomata	0.22	0.87**	0.01	0.04
Cuttles	0.22	0.88***	-0.08	-0.27
Carangidae	0.22	0.89***	-0.05	-0.19
Chirocentridae	0.21	0.84**	-0.14	0.49
Sharks and rays	0.21	0.86**	0.07	0.24
Squids	0.18	0.74	0.08	0.27
Synodontidae	0.18	0.71	0.15	0.52
Gerreidae	0.18	0.74	0.17	0.60
Drepanidae	0.16	0.65	-0.12	-0.42
Other food fish	0.16	0.64	-0.14	-0.49
Pomadasyidae	0.15	0.60	-0.01	-0.04
Nemipteridae	0.15	0.60	0.21	0.74
Sphyraenidae	0.14	0.56	-0.21	-0.73
Other invertebrates	0.14	0.57	-0.18	-0.63
Rachycentridae	0.14	0.59	-0.19	-0.64
Lutjanidae	0.14	0.59	-0.06	-0.22
Trash fish	0.14	0.58	0.22	0.76
Scombridae	0.12	0.47	-0.09	-0.30
Sciaenidae	0.11	0.45	0.17	0.60
Clupeidae	0.11	0.45	-0.21	-0.73
Muraenesocidae	0.10	0.42	0.24	0.83**
<i>Priacanthus</i> spp.	0.10	0.41	0.24	0.84**
Balistidae	0.10	0.39	0.20	0.68
Pentapodidae	0.09	0.37	0.26	0.90***
Polynemidae	0.09	0.38	-0.17	-0.60
Mullidae	0.08	0.34	0.27	0.93***
Serranidae	0.05	0.20	-0.04	-0.13
<i>Thenus orientalis</i>	-0.00	-0.02	0.27	0.93***
<i>Formio niger</i>	-0.02	-0.09	-0.24	-0.83**
Stromateidae	-0.05	-0.22	-0.25	-0.86**
Ariidae	-0.15	-0.60	0.15	0.53
Terapontidae	-0.18	-0.75	-0.13	-0.43
Shrimps	-0.21	-0.83**	0.09	0.32
Leiognathidae	-0.23	-0.91***	0.07	0.26
Trichiuridae	-0.24	-0.99***	-0.01	-0.03
Engraulidae	-0.24	-0.95***	0.07	0.25
Crabs	-0.24	-0.97***	-0.01	-0.05

*** significant at $p < 0.05$; ** significant at $p < 0.01$.

Table 7. Structure of taxonomic principal components for the demersal resources in areas E-F (South Kalimantan), ranked by their eigenvector; groups with similar eigenvectors tend to co-occur.

[Tabel 7. Struktur komponen pokok berdasarkan pembagian taksonomi dari sumberdaya demersal di daerah E-F (pantai Kalimantan Selatan) diklasifikasi berdasarkan nilai eigenvector; grup dengan nilai eigenvector yang berdekatan cenderung untuk muncul bersama.]]

Taxonomic group	PC-1 (48.7% of var.)		PC-2 (27.7% of var.)	
	Eigen-vector	Correlation ^a	Eigen-vector	Correlation ^a
Sciaenidae	0.23	0.99**	0.01	0.03
Polynemidae	0.22	0.93**	-0.03	-0.10
Clupeidae	0.22	0.93**	-0.00	-0.01
Sharks and rays	0.21	0.89**	0.06	0.18
Leiognathidae	0.21	0.88**	0.13	0.43
Ariidae	0.21	0.90**	0.06	0.19
Pomadasyidae	0.20	0.88**	-0.09	-0.29
<i>Lactarius lactarius</i>	0.20	0.88**	0.14	0.46
Scombridae	0.20	0.85*	0.09	0.29
Drepanidae	0.20	0.86*	-0.12	-0.39
Muraenesocidae	0.19	0.81*	-0.16	-0.51
Heterosomata	0.19	0.82*	0.10	0.33
Shrimps	0.19	0.82*	0.02	0.05
Cuttles	0.19	0.82*	0.15	0.50
Theraponidae	0.18	0.77	0.07	0.24
Synodontidae	0.18	0.78	0.17	0.56
Other food fish	0.16	0.70	0.12	0.40
<i>Formio niger</i>	0.16	0.70	0.14	0.46
Trichiuridae	0.15	0.65	-0.16	-0.53
Crabs	0.15	0.64	0.01	0.04
Lutjanidae	0.15	0.63	-0.18	-0.58
Squids	0.12	0.51	0.22	0.70
Carangidae	0.10	0.44	0.27	0.87*
Serranidae	0.10	0.41	-0.06	-0.18
Sphyraenidae	0.08	0.33	0.20	0.65
Pentapodidae	0.07	0.29	0.12	0.39
Other invertebrates	0.05	0.22	-0.27	-0.89**
Mullidae	0.02	0.09	0.26	0.84*
Trash fish	-0.03	-0.11	0.26	0.85*
Rachycentridae	-0.05	-0.20	-0.22	-0.70
Gerreidae	-0.05	-0.21	0.29	0.94**
Nemipteridae	-0.07	-0.30	0.28	0.89**
Chirocentridae	-0.13	-0.54	0.25	0.81*
Stromateidae	-0.13	-0.55	0.10	0.33
Engraulidae	-0.18	-0.76	0.12	0.39
Balistidae	-0.19	-0.81*	0.16	0.51
<i>Priacanthus</i> spp.	-0.19	-0.82*	0.11	0.34
<i>Thenus orientalis</i>	-0.20	-0.89**	0.14	0.44

* significant at $p < 0.10$; ** significant at $p < 0.05$.




Fig. 9. Trends in abundance of some taxonomic groups in areas E and F (South Kalimantan) showing high correlations with PC-1 and PC-2 (see also Table 7); A: positive correlations with PC-1; B: negative correlations with PC-1; C: positive correlations with PC-2; D: negative correlations with PC-2.

[Gambar 9. Perubahan kelimpahan beberapa grup ikan (menurut taksonomi) di daerah E dan F (Kalimantan Selatan) yang menunjukkan korelasi tinggi dengan PC-1 dan PC-2 (lihat Tabel 7); A: korelasi positif dengan PC-1; B: korelasi negatif dengan PC-1; C: korelasi positif dengan PC-2; D: korelasi negatif dengan PC-2.]

The patterns that were identified here differed from those detected in the Gulf of Thailand: the consistent declining trend for nearly all individual taxonomic groups reported by Pope (1979) did not occur in the present analysis. This may be attributed to:

- fishing pressure on the demersal resources off the north coast of Java and southeast coast of Sumatra was not as heavy as in the Gulf of Thailand;
- fishing operations in the Gulf of Thailand were very much dominated by bottom trawling, whereas bottom trawl catches contribute only 35-38% of the total demersal catches of northern western Java and southeast Sumatra (see above).

NORTH COAST OF CENTRAL AND EAST JAVA

The PCA indicates that 74.5% of the total variance is explained by the first two principal components (Table 6). Sharks and rays, Carangidae, Chirocentridae, Heterosomata, *Lactarius lactarius* and cuttles make a significant positive contribution to the first principal component, i.e., increased their abundance (Fig. 8A). The Engraulidae, Leiognathidae, Trichiuridae, shrimp and crab make a significant negative contribution, i.e., exhibited declining trends (Fig 8B).

For the second principal component, declining abundance trends were observed for the Mullidae, Muraenesocidae, Pentapodidae, *Priacanthus* spp. and *Thenus orientalis* (Fig. 8C), while *Formio niger* and Stromateidae increased (Fig. 8D). However, here again, more important groups declined, and thus the entire demersal group declined as well.

Table 8. Assumed trophic levels (TL) and relative catch rate (in %) of taxonomic groups in areas A-H, in 1976, ranked by (1) trophic level and (2) % of catch rates within each trophic level.

[Tabel 8. Jenjang rantai makanan (TL) dan laju penangkapan relatif (dalam %) dari grup taksonomi di daerah A-H, tahun 1976, diklasifikasi berdasarkan (i) jenjang rantai makanan dan (ii) % laju penangkapan dalam setiap jenjang rantai makanan.]

Taxonomic group	TL	A	B	C	D	E	F
Rays	4	7.85	10.50	6.88	3.05	3.66	5.94
Ariidae	4	0.60	4.71	2.51	6.04	6.16	15.58
<i>Priacanthus</i> spp.	4	2.40	1.86	4.82	24.23	0.37	0.11
Pomadasyidae	4	0.87	4.71	1.80	0.62	3.45	2.91
Synodontidae	4	2.72	3.09	3.86	2.11	0.80	1.43
Sharks	4	4.20	0.39	0.06	0.49	3.40	0.97
Sciaenidae	4	0.82	0.88	1.74	3.18	1.17	1.31
Sphyraenidae	4	0.00	1.23	0.96	0.36	5.25	0.68
Trichyuridae	4	0.22	1.28	1.35	0.75	1.17	0.51
Pentapodidae	4	0.49	0.10	0.00	0.52	0.27	0.06
Chirocentridae	4	0.33	0.59	0.06	0.00	0.16	0.29
Serranidae	4	0.05	0.20	0.00	0.03	0.53	0.06
Muraenesocidae	4	0.00	0.39	0.06	0.19	0.21	0.00
Rachycentridae	4	0.00	0.00	0.06	0.00	0.05	0.29
Mullidae	3	5.56	16.18	7.65	6.72	9.34	10.45
Carangidae	3	6.10	9.56	6.05	7.89	6.42	4.57
Trash fish	3	10.35	4.90	4.89	7.96	4.67	2.74
Nemipteridae	3	5.40	3.87	2.12	2.34	2.02	1.37
Squids	3	1.42	1.18	5.40	1.07	2.44	2.74
Other food fish	3	0.54	1.18	0.19	0.49	1.27	0.63
Terapontidae	3	1.25	1.08	0.58	0.13	0.53	0.57
Balistidae	3	0.54	0.25	0.39	1.43	0.80	0.23
Stromateidae	3	0.93	0.69	0.45	0.10	0.27	0.68
Drepanidae	3	0.00	0.15	0.06	1.30	1.43	0.06
Cuttles	3	0.27	0.59	0.45	0.32	0.11	0.34
<i>Lactarius lactarius</i>	3	0.00	0.34	0.71	0.26	0.11	0.46
Heterosomata	3	0.33	0.54	0.39	0.13	0.11	0.17
<i>Formio niger</i>	3	0.27	0.10	0.13	0.03	0.27	0.46
Polynemidae	3	0.33	0.05	0.00	0.03	0.64	0.06
<i>Thenus orientalis</i>	3	0.33	0.20	0.19	0.16	0.16	0.00
Crabs	3	0.05	0.05	0.19	0.03	0.00	0.00
Leiognathidae	2	31.61	8.68	28.17	19.78	14.54	15.64
Clupeidae	2	5.72	1.42	1.99	1.82	14.07	11.53
Gerreidae	2	6.49	13.68	3.34	1.46	3.50	4.11
Scombridae	2	0.71	2.26	4.82	5.29	1.38	1.88
Engraulidae	2	0.60	1.62	6.50	0.32	3.08	3.31
Lutjanidae	2	0.05	1.28	0.19	0.49	4.46	4.91
Shrimps	2	0.38	0.29	0.19	0.06	0.74	0.29
Other invertebrates	2	0.11	0.00	0.13	0.00	0.00	0.06
Total catch/effort (kg·hour ⁻¹)	(2.88) ^a	184	204	156	308	188	175

^aOverall mean weighted by the mean C/f over all areas.




Fig. 10. Trend of % contribution to total demersal biomass of the fish of three different trophic levels (TL = 2: preys; TL = 3: intermediate predators; TL = 4: top predators; see also Table 8) by area.
 [Gambar 10. Perubahan presentase kontribusi terhadap biomass total ikan-ikan demersal yang terbagi dalam beberapa kelompok rantai makanan (TL = 2 mangsa; TL = 3 pemangsa antara; TL = 4 pemangsa puncak, lihat Tabel 8) dan menurut daerah.]

Table 9. Number of taxonomic groups contributing significantly to the first two principal components (n), relative to the total number of groups at each trophic level (N), by area (A, C-D, E-F).

[Tabel 9. Jumlah grup taksonomi pendukung penting dua komponen pokok awal (n), dibandingkan dengan jumlah total grup dalam tiap jenjang rantai makanan (N) berdasarkan daerah (A, C-D, E-F).]

Trophic level/area	A		C-D		E-F	
	n	N	n	N	n	N
2 (prey fish)	4	8	3	8	5	8
3 (intermediate predators)	7	17	9	17	9	17
4 (top predators)	5	13	6	13	6	13

SOUTH COAST OF KALIMANTAN

Although surplus-production analysis could not be applied to the sparse data from the south coast of Kalimantan, PCA was performed to examine seasonal variations of the stocks. The results indicate that 76.4% of the variation of the total demersal resources can be explained by the first two principal components (Table 7). Significant positive contributions to PC-1 were made by the sharks and rays, Ariidae, Clupeidae, Drepanidae, Heterosomata, *Lactarius lactarius*, Leiognathidae, Muraenesocidae, Polynemidae, Pomadasyidae, Sciaenidae, Scombridae, cuttles and shrimp (Fig. 9A), while negative contributions were made by the Balistidae and *Thenus orientalis* (Fig. 9B).

PCA AND FUNCTIONAL GROUP ANALYSIS

The foregoing shows that PCA could be used to reduce much of the multivariate data matrix of trawl catches to two principal components, and to identify taxonomic groups that made significant contributions to the change in abundance of the demersal group as a whole.




Fig. 11. Plots of % contribution to total demersal biomass of various trophic levels against second principal component scores in two areas of the North Java coast: left: area C, 1975-1979; right: area D, 1976-1979.
 [Gambar 11. Grafik persentase kontribusi terhadap biomass total ikan-ikan demersal dari berbagai tingkat rantai makanan terhadap nilai komponen utama yang kedua di dua daerah utara Jawa: kiri: daerah C, 1975-1979; kanan: daerah D, 1976-1979.]




Fig. 12. Rank correlations among areas, from 1975 to 1979 and among years, for various areas, of the taxonomic components in Java Sea trawl survey catches (all significant at $p < 0.01$).

[Gambar 12. Korelasi berdasarkan ranking antar daerah, dari tahun 1975 hingga 1979, dan antar tahun, untuk berbagai daerah, dari grup taksonomi di Laut Jawa (semua nyata pada $p < 0.01$).]




Fig. 13. Surplus-production analyses of the demersal fisheries along the Southeast Sumatra and North Java coasts (areas A-D), 1975-1979, using different indices of catch/effort and effort: Index - 1: based on catch (from fisheries statistics)·trawler⁻¹·year⁻¹; Index - 2: based on trawl survey catch·hour⁻¹; Index - 3: based on transformed trawl survey catch·hour⁻¹ (see also Fig. 6).

[Gambar 13. Analisis surplus produksi dari perikanan demersal didaerah tenggara Sumatra dan pantai utara Jawa (daerah A - D), 1975-1979, dengan beberapa indeks hasil tangkapan per upaya. Indeks - 1 : berdasarkan hasil tangkapan (data statistik perikanan)·kapat⁻¹·tahun⁻¹; Indeks - 2 : berdasarkan hasil tangkapan trawl (survei)·jam⁻¹; Indeks - 3 : berdasarkan transformasi hasil tangkapan trawl·jam⁻¹ (lihat Tabel 6).]

Table 10. Correlation among pelagic, demersal, total catch and principal component (1 or 2)^a.

[Tabel 10. Korelasi antara hasil tangkapan sumberdaya pelagis, demersal, total hasil tangkapan dan komponen pokok^a.]

Dependent variable	PC	Areas A-D	Areas C-D	Areas E-F
Pelagic group	1	-0.87* (-0.97)**	0.29 (0.24)	0.89** (0.92)**
	2	-0.26 (-0.22)	-0.49 (-0.87)*	0.34 (0.17)
Demersal group	1	0.78* (0.84)*	0.51 (-0.27)	0.87* (0.98)**
	2	0.59 (0.47)	0.85* (0.85)*	0.38 (0.08)
Total catch	1	0.67 (0.70)	-0.50 (-0.25)	0.89** (0.99)**
	2	0.74 (0.57)	0.84* (0.74*)	0.38 (0.10)

^aValues within brackets were based on 17 taxonomic groupings; * significant at $p < 0.10$; ** significant at $p < 0.05$.




Fig. 14. Surplus-production analyses of the demersal fisheries along northern coast of Central and East Java (areas C-D), 1975-1979, using different indices of catch/effort and effort: Index - 1: based on catch (from fisheries statistics)-trawler⁻¹·year⁻¹; Index - 2: based on trawl survey catch·hour⁻¹; Index - 3: based on transformed trawl survey catch·hour⁻¹ (see also Fig. 6).

[Gambar 14. Analisis surplus produksi dari perikanan demersal di pantai utara Jawa Tengah dan Timur [daerah C-D], 1975-1979, dengan beberapa indeks hasil tangkapan per upaya. Indeks - 1: berdasarkan hasil tangkapan (dari data statistik perikanan)-kapal⁻¹·tahun⁻¹; Indeks - 2: berdasarkan hasil tangkapan trawl (survei)-jam⁻¹; Indeks - 3: berdasarkan transformasi hasil tangkapan trawl·jam⁻¹ (lihat Tabel 6)].

The present study was, unfortunately, based on only five years' worth of data. Still, it was possible to identify those taxonomic groups which significantly contributed to the total variation of demersal groups in different areas experiencing different degrees of fishing pressure. The percentage contribution of the first two principal components was around 75% of total variation both for the lightly exploited areas (areas E and F) and the highly exploited areas (areas C and D). However, there was a large difference in the number of taxonomic groups significantly contributing to the first two principal components. Whether this difference was caused by differences in fishing pressure, by interspecific interactions among taxonomic groups, or by a combination of these cannot be resolved.

An attempt was also made, based on literature data, to assign trophic levels (TL) to the various taxonomic groups, i.e., prey fish (TL = 2), intermediate predator (TL = 3) and top predator (TL = 4), i.e., assuming small prey fish to feed directly on primary producers and on detritus (for which TL = 1 in both cases, see contributions in Christensen and Pauly 1993). The hypothesis to be tested here was whether the fishes of different trophic levels (Table 8) to the total variance in three areas were exposed to different degrees of fishing pressure.

No significant difference was detected (using an χ^2 analysis) for any of the trophic levels (Table 9).

Another way to test whether the stocks underwent changes in trophic structure over time is through an examination of the trophic level contribution to the time series. Fig. 10 shows that, in general, only minor changes of trophic level contribution to overall biomass occurred, except in Area D, where only a

small number of trawl hauls were made in 1978. This suggests that increased fishing, at least from 1975 to 1979, has had only a minor effect on the trophic structure of the demersal communities of the Java Sea, as also confirmed by the lack of trends in Fig. 11, and by Fig. 12, documenting the stability of the ranking of different taxa among years and areas.

PCA AND SURPLUS-PRODUCTION MODELS

The entire "demersal" catch has been used for the surplus-production analyses documented in Figs. 13 and 14. These indicate that overall fishing effort was, in the late 1970s, not in excess of that needed to generate MSY, irrespective of which measures of catch and effort are used.

Table 10 shows that the "demersal" groups have, in Areas A-D, C-D, and E-F, significant positive correlations with PC-1 and PC-2. Significant positive correlations also occur between the "total catch" and the principal components in Areas C-D and E-F. The latter appears to be caused by a strong negative correlation between the "pelagic group" and the principal components.

In general, the "pelagic group" is negatively correlated with the principal components (except in Areas E-F where the pelagic fishing effort is slight) which may be attributed to a significant contribution of some pelagic fishes to the variation of the total catch (see above).

Application of the surplus-production model to total biomass of the demersal resources in the tropics has generally been based on the index of abundance derived from the total catch rate of a bottom trawl, as done in the analysis of demersal

resources in the Gulf of Thailand (FAO 1978), and those off the north coast of Java (Dwiponggo 1978). Since some pelagic fishes are represented in the trawl catch, the estimate of MSY for the demersal resources will obviously include the pelagic component. Thus, unless the pelagic group represents a small portion of the total catch, estimates of MSY for the trawl fishery must be taken with a grain of salt, particularly when the pelagic group is also negatively correlated with the principal component, as in the present analysis.

Overall, it would thus seem best, when dealing with catch and effort data from bottom trawl fisheries, to separate the truly demersal fishes from the pelagics caught incidentally, and which can be defined both in taxonomic/ecological terms, and through their correlation with the first and second components of a PCA.

Acknowledgements

This contribution was extracted from my PhD thesis, submitted in July 1982, entitled "Fishery dynamics of the demersal resources of the Java Sea" and supervised by Prof. A. Hanson at Dalhousie University, Halifax, Canada. I take this belated opportunity to thank all involved in this venture, especially Prof. Hanson, and also Dr. J. McManus (ICLARM) for a thorough review of the draft of this contribution.

References

- Baum, G.A. 1978. A cost/benefit calculation for a "Bagan Siapi-api" purse seiner operating from Pekalongan (Central Java). Contributions to the Symposium on Modernization of Small-scale Fisheries, 27-30 June 1978, Jakarta, Indonesia. 15 p. (Mimeo).
- Berlage, H.P. 1927. Monsoon-currents in the Java Sea and its entrances. Konin. Magnet. Met. Observ. Batavia Verhandelingen 19: 1-28.
- Birowo, S., A.G. Iahude and A. Nontji. 1975. Status pengetahuan dalam ilmu Laut di Indonesia, p. 1-79. In A. Soegiarto and S. Birowo (eds.) Atlas oseanologi perairan Indonesia dan sekitarnya, LON-LIPI. Buku I.
- Bliss, C.I. 1967. Statistics in biology: statistical methods for research in the natural sciences. Vol. I. McGraw-Hill, New York.
- Budihardjo. 1978. Analisa ekonomi perikanan trawl type Bagan di Java, suatu studi kasus. Contributions to the Symposium on Modernization of Small-scale Fisheries, 27-30 June 1978, Jakarta, Indonesia. 30 p. (Mimeo).
- Christensen, V. and D. Pauly, Editors. 1993. Trophic models of aquatic ecosystems. ICLARM Conf. Proc. 26, 390 p.
- DGF (Directorate General of Fisheries). 1981. Fisheries statistics of Indonesia (1977-1981). Directorate General of Fisheries, Ministry of Agriculture, Jakarta, Indonesia.
- Doty, M.S., R.E. Soeriatmadja and A. Soegiarto. 1963. Observations on the primary productivity of northwestern Indonesian waters. Mar. Res. Indones. 5: 1-25.
- Dwiponggo, A. 1978. Status perikanan demersal diperaian pantai Laut Jawa, potensi dan tingkat peugusahaannya. Contributions to the Symposium on Modernization of Small-scale Fisheries, 27-30 June 1978, Jakarta, Indonesia. 13 p. (Mimeo).
- Dwiponggo, A. and M. Badrudin. 1980. Data of the Java Sea inshore monitoring survey by *Mutiara IV*. Mar. Fish. Res. Rep. Contrib. Demersal Fish. Proj. 7A, 88 p.
- Emery, K.D. 1969. Distribution patterns of sediments on the continental shelves of Western Indonesia. ECAFE Tech. Bull. 2: 79-82.
- FAO. 1978. Some scientific problems of multi-species fisheries. FAO Fish. Tech. Pap. 181, 42 p.
- Gulland, J.A. 1961. Fishing and the stocks of fish at Iceland. Fish. Invest. Minist. Agric. Fish. Food UK (Ser. 2) 23(4):1-52.
- Gulland, J.A. 1971. The fish resources of the ocean. Fishing News (Books) Ltd., Surrey, England. 255 p.
- Losse, F.G. and A. Dwiponggo. 1977. Report on the Java Sea southeast monsoon trawl survey, June - December 1976. Mar. Fish. Res. Rep., Contrib. Demersal Fish. Proj. 1, 119 p.
- Martosubroto, P. 1978. Perkembangan perikanan semi-industri di pantai Utara Jawa Tengah dan pengaruhnya terhadap perikanan rakyat. Contributions to the Symposium on Modernization of Small-scale Fisheries, 27-30 June 1978, Jakarta, Indonesia. 16 p. (Mimeo).
- Martosubroto, P. and D. Pauly. 1976. *R.V. Mutiara IV* survey data, November 1974 - July 1976. Mar. Fish. Res. Rep., Contrib. Demersal Fish. Proj. 2, 136 p.
- May, R.M., J.R. Beddington, C.W. Clark, S.J. Holt and R.M. Law. 1979. Management of multi-species fisheries. Science 205: 267-277.
- Naamin, N. and J. Uktolseya. 1976. Status perikanan udang diperaian Kalimantan Selatan dan Timur (with English abstract). Laporan Penelitian Perikanan Laut 2: 1-82.
- Pauly, D. 1979. Theory and management of tropical multi-species stocks. ICLARM Stud. Rev. 1, 35 p.
- Pauly, D. 1984. Fish population dynamics in tropical waters: a manual for use with programmable calculators. ICLARM Stud. Rev. 8, 325 p.
- Pauly, D. 1987. Theory and practice of overfishing: a Southeast Asian perspective, p. 146-163. In Proceedings of the 22nd Session of the Indo-Pacific Fisheries Commission, 16-26 February 1987, Darwin, Australia. RAPA Rep. 10. FAO Regional Office for Asia and the Pacific, Bangkok.
- Pielou, E.C. 1977. Mathematical ecology. John Wiley and Sons, New York. 385 p.
- Pope, J.G. 1979. Stock assessment in multispecies fisheries, with special reference to the trawl fishery in the Gulf of Thailand. SCS/DEV.79/19, 106 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191, 382 p.
- Saeger, J., P. Martosubroto and D. Pauly. 1976. First report of the Indonesian-German Demersal Fisheries Project. (Results of a trawl survey in the Sunda Shelf area). Mar. Fish. Res. Rep., Contrib. Demersal Fish. Proj. 1: 1-16.
- Sardjono, I. 1980. Trawlers banned in Indonesia. ICLARM Newsl. 3(4): 3.
- Schaefer, M.B. 1954. Some aspects of the dynamics of populations important to the management of commercial marine fisheries. Bull. I-ATTC 1(2): 26-56.
- Schaefer, M.B. 1957. A study of the dynamics of the fishery for yellowfin tuna in the eastern tropical Pacific Ocean. Bull. I-ATTC 2(6): 247-285.
- SCSP. 1979. Report of the Workshop on Demersal and Pelagic Fish Resources of the Java Sea, 5-9 December 1978, Semarang. SCS/GEN/79/20, 15 p. South China Sea Fisheries Development and Coordinating Programme, Manila.
- Sneath, P.H.A. and R.R. Sokal. 1973. Numerical taxonomy: the principles and practice of numerical classification. W.H. Freeman Co., San Francisco, California. 573 p.
- Soeriatmadja, R.E. 1956. Seasonal fluctuations in the surface salinity off the north coast of Java. Mar. Res. Indones. 1: 1-19.
- Sokal, R.R. and F.J. Rohlf. 1969. Biometry. The principles and practice of statistics in biological research. Second ed. W.H. Freeman Co., San Francisco, California. 859 p.
- Sjarif, S. 1959. Seasonal fluctuations in the surface salinity along the coast of the southern part of Kalimantan (Borneo). Mar. Res. Indones. 4:1-23.
- Steel, R.G.D. and J.H. Torrie. 1960. Principles and procedures of statistics, with special reference to the biological sciences. McGraw-Hill, New York.
- Sujastani, T. 1978. Penhitungan besarnya stock sumber-sumber perikanan di Laut Jawa berdasarkan data statistik perikanan daerah. Contributions to the Symposium on Modernization of Small-scale Fisheries, 27-30 June 1978, Jakarta, Indonesia. 15 p. (Mimeo).
- Tiews, K. 1966. On the possibilities for further developments of the Southeast Asian fisheries. IPFC Curr. Aff. Bull. 47: 1-13.
- van Riehl, P.M. 1932. The Snellius expedition. J. Cons. CIEM 7: 212-217.
- Wyrtki, K. 1961. Physical oceanography of the Southeast Asian waters. Scripps Inst. Oceanogr., Naga Rep. 2, 195 p.
- Zachman, N. 1973. Fisheries development and management in Indonesia. J. Fish. Res. Board Can. 30 (12/2): 2335-2340.

Narrative and Major Results of the Indonesian-German Module (II) of the JETINDOFISH Project, August 1979 to July 1981

UWE LOHMEYER

*Deutsche Gesellschaft für Technische
Zusammenarbeit
GmbH, Eschborn, Germany*

LOHMEYER, U. 1996. Narrative and major results of the Indonesian-German Module (II) of the JETINDOFISH Project, August 1979 to July 1981 [*Sekilas tentang hasil pokok kerjasama Indonesia-Jerman (Modul II) dari proyek JETINDOFISH, Agustus 1979 hingga Juli 1981*], p. 77-90. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

An account is presented of a series of Indonesian-German trawl survey cruises conducted from August 1979 to July 1981 in the framework of the multiagency "Joint Eastern Indian Ocean Fisheries Survey" (JETINDOFISH).

The area surveyed - mainly through demersal trawling - covered 70,000 km² of shelf from the central coast of Sumatra to Java and Bali.

The survey and sampling methodologies are described in some details, such as to allow replication of estimates of what was, during the survey period, a largely unfished biomass.

Abstrak

Sekilas pemaparan tentang hasil-hasil pokok dari rangkaian survei Indonesia-Jerman dari bulan Agustus 1979 hingga Juli 1981 dalam rangka kerjasama antar beberapa instansi yang dikenal dengan nama "Joint Eastern Indian Ocean Fisheries Survey (JETINDOFISH)".

Daerah yang disurvei - terutama melalui trawling sumberdaya ikan demersal - meliputi wilayah seluas 70,000 km², yakni daerah paparan perairan dari pantai tengah Sumatra hingga ke Jawa dan Bali.

Metode survei dan cara pengambilan contoh diketengahkan secara rinci sedemikian rupa sehingga memungkinkan replikasi penghitungan estimasi dalam kurun waktu survei terhadap biomassa sumberdaya yang mana pada waktu survei dilakukan sebagian besar masih belum tereksplorasi.

Introduction

The work reported upon here was performed within the framework of the "Joint Eastern Indian Ocean Fisheries Survey" (JETINDOFISH), a project conceptualized by members of the Indian Ocean Fishery Executive Committee during meetings held in Rome and Mombasa in 1974. During these workshops, it became apparent that there was very little information available on fish stocks of the eastern and southeastern Indian Ocean, i.e., south of the Indonesian archipelago and on the northwestern shelf of the Australian continent.

The committee and representatives of the Food and Agriculture Organization (FAO) and the United Nations Development Program (UNDP) therefore agreed to launch a major exploratory fishing survey in the area, based on the assumption that it contained large resources so far not utilized because of the very narrow shelves, unsuitable trawling grounds, rough seas, inaccessible coasts, and lack of fisheries

support facilities and markets.

This was especially applied to the southern waters of Indonesia which, by that time, had only a limited fishery on pelagic stocks. These were thought to be abundant, as suggested by the occurrence of a small upwelling in the Bali Strait, then thought to extend well beyond that area (Cushing 1971). Evidence to the contrary (Venema 1976) came too late to affect this project (see Venema, this vol.).

The project was implemented in three phases:

The Preparatory Phase. This was funded and carried out in 1976 by the Indian Ocean Programme (IOP) of the Indian Ocean Fisheries Commission, with assistance from the Free Hanseatic City of Bremen (Federal Republic of Germany).

The Survey Phase. Contrary to original plans, UNDP was in no position to fund this phase, and therefore, a multinational effort was undertaken to find funding for the project. This led to the Governments of Indonesia and Australia agreeing to embark on a major joint fishery survey in eastern Indonesia and northwestern Australia. Also, the Government

of Indonesia received assistance to conduct a survey through a technical cooperation project from the Government of the Federal Republic of Germany, in the form of a chartered research vessel, and of personnel and equipment. The implementing agencies were to be the Directorate General of Fisheries (DGF), Ministry of Agriculture, on the Indonesian side, and the Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ), GmbH (German Agency for Technical Cooperation) on the German side. This project phase lasted from August 1979 to July 1981.

The **Evaluation Phase**. This stage was for the data to be analyzed and a first set of results to be published. Subsequent to this, recommendations derived from detailed analysis of the data obtained in the course of the surveys were to be provided concerning the improved utilization of the resources, and various pilot projects were to be implemented to increase catches, to improve processing and marketing methods and facilities, and to develop the fisheries infrastructure.

Coordination and standardization of the survey activities were entrusted to FAO, which seconded coordinators (Dr. W.

Brandhorst, later Dr. T. White).

This final phase was not implemented as anticipated. Particularly, a software package previously developed for analysis of trawl data in the Persian Gulf failed to perform, and only averaged catch rates (by area, by depth, etc.) became available for analysis and reporting before the project ended, hence this contribution and the others in this book (see also Pauly, this vol.).

Materials and Methods

The Surveys

The area surveyed by the JETINDOFISH project stretches from Sumatra to the northwestern shelf of the Australian continent. This area was subdivided into three "Modules". One of these (Module II) extended from Sumatra to the degree of longitude that crosses the island of Bali (Fig. 1). The fisheries research vessel, *FRV Jurong*, with a displacement of 300 GT and LOA of 30 m was commissioned




Fig. 1. Definition of that part of the Indian Ocean coast of Indonesia covered by areas 3-7, Module II of JETINDOFISH. Areas 1 and 2, as initially defined, were not surveyed and are not shown.

[Gambar 1. Bagian dari Samudra Hindia wilayah Indonesia (area 3-7), JETINDOFISH Modul II. Daerah 1 dan 2, sebagaimana dijelaskan sebelumnya, tidak diliput didalam survei, oleh karena itu tidak ditampilkan ini.]

to conduct the survey in Module II.

Module I was located to the east of Module II, and extended to the northwestern shelf of the Australian Exclusive Economic Zone in the Arafura Sea. Module I was surveyed by *FRV Bawal Putih 2* (see Martosubroto, this vol.).

The Australian northwestern shelf (Module III) was first surveyed by *FRV Courageous*, later by *FRV Soela*. Both vessels were made available by the Australian Ministry of Primary Industry.

Modules I, II and III were surveyed independently from each other; however, most activities and sampling protocols were standardized by an Advisory and Technical Committee composed of members of all agencies involved in the JETINDOFISH project. All modules were subdivided into "areas". Only the results of the survey performed in Module II are presented here.

The main objective of the survey was to gather data for an assessment of stock density. This was to be achieved by collecting catch/effort data and combining these with readings from echointegrators, as gathered during acoustic surveys.

It was clear from the beginning of the project that the vast expanses of the survey area and the limited time available would allow to take samples only at a relatively small number of stations. It was assumed that this constraint would curtail the accuracy and precision of the findings, but would yield at least a rough estimate of stock density and abundance, sufficient for a decision on whether or not to encourage the development of commercial fisheries, and possibly to decide on some of their technical features.

In addition, an attempt was made to describe the hydrography of the area by collecting data from a bathythermograph (BT) and by regularly recording sea surface temperatures (SST).

The survey was also expected to help answer the

question whether the acoustic equipment installed on board the vessel was a cost-efficient tool that could be employed to explore and monitor fish stocks. The answer to this question was of particular importance since a portion of the resource was already subjected to exploitation through established, if ill-documented artisanal fisheries (see Venema, this vol., and Ghofar and Mathews in Pauly et al., this vol.).

The Survey Vessel

FRV Jurong, used for surveying Module II, had the characteristics of a stern trawler (Box 1). This vessel was formerly commissioned as a fisheries training vessel and was chartered for the survey by GTZ from the UNDP/FAO vessel pool.

Given its original design and previous utilization, it had sufficient space to accommodate the crew and a number of scientists and technicians, and to process catch samples according to established scientific standards. The vessel's engine, rated at 555 kW, was adequate to tow the bottom trawl at speeds of up to 5 kn. With increasing depth, trawling speed however became reduced; at depths of more than 250 m the standard speed of 3 kn could not be maintained, and at the maximum fishing depth (380 m), the trawling speed was only 1.2 kn.

The vessel was of a shelter-decked type. The catch was hauled in via a stern ramp to the upper deck, then cleaned and brought forward to a hatch from where it came down to the steerage and processing deck, where the fish were sorted and recorded. Individual fishes suitable for processing were preserved, i.e., frozen in the deep-freezing compartment. Navigational equipment included a gyro compass, radar, echosounder, sonar, and satellite and radio navigation systems.

The vessel had a full complement of Indonesian officers and crew. The German counterparts consisted of a captain/masterfisher, a chief mate, two engineers who trained their counterparts in vessel operations and fishing techniques, and an electronics specialist responsible for operation and maintenance of the electronic equipment. Another land-based expatriate, a vessel operation manager, was responsible for logistics.

Three Indonesian fisheries biologists and three technicians were permanently assigned as counterparts to the German team. In addition, a total of 20 Indonesian biologists, technicians and one student received hands-on training during 32 of the cruises.

The FAO/UNDP coordinator participated in several cruises with the main objective of standardizing the data collection according to the criteria set by the coordinating FAO/UNDP body. During three cruises, a total of 11 guests and short-term consultants joined the survey, particularly to assist in the taxonomic work.

Box 1. Vessel specifications of *R/V Jurong*.

[Boks 1. Spesifikasi kapal penelitian *Jurong*.]

Type stern trawler, Class + 1A	
Main dimensions:	
Length O.A.	35.0 m
Length B.P.P.	31.7 m
Breadth MLD	7.9 m
Tonnage	213.25 BRT
Capacity:	
Freshwater tanks	27 m ³
Fuel oil tanks	57 m ³
Lubricant oil tanks	2 m ³
Tunnel freezer	-30°C (3 t day ⁻¹)
Freezer	-18°C
Engine and performance	
Main Engine CATERPILLAR Type D 398 TEF	
750 HP	1225 RPM
(controllable pitch propeller)	
Max. speed	12.5 kn
Main fishing winch	Hydraulik A/S Dia 10/4
	Wire capacity 1400 m; ϕ 18 mm

Box 2. Navigation and echosounding equipment of FRV Jurong.

[Boks 2. Peralatan navigasi dan echosounding dari FRV Jurong.]

Navigation:

DECCA Radar D 202
KELVIN HUGHES Radar 18/12
MAGNAVOX Satellite Navigator MX 1102 - NV
SIMRAD Sounder EQ 50
TOKYO KEIKI Gyro-Compass
BERGEN NAUTIK Magnetic Compass
KELVIN HUGHES Pentland Bravo Radio DRG
KELVIN HUGHES Foreland VHF Radion 801 A
BERGEN NAUTIK Electric Log

Echosounding:

SIMRAD Scientific Sounder EK 38 (38 kHz)
SIMRAD Scientific Sounder EK 120 (120 kHz)

SIMRAD SK 3 Sonar

SIMRAD CI - Scope connected to the sounder, and to a SIMRAD MC
- Magnifier.

SIMRAD Echo Integrator QM - MK II
HEWLETT PACKARD 7702 B Recorder

SIMRAD EX Netsonde

Test and calibration equipment:

HEWLETT PACKARD 141 B Storage Oscilloscope
WAVETECK 116 Signal Generator
Frequency Counter
Voltmeter
Hydrophone

KAHL Bathythermograph (0-250 m)
KAHL 250 WA 100 Meter Wheel

The Electronic Equipment

A sonar and a Simrad QM-MK II echointegrator were used for locating and assessing fish aggregations (Box 2). Basically, the instruments consists of a specially designed transducer coupled to an electronic processor. All electroacoustic equipment, with exception of the sonar and the net sonde, were kept in *Jurong's* airconditioned chart room for protection against excessive humidity. For fish detection, either one of the two scientific sounders - SIMRAD EK 38 (operating at 38 KHz) and SIMRAD EK 120 (120 KHz) - or the navigational sounder, SIMRAD EQ 50 KHz was operated. For actual sampling, the EK 38 was used because the shortwave oscillations of the EK 120 had a depth range limited to about 100 m. During operation of the integrator, the types and spatial distributions of echoes were logged in form of echograms. Thus, it was possible to directly associate the integrator readings with the characteristics of the echoes.

The echosounder converts into electric impulses the echoes that are received by the transducer from individual fish or fish aggregations. The integrator then sums these impulses, while compensating for distance and other variables. Readings from the latter instrument were assumed proportional to the biomass of the particular fish aggregations in question.

An MC-Magnifier screen was attached to the integrator to display simultaneously and continuously both fish aggregations and signal strengths. Detailed analysis of individual signals was performed using a fish scope. The navigational echosounder working at a frequency of 50 kcycles was used during trawling operations and during acoustic surveys to detect concentrations of fish at the shelf's margin and slope. However, when switched on for fish detection in the surface layer, the sonar (SIMRAD SK 3) often caused interference with the echosounder. One unit each of oscilloscope, signal generator, frequency counter, voltmeter and hydrophone were used to test and calibrate the electroacoustic equipment.

Individual fish in the vicinity of the net opening could be detected by a net sonde. This instrument, however, was attached to the headrope only when the risk of losing gear and instrument was low, i.e., when the bottom was smooth.

The vessel was also equipped with a winch for a bathythermograph: recordings to depths of 200 m were made regularly throughout the survey.

The Demersal Trawl

A high-opening bottom trawl of 527 meshes circumference with 200 mm (stretched) mesh was used, with a rather large codend mesh size of 40 mm (Fig. 2). The headline carried floats of 20-cm diameter; the ground rope was mounted alternately with rubber discs of 30- and 50-cm diameter at a spacing of approximately 50 cm, and fitted with chains. The dry weight of the ground rope was approximately 675 kg. The bridles were 53-m long at 14-mm diameter. The vertical net opening at a trawling speed of 3 kn was 4.5 m. The horizontal net opening was 25 m on the average, depending on the nature of the sea bottom, and was measured by gauging the angle of spread of the warplines. The area swept during a one hour tow at 3 kn was estimated at 0.1389 km².

The otter boards measured 2.6 x 1.5 m, were oval-shaped, had interchangeable steel shoes, and weighed between 600 and 650 kg each.

The Pelagic Trawl

The four-seam midwater trawl used during the surveys had 308 meshes circumference at 800 mm (stretched mesh), was made of 100% high-quality nylon twine for optimum elasticity and was designed for the capture of small pelagic fishes (Fig. 3, inset). Sets of five floats each were mounted to the headrope for static buoyancy. The codend had a mesh size of 40 mm (stretched mesh) and was equipped with a splitting strap. Bridles were 70-m long at 14-mm diameter. Steel weights of 300 kg each were attached to the lower tips of the wings in order to give the mouth opening of the net a more or less permanent rectangular shape. The otterboards were 2.55 x 1.5 m in size, symmetrical and could be used on either side of the ship.




Fig. 2. Schematic representation of the otter trawl used for sampling demersal fish.
 [Gambar 2. Diagram otter trawl yang dipakai untuk pengambilan contoh ikan demersal.]




Fig. 3. Pelagic trawl stations covered by Module II of JET:INDOFISH (inset: the sampling gear).
 [Gambar 3. Stasiun trawl pelagis dari JET:INDOFISH Modul II (sisipan: gambar trawl, alat tangkap yang digunakan).]




Fig. 4. Cumulative cruise tracks of *R/V Jurong*, as followed during the combined acoustic, trawling and hydrographic surveys lasting from August 1979 to July 1981. Note rectangular shapes, allowing even coverage of the shelf, from inshore to deeper waters.

[Gambar 4. Liputan kumulatif pelayaran kapal *Jurong* selama kegiatan survei terpadu yang menggunakan peralatan akustik, trawl dan hidrografi, yang berlangsung dari bulan Agustus 1979 hingga Juli 1981. Perhatikan garis pelayaran yang menyerupai persegi panjang, menandakan liputan paparan yang merata dari perairan dangkal hingga ke perairan dalam.]

Some 154 midwater trawl operations were performed during the survey (Fig. 3). Cruise tracks were determined ahead of each voyage. Sailing along these tracks and with the echointegrator continuously running, large fish aggregations were supposed to be caught once they were detected by the sonar and/or the sounders. Whenever bigger schools of fish were detected, the net was shot regardless of the hour of the day. However, only a very limited number of such hauls were made due to the absence of large fish concentrations.

A standard procedure of midwater sampling was implemented to routinely verify readings received by the integrator. These so-called "standard fishing operations" were carried out early mornings, at noon and late in the afternoon and/or during night time. The main purpose of these operations was to sample the upper 8-m layer of the water column since this horizon could not be surveyed by the electroacoustic equipment. Once the net was hauled in, the vessel returned to the position where the net was shot and continued with the acoustic survey, following the predetermined cruise tracks. Nearly 12,000 nm of tracks were covered this way, twice from Bali up to Nias Island, and once each during the southeast and northwest monsoon periods (Fig. 4).

Fish Species Identification

Species identification required a considerable effort. There were quite a number of identification keys dealing with the fish fauna of the tropical Indo-Pacific. However, with a few exceptions (see below), these works were not appropriate for shipboard use. Neither was a single comprehensive key available which would have allowed for reliable identification of the species likely to be encountered in the survey area. Quick and correct identification was therefore very difficult. Most shipboard identifications were based on the *FAO species identification sheets for fishery purposes: Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71)* edited by Fischer and Whitehead (1974); *Coastal fishes of South Japan* (Masuda et al. 1980), *Coastal fishes of New Guinea* (Munro 1967) and *The fishes of the Indo-Australian Archipelago* (Weber and de Beaufort 1913, 1916, 1922, 1929, 1931, 1936).

Doubtful or unknown species were marked with a temporary code number, preserved and sent to a specialist. In all cases, this procedure ensured an identification down to the species level. Also, almost all species were photographed and

one reference specimen each was preserved for later referral. Thus, a project-specific "field guide" emerged which gradually replaced the other identification works. This "field guide" was later published as *Trawled fishes of southern Indonesia and northwestern Australia* (Gloerfelt-Tarp and Kailola 1984).

Trawl Sampling

The *Jurong's* draught of 7.8 m precluded an investigation of nearshore waters, lagoons and reefs. On the other hand, the equipment installed on board did not permit sampling of deep water zones: for technical reasons, the echointegrator was calibrated to depths ranging between 15 and 100 m.

Likewise, the biomass of pelagic fish species near the sea surface could not be assessed reliably. Fast swimming pelagic species near the surface avoided both the trawl gear and acoustic detection. Exception to this were extensive aggregations of *lemuru* (*Sardinella lemuru*) found in Bali Strait. These caused strong signals in the echosounders and were very well sampled by the high opening bottom trawl.

FRV Jurong was commissioned for the survey of Module II for two years, from August 1979 to July 1981. During this period, 32 research cruises and 743 fishing operations were conducted, 154 of which with the pelagic trawl. The data from 515 of the remaining 592 (bottom) trawl hauls were considered to be suitable for purposes of stock assessment; 77 hauls were conducted as "simulated commercial" operations, i.e., their location was nonrandom.

For each area, the catch/effort data were used to estimate standing stock size. The decision on where fishing stations should be located was done in random fashion, as follows: prior to the survey, each area was subdivided into squares of 5 x 5 nm, and the squares numbered consecutively. Using a random number generator, the squares to be investigated were then identified. When the vessel actually visited the identified square and the echosounder indicated that the sea bottom would be too rough to allow smooth trawling, that particular square was marked in the sea chart, and the next square previously selected by the random process was visited. Later analysis of echosounder recordings, including those from untrawable grounds, were used to classify and document the bottom topography according to its general suitability for trawl fishing.

In certain parts of the survey area - e.g., in Grajagan Bay, on the southeast coast of Java - trawable grounds were extremely scattered and could not be sampled at random. In such cases a systematic search for smooth bottoms was conducted with trawling operations made whenever and wherever possible. Data from these hauls were not used for the estimation of biomass.

"Simulated commercial" trials with the bottom trawl were made up to three times per cruise. In these cases, the decision on where and how long to fish was left to the captain. Quite often these commercial trials were made where random fishing operations had indicated an abundance of commercially valuable species. Sampling of those catches followed the

standard procedure; however, the data and results originating from these operations were also kept separate from those used for biomass assessments.

Cruise No. 24 served for intercalibration of fishing gear between *FRV Jurong* and *FRV Bawal Putih 2*. Both vessels attempted to catch as much fish as possible under simulated commercial conditions. Their fishing stations were therefore not selected at random, and the data thus gathered were likewise not used for biomass assessments. The operations included target fishing for deep water resources (<< 200 m), especially for deep sea shrimp.

Catch sampling followed the standard procedures described in Losse and Dwiponggo (1977; see also Pauly 1983). The presorted catches were weighed, and subsamples were recorded according to species and weight.

In cases where the standard time schedule of one hour trawling could not be adhered to, the weights of that haul and of its component species were corrected by applying the appropriate multiplicative factor(s). Trawling speed and net opening were usually assumed to remain constant, implying an equal average volume of water being filtered by the net and an equal bottom area being swept (see below for cases where haul-specific statistics were used to estimate the areas swept).


Whenever time and resources permitted, length-frequency measurements were taken. Total lengths were recorded when applicable; in cases where the caudal fin was heterocercal or where the tail end allowed no clear definition of total length, standard lengths were taken.

Assessment of Biomass

The catch/hour data were divided by the area swept to obtain estimates of relative density ($t \cdot km^{-2}$), which were averaged within previously defined depth ranges and seasons, using either arithmetic means, or geometric means in cases where the data were highly variable (see contributions in Doubleday and Rivard 1981 and Pauly 1984).

Area 7 is that part of Module II where the grid of fishing stations was most dense, compared to the other areas. Here, unit catches were directly converted into density estimates. Also, in all cases where detailed recordings were available, the actual distance which the gear travelled over ground and the actual horizontal net opening (estimated from the angle and distance between the warp lines) were used for the computation of the area swept during individual hauls.

In the other areas, and/or whenever the area swept by individual hauls could not be estimated, the mean values presented above (area covered, net opening, etc.) were used. Either approaches yielded estimates of *relative* density, which may not be mistaken for estimates of *absolute* stock density, i.e., biomass- km^{-2} . However, one can derive approximate values of absolute stock density by multiplying relative density with the assumed catchability coefficient of the gear. This coefficient was assumed to be 0.5, a value that was subsequently verified for the similar species assemblage occurring in the Gulf of Thailand, and nets resembling ours (Pauly 1980).



Results and Discussion

Fig. 5 summarizes some of the temperature measurements taken during the survey. The average surface temperatures of the area was established to be 29°C, confirming Wyrcki (1961). The highest temperature measured was 31°C and the lowest was 28°C; 95% of the temperature recordings deviated from the mean value of 29.5°C by only $\pm 0.5^\circ\text{C}$. Seasonal fluctuations of surface temperatures were hardly noticeable.

The depth of the mixed layer tended to increase from north to south, again confirming Wyrcki (1961), and varied seasonally, especially in the Sunda Strait, where during the northwest monsoon, the thermocline was found to be 60 m higher than during the southeast monsoon, where it occurred at a depth of 120 m.

Two transects were run on August 1979 southward from the Java coast, with BT's shot every 20 nm. The resulting temperature profiles (of which one is shown on Fig. 5) show isotherms inclined toward the coast, suggesting a balanced geostrophic current.

No evidence of upwellings was detected outside of the Bali Strait area, where this phenomenon is well-documented (Nontji and Ilahude 1975). This disproves earlier suggestions of upwellings along the coast of southern Java (Wyrcki 1962; Cushing 1971), but also points out that our BT-based approach - which did not include nutrient measurements - was insufficient for serious oceanographic research. The following paragraphs, dealing with primary and secondary production, were thus assembled from the published literature.

Direct measurements of primary productivity in the survey area are rare. Productivity, integrated over the entire euphotic zone, was estimated as $0.43 \text{ g}\cdot\text{Cm}^{-2}\cdot\text{day}^{-1}$ in the area south of Java and $3.7 \text{ g}\cdot\text{Cm}^{-2}\cdot\text{day}^{-1}$ south of Bali (Stehmann-Nielsen and Jensen 1957). Krey and Babenerd (1979) present

productivity values of $0.5 \text{ g}\cdot\text{Cm}^{-2}\cdot\text{day}^{-1}$ for the area off eastern Java and around Bali during the southeast monsoon. During the northwest monsoon, there is generally a lower productivity, which hardly ever exceeds $0.25 \text{ g}\cdot\text{Cm}^{-2}\cdot\text{day}^{-1}$.

According to Nontji (1977) and Nontji and Ilahude (1975), the highest concentrations of phytoplankton in the area occur south of Bali ($2.36 \text{ mg}\cdot\text{m}^{-3}$) and off central Java ($1.4 \text{ mg}\cdot\text{m}^{-3}$), during the southeast monsoon period. Leyndekkers (1964) and Tranter (1962) report that zooplankton reach their highest biomass during the southeast monsoon. This happens when the south equatorial current is strongest, as are the upwelling along its northern flank and south of Java (Cushing 1971). Box 3 summarizes the incidental information gathered on zooplankton and micronekton.

Pelagic Fish Distribution

Fig. 3 presented earlier shows the distribution of stations sampled using the pelagic midwater trawl, while Fig. 4 shows the cruise tracks of the acoustic surveys.

The bulk of the pelagic trawl hauls yielded zero catches, while the acoustic survey did not yield absolute estimates of biomass due to unavailability of estimates of fish target strength.

However, estimates of relative abundance were obtained (Fig. 6). These are presented, as mm deflection per nautical mile (nm), and on a seasonal basis, since strong seasonal differences of relative abundance were observed (Table 1), probably reflecting real change in biomass and not only our different sampling grids (Fig. 6). Some information on squid is gathered in Box 4 (see also the database presented in Torres et al., this vol.).

Demersal Fish Distribution and Abundance

The two gears used to map the distribution of demersal fish were bottom trawl and echosounding; the latter was important in view of the fact that the "demersal" fish stocks in the area included many species of semipelagic fishes, spending much of their time clear off the sea bottom, and because much of the survey area was too rocky and/or uneven to be sampled by bottom trawling. Integrator deflections were assigned to the pelagic category if they occurred over 10 m away from the bottom; otherwise the echoes were added to those of the true bottom fish (Fig. 9).

No particular study was made of the sea bottom, but analyses of samples that were brought to the surface by the nets and the otterboards allowed some conclusions. Supplemented by recordings of the echosounder, these provided the following picture:

The sea bottom off the southwestern coast of Sumatra is sandy to muddy whereas the shelf around the islands of Nias, Mega and Engano is of a rough and rocky nature. The bottom is interspersed with coral reef formations in the shallow waters around the islands of Tanah Bala and Pulau Pini. Adjacent to the south, the Sunda Strait is characterized in its central parts by rough banks, parts of the exploded Krakatau volcano. The

Box 3. Zooplankton and micronekton.

[Boks 3. Zooplankton dan micronekton.]

During the activities of Module II of JETINDOFISH, not enough resources were available to do any research on the zooplankton and microplankton, although this would have provided some knowledge of the distribution of the early life history stages of fishes and/or of their food. The following observations are based on the fact that some of the larger zooplankton stuck to the meshes of the trawls (especially so in the case of the midwater trawls):

More than 70 different taxa of macroplankton were recorded, mainly fish larvae and young fishes, but also squids and crustaceans (see Boxes 4 and 5). At various stations, scattering layers were observed at night, both at depths between 120 and 150 m and close to the surface. Some of the larger components of these layers retained by our midwater trawl were typical deepwater species, such as *Chauliodus*, juvenile and adult myctophids, Gonostomatidae and postlarval and juvenile fishes of various other families, notably acanthurids, gempylids and priacanthids. Several times, huge amounts of leptocephali, squids, salps and *Pyrosoma* were caught. Particularly high number of leptocephali, over 20 liters per haul, were found around the Mentawai Islands, off southeast Java, and in the Bali Strait. Details may be found in the database described in Torres et al. (this vol.).




Fig. 6. Relative abundance of pelagic fish during the rainy and dry seasons from August 1979 to July 1981, as indicated by echointegration. Note that there is a higher relative abundance of pelagic fish during the rainy season than during the dry season.

[Gambar 6. Kelimpahan relatif ikan-ikan pelagis selama musim hujan dan kering dalam kurun waktu Agustus 1979 hingga dengan Juli 1981, sebagaimana kelimpahan relatif ini muncul lebih tinggi dalam musim hujan daripada musim kering.]

Table 1. Areas with low, medium and high integrator deflection, indicating relative abundance of pelagic fish during the rainy (November-March) and dry seasons (May-September).

[Tabel 1. Perbandingan rata-rata penyimpangan integrator, yang menunjukkan kelimpahan relatif ikan pelagis (dalam mm per nm) selama musim hujan (November-Maret) dan musim kering (Mei-September).]

Deflection (mm/nm)	Sumatra		Java	
	Rainy(Season) nm ² .10 ³	Dry nm ² .10 ³	Rainy(Season) nm ² .10 ³	Dry nm ² .10 ³
0	76.0	25.4	27.6	41.0
1	8.6	0	10.8	1.1
2	0.2	0	2.2	0
Relative abundance ^a	0.106	0.000	0.374	0.026

^aObtained from area covered x mm deflection total area surveyed in that season.

Box 4. Squids caught during the JETINDOFISH survey.

[Boks 4. Cumi-cumi yang tertangkap dalam survei JETINDOFISH.]

After fish, squids were the most important marine organisms caught during the survey. The distribution of squids appeared rather homogeneous, as they were taken by bottom and by midwater trawls in every depth layer, in the entire survey area. At night, the midwater trawl often caught squids at depths of less than 150 m but these were mostly small specimens, i.e., planktonic juveniles. Squid eggs nearly ready to hatch were taken at 7 bottom trawl stations. As the mesh size used was such that many squids undoubtedly escaped through the net, these results could not be quantified.

Overall, squids were captured at more than 280 of all stations (45% of the total). The highest catch rates were obtained off Central Java, where between 30 and 129 kg-hour⁻¹ were caught by bottom trawls at stations of about 120 m depth. Catches of 95 kg squid-hour⁻¹ were obtained on the Sumatra Shelf, at 70 m and of 76 kg-hour⁻¹ in the southern part of the Bali Strait, near Blambangan Peninsula, at a depth of 60 m. Squids were caught in 49 midwater trawl hauls; in five of these hauls, the gear was towed through a scattering layer which contained small pelagic squids (see Box 3).

Table 2. Summary of information on the trawlable vs nontrawlable bottoms in the survey area, as estimated from trawling experiments and acoustics (all areas in km².10³).

[Tabel 2. Ringkasan informasi keadaan daerah survei antara yang dapat ditrawl dan yang tidak, berdasarkan estimasi dari percobaan penggunaan trawl dan pengamatan akustik (semua daerah dalam unit km² 10³).]

Subarea	Total shelf	Surveyed area	Shallows to 99 m		100-200 m	
			Trawlable	Untrawlable	Trawlable	Untrawlable
3	40	20.5	10.0	2	2.5	6
4	18	13.0	5.0	1	2.0	5
5	1	13.5	4.5	2	1.0	6
6	13	12.0	4.5	1	1.5	5
7	12	11.0	4.0	1	2.0	4
Σ	84	70.0	28.0	7	9.0	26

Table 3. Average total catch per effort (\bar{x}) of bottom trawl, by area, depth and season (in kg-hour⁻¹).

[Tabel 3. Ringkasan hasil tangkapan total per upaya dari trawl dasar (kg per jam) berdasarkan daerah, kedalaman dan musim.]

Sub-area	Depth (m)	NW monsoon (November-February)			Intermonsoon I (March-April)			SE monsoon (May-September)			Intermonsoon II (October)		
		n	\bar{x}	s.d.	n	\bar{x}	s.d.	n	\bar{x}	s.d.	n	\bar{x}	s.d.
3	0-49	1	220	-	18	315	159	7	434	543	-	-	-
	50-99	7	171	57	28	164	122	8	175	140	-	-	-
	100-149	-	-	-	3	71	40	-	-	-	-	-	-
	150-199	-	-	-	-	-	-	1	241	-	-	-	-
	200+	-	-	-	3	74	133	2	506	345	-	-	-
4	0-49	-	-	-	7	658	484	6	246	151	-	-	-
	50-99	-	-	-	5	109	81	6	179	140	-	-	-
	100-149	-	-	-	-	-	-	1	31	-	-	-	-
	150-199	3	71	13	-	-	-	3	54	41	-	-	-
	200+	3	127	110	2	100	111	2	241	151	-	-	-
5	0-49	9	203	218	-	-	-	10	326	326	5	195	128
	50-99	3	463	96	-	-	-	9	180	221	2	96	56
	100-149	1	608	-	-	-	-	4	94	104	-	-	-
	150-199	2	441	283	-	-	-	2	233	137	-	-	-
	200+	1	759	-	-	-	-	2	64	66	-	-	-
6	0-49	7	1,161	1,346	-	-	-	17	275	155	-	-	-
	50-99	11	301	318	-	-	-	27	307	392	-	-	-
	100-149	1	394	-	-	-	-	23	145	117	-	-	-
	150-199	5	204	129	-	-	-	-	-	-	1	83	-
	200+	1	155	-	-	-	-	-	-	-	-	-	-
7	0-49	4	625	796	12	311	368	6	1,208	1,369	-	-	-
	50-99	10	18	735	397	475	309	11	509	386	-	-	-
	100-149	7	1,496	1,289	16	265	281	-	-	-	-	-	-
	150-199	6	397	207	11	270	197	2	316	247	-	-	-
	200+	11	360	242	19	281	140	1	228	-	-	-	-
Σ	-	77	-	-	158	-	-	150	-	-	16	-	-




Fig. 7. Showing the distribution of smooth (trawlable) and rough (untrawlable) grounds, as assessed from echosounding records and bottom trawling dur
[Gambar 7. Penyebaran dasar perairan yang layak dan tak layak untuk operasi trawl berdasarkan pengamatan echosounding dan operasi trawl dasar se




Fig. 8. Bottom trawl stations covered in Module II of JETINDOFISH, from August 1979 to July 1981.
 [Gambar 8. Penyebaran stasiun trawl dasar yang diliput selama JETINDOFISH Modul II, dari Agustus 1979 hingga Juli 1981.]




Fig. 9. Relationship between demersal trawl abundances estimates through bottom trawling (in kg·hour⁻¹) and acoustics (mm deflection of integrator). Based on records from locations where both measures could be unequivocally related to each other.

[Gambar 9. Hubungan antara penunjuk kelimpahan sumberdaya ikan demersal (kg/jam) berdasarkan pengamatan trawl dasar dan pengamatan akustik (mm defleksi dari integrator), sesuai dengan catatan pengamatan dari stasiun-stasiun dimana kedua parameter sangat berkaitan satu sama lain.]

coastal areas range from sandy to sandy-muddy. The western shelf of the Java Island is very narrow, rocky, and scarred by ravines. These conditions make trawl fisheries almost impossible.

The sea bottom to the south of Central Java is an extended shelf with a substratum that is predominantly muddy. The shelf in front of the East Java coast is generally about 4 nm wide, and the bottom usually rocky; wherever the sea bottom is level, it is almost always muddy-sandy. The adjacent Bali Strait has a predominantly shallow, sandy bottom; whereas

Box 5. Lobsters and deep sea shrimps.

[Boks 5. Udang barong dan udang laut dalam.]

Bottom trawling is not a good method for aimed fishing of palinurid lobsters, which generally hide in the crevices of rocky areas during daytime, from which they emerge to forage at night.

Still, adult spiny lobsters occurred in 6% of all successful trawl catches, notably 26 individuals from a depth of 211 m off Central Java, and 12.5 kg·hour⁻¹ at a depth of 320 m in the Bali Strait.

The depth preference of spiny lobsters is illustrated by the following:

Trawling depth (m)	Number of valid hauls	Hauls with adult palinurid (%)
0-49	340	7 (2)
100-199	97	8 (8)
200+	78	16 (21)

Phyllosoma larvae were occasionally observed clinging to the trawl net and the locations where this occurred generally corresponded to those areas where adults were caught.

Shovelnose lobsters (*Thenus* sp.) were caught in 35 bottom trawl hauls; only at three of these stations were they taken together with spiny lobsters.

In contrast to spiny lobster, most shovelnose lobsters were caught in shallow waters, i.e., in nearly 50% of all stations down to 49 m, 20% of those between 50 and 99 m, and in less than 5% of the deeper stations.

Unidentified deep sea shrimps (*Solenocera*?) were caught in 160 of the valid bottom trawl hauls, with the highest concentration (28 to 108 kg·hour⁻¹) occurring along the southern coast of Central Java, at depths of 180-290 m.

south of Bali, the shelf is very rocky and with a rugged structure. The continental slope is generally very steep with ravines perpendicular to the coastline. Fishing with a bottom trawl is possible only in few selected places. Fig. 7 and Table 2 summarize these findings.

Fig. 8 shows the distribution of bottom trawl hauls by FRV

Jurong during the survey area, and Table 3 presents the corresponding mean catch rates, by area, depth and season.

Fig. 9 shows the relationship between trawl catch rates and the corresponding acoustic estimates of density. The (geometric mean) regression line (Ricker 1973) shows a reasonably good fit ($r = 0.70$; $P < 0.01$), suggesting that the acoustic estimates do reflect relative densities, and thus allow conversion of acoustic density estimates into (rough) estimates of catch/hour and *vice versa*.

The biomasses estimated for the entire shelf area covered by Module II were 140,000 t for area 3; 43,000 t for area 4; 62,000 t for area 5; 69,000 t for area 6 and 97,000 t for area 7 (areas 1 and 2 were not surveyed).

The final survey report submitted by the project team to the DGF (Lohmeyer 1982) presented, beyond those recalled here, detailed tables with catch/hour data for various commercial categories of fish, and estimates of potential yield (P_y) based on the Gulland equation of 1971, i.e., $P_y = 0.5 \cdot M \cdot B_o$, where with our biomass estimates serving as unexploited stock sizes (B_o) and "M" set at 1 year⁻¹.

This material is not presented here: we now know that the equation of Gulland (1971) does not predict *sustainable* yields (Beddington and Cooke 1983), and this also invalidates our initial estimates of potential yields for various groups of commercial fish.

I refer instead to McManus (this vol.) who presents community analyses based on the data we sampled, and to Pauly et al. (this vol.), who discuss the biology, distribution and abundance of various fish groups throughout Western Indonesia, based on data collected during *FRV Jurong's* and other trawl surveys (and see Box 5 for notes on large crustaceans).

Finally, I wish to draw the reader's attention to the contribution of Torres et al. (this vol.), which documents the (MS-DOS) computer files containing most of the data we gathered, including the trawl catches, haul-by-haul and species-by-species. These data are available to anyone interested in their further analysis.

Such (re)analysis would be to the credit of all those who participated in Module II of JETINDOFISH.

Acknowledgements

I would like to thank Drs. Jürgen Saeger and Daniel Pauly for extracting the above summary from my earlier accounts of Module II of JETINDOFISH (Lohmeyer 1982, 1987); also if belatedly *Terima kasih* and *Danke schön* to all the Indonesian and German colleagues who contributed to this project.

References

- Beddington, J.R. and J.G. Cooke. 1983. The potential yield of fish stocks. FAO Fish. Tech. Pap. 242, 47 p.
- Cushing, D.H. 1971. Survey resources in the Indian Ocean and Indonesian area. IOFC/DEV/71/2, 123 p. Indian Ocean Program, Food and Agriculture Organization, Rome.
- Doubleday, W.G. and D. Rivard, Editors. 1981. Bottom trawl surveys. Can. Spec. Publ. Fish. Aquat. Sci. 58, 273 p.
- Fischer, W. and P.J.P. Whitehead, Editors. 1974. Species identification sheets for fisheries purposes. Eastern Indian Ocean and Western Central Pacific. Vol. 4. Food and Agriculture Organization, Rome.
- Gloerfelt-Tarp, T. and P.J. Kailola. 1984. Trawled fishes of Southern Indonesia and Northwestern Australia. Directorate General of Fisheries, Indonesia and German Agency for Technical Cooperation, Federal Republic of Germany.
- Gulland, J.A. 1971. The fish resources of the ocean. Fishing News Books for FAO, West Byfleet, Surrey, England. 255 p.
- Krey, J. and B. Babenerd. 1979. Atlas of the international Indian Ocean expedition. Intergovernmental Oceanographic Commission, UNESCO, Paris.
- Leyndekkers, J.V. 1964. The Indian Ocean expedition. Aust. J. Sci. 27(6):153-161.
- Lohmeyer, U.P., Editor. 1982. JETINDOFISH - joint eastern tropical Indian Ocean survey. Survey report module II. Directorate General of Fisheries, Jakarta, Indonesia, and GTZ, Eschborn, Federal Republic of Germany. (Unpublished report, coauthored by Dr. F. Woerner and Messrs. B. Ziegler, T. Gloerfelt-Tarp and Bambang Kartiko). 277 p.
- Lohmeyer, U.P. 1987. Bestandkundliche Untersuchungen der Bodenfische des östlichen Indischen Ozeans in vorgelagerten Schelf der Inseln Sumatra, Jawa und Bali. Universität Kiel, Kiel Germany. 138 p. Doctoral thesis.
- Losse, G. and A. Dwippongo. 1977. Report on the Java Sea southeast monsoon trawl survey, June-December 1976. Marine Fisheries Research Reports/Contribution of the Demersal Fisheries Project No. 3:1-119.
- Masuda, H., C. Araga and T. Yoshino. 1980. Coastal fishes of Southern Japan. Revised edition. Tokai University Press, Tokyo.
- Munro, I.S.R. 1967. The fishes of New Guinea. Department of Agriculture, Stock and Fisheries, Port Moresby, New Guinea. 651 p.
- Nontji, A. 1977. Notes on the chlorophyll distribution around Java. Oseanol. Indon. 7:43-47.
- Nontji, A. and A.G. Ilahude. 1975. Ekologi fitoplankton di Selat Bali. Oseanol. Indon. 5:25-42.
- Pauly, D. 1980. A new methodology for rapidly acquiring basic information on tropical fish stocks: growth, mortality and stock-recruitment relationships, p. 154-172. In S. Salla and P. Roedel (eds.) Stock Assessment for Tropical Small-scale Fisheries Workshop, 17-21 September 1979, University of Rhode Island. International Center for Marine Resource Development, Kingston, Rhode Island. 198 p.
- Pauly, D. 1983. Some simple methods for the assessment of tropical fish stocks. FAO Fish. Tech. Pap. (243):33-34.
- Pauly, D. 1984. Methods for assessing the marine stocks of Burma with emphasis on demersal species. FAO FI/Dp/BUR/77/003.
- Ricker, W.E. 1973. Linear regression in fisheries research. J. Fish. Res. Board Can. 30: 409-434.
- Stehmann-Nielsen, E. and E.A. Jensen. 1957. Primary oceanic production. Galathea Rep. 1:49-125.
- Tranter, D.J. 1962. Zooplankton abundance in Australian waters. Aust. J. Mar. Freshwat. Res. 13, 106 p.
- Venema, S.C. 1976. Report of the operations and results of the UNDP/FAO vessel *Lemuru* in Indonesian waters. A full account of all fishing operations, echo-recordings, observations, cruise tracks and oceanographic data. INS/72/064. 10 parts. pag. var. Development and Management Project. (Mimeo).
- Weber, M. and L.F. de Beaufort. 1913. The fishes of the Indo-Australian Archipelago. Vol. 2. E.J. Brill, Leiden. 404 p.
- Weber, M. and L.F. de Beaufort. 1916. The fishes of the Indo-Australian Archipelago. Vol. 3. E.J. Brill, Leiden. 455 p.
- Weber, M. and L.F. de Beaufort. 1922. The fishes of the Indo-Australian Archipelago. Vol. 4. E.J. Brill, Leiden. 410 p.
- Weber, M. and L.F. de Beaufort. 1929. The fishes of the Indo-Australian Archipelago. Vol. 5. E.J. Brill, Leiden. 458 p.
- Weber, M. and L.F. de Beaufort. 1931. The fishes of the Indo-Australian Archipelago. Vol. 6. E.J. Brill, Leiden. 448 p.
- Weber, M. and L.F. de Beaufort. 1936. The fishes of the Indo-Australian Archipelago. Vol. 7. E.J. Brill, Leiden. 607 p.
- Wyrtki, K. 1961. Naga report: scientific results of marine investigations of the South China Sea and the Gulf of Thailand 1959-1961. Vol. 2. Scripps Institution of Oceanography, La Jolla, California.
- Wyrtki, K. 1962. The upwelling in the region between Java and Australia during the south-east monsoon. Aust. J. Mar. Freshwat. Res. 13:217-225.

Marine Bottomfish Communities from the Indian Ocean Coast of Bali to Mid-Sumatra^a

JOHN W. McMANUS
International Center for Living
Aquatic Resources Management
MCPO Box 2631, 0718 Makati City
Philippines

McMANUS, J.W. 1996. Marine bottomfish communities from the Indian Ocean coast of Bali to mid-Sumatra [*Komunitas ikan dasar di perairan pantai Samudra Hindia dari Bali hingga pertengahan Sumatra*], p. 91-101. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

The fish communities of Bali to mid-Sumatra region are characterized by an assemblage extending reasonably uniformly across all depths overlain by a shallow fish community of more mobile species which tapers off sharply at approximately 100 m depth. The physical explanation for this limitation is not clear, but behavioral adaptations to periodic oxygen limitations cannot be discounted. Both the shallow and depth-ubiquitous communities can be subdivided further, but the resulting assemblages exhibit high between-group similarities. The shallow assemblages are both species-rich and highly even in distribution of biomass among species. The ubiquitous assemblages are relatively species-poor and exhibit low evenness. The waters below 100 m could probably be fished by commercial interests without substantially affecting the fish communities in shallow waters and vice-versa, but the low biomass of fish below 100 m probably makes this impractical.

Abstrak

Komunitas ikan dari Bali hingga daerah pertengahan Sumatra ditandai dengan suatu asosiasi kelompok yang tersebar secara seragam di semua kedalaman dan ditandai dengan komunitas ikan daerah dangkal yang merupakan spesies yang aktif bergerak dimana secara berangsur-angsur berkurang kelimpahannya secara tajam pada kedalaman sekitar 100 m. Keterbatasan secara fisik tidak jelas, namun perilaku adaptasi terhadap keterbatasan oksigen tidak boleh diabaikan. Baik komunitas laut dangkal maupun komunitas yang tersebar merata berdasarkan kedalaman dapat diperinci lagi lebih lanjut, namun demikian kelompok pembagian yang dihasilkan menunjukkan persamaan antar grup yang tajam. Kelompok laut dangkal pada umumnya kaya akan spesies dan mempunyai penyebaran biomassa yang merata antar spesies. Kelompok yang tersebar merata pada semua kedalaman umumnya miskin akan spesies dan penyebarannya kurang merata. Kegiatan penangkapan pada kedalaman lebih dari 100 m dapat dilaksanakan tanpa mempengaruhi perairan wilayah dangkal, demikian pula sebaliknya, namun rendahnya biomassa sumberdaya perikanan pada perairan kedalaman dibawah 100 m, menyebabkan kegiatan penangkapan ikan menjadi tidak praktis dilakukan.

Introduction

Trawl fishing grounds are often heterogeneous environments consisting of a variety of identifiable habitats and their associated species assemblages. An important step toward informed management of such grounds is the identification of these species-habitat combinations. Species which tend to co-occur, the "recurrent groups" of Fager and Longhurst (1968), can serve as fundamental fishery management units, or Assemblage Production Units (APU - Tyler et al. 1982). Indeed, models have shown that efforts to optimally harvest a diverse assemblage of species with different production capabilities as a single, combined unit will tend to lead to a reduction in the number of species contributing to production. Such an imposed shift in the relative abundances among species may not nec-

essarily be reversible upon reduction of fishing effort (Tyler et al. 1982; Beddington 1984). Ralston and Polovina (1982) showed that production models analyzed for a heterogeneous demersal fishery as a whole may be less informative and useful than similar models applied individually to groups of species identified by cluster analysis. Finally, it has been suggested that site-species groupings may be used as a basis for dividing a fishery between competing fishing interests, such as between large-scale commercial trawling and small-scale fishing (McManus 1985a). In this case, the boundaries between species distributions can be treated as "ultimate stock boundaries", and optimally chosen lines dividing areas containing different sets of species can potentially be used to regulate fisheries on a geographic basis.

Southeast Asian waters form a biogeographic unit of high diversity (the highest marine diversity globally), with strong

affinities to communities throughout the Indian Ocean (McManus 1985b, 1993). Previous classificatory studies of demersal fish communities of these two regions include McManus (1985a, 1986, 1989), Bianchi (1992), Bianchi et al. (this vol.) and Federizon (1992). Other studies are summarized in Longhurst and Pauly (1987). The study of McManus (1985a, 1986) involved analyses of published data from 15 trawl surveys made at the same 28 sites in the Samar Sea (Philippines) over 1.5 years. This division involved principally species within genera, and only some families (e.g., Nemipteridae) were restricted by the line. The study demonstrated that a major feature of the community, a division at approximately 40 m depth, was a stable feature throughout the period and hence not affected by seasonality. This helped to substantiate the usefulness of community structural analysis methods originally designed for stationary land plant communities in analyses of communities of highly mobile fish. However, the sites sampled were primarily shallower than 100 m. A preliminary study of the data from the JETINDOFISH survey from the Indian Ocean side of Indonesia (McManus 1989) showed that a far more substantial division existed at approximately 100 m, involving several tens of families and hundreds of species which did not extend deeper than this. Subsequently, Federizon (1992) demonstrated that a 100-m division occurred in the Ragay Gulf of the Philippines as well, a preliminary indication that the division may be a widespread phenomenon. Bianchi's (1992) studies of research trawl data from the western Indian Ocean (Pakistan, Oman, Yemen, Ethiopia and Somalia) indicated a tendency for many fish species to shift location in order to avoid seasonal oxygen depletions. This was particularly apparent in areas of seasonal upwelling, and depth-based community structure could be expected to be far more stable in nonupwelling areas, such as the eastern Indian Ocean. Bianchi (this vol.) analyzed research trawl data from the northwest Sumatra, and found a community break at 40 m quite similar to that found in the Samar Sea (McManus 1985a, 1986). That study again included few samples below 100 m.

The current study is a more complete analysis of the JETINDOFISH data aimed at elucidating the trends identified in the preliminary analysis (McManus 1989). In particular, the study emphasizes the grouping of sites and species by co-occurrence so as to facilitate the management of this extensive fishery, as well as the future investigation of the specific ecological and evolutionary causes of the depth limitation.

Methods

The study area (Fig. 1) was described by Lohmeyer (this vol.), who provided a de-

scription of the trawls, tracks and other useful information; the material discussed here was gathered through over 860 research trawl hauls made from 1979 to 1981. For this study, hauls which were more than 10 m above bottom at their start or finish were omitted, as were hauls for which no fish families were identified. This left 534 trawls for analysis. This being an awkwardly large number of objects to display and interpret effectively in a classification analysis, it was decided to average the species abundances found within each area and each 10 m depth zone. This left 119 area-depth groupings or "sites" (Table 1). The data on numbers of fish were incomplete or inaccurate in many cases (e.g., 20 kg, 0 individuals), reflecting the focus of the research program on locating large biomasses of trawlable fish rather than on determining community structural properties. Bianchi (1992) showed that when dealing with large-scale gradients and heterogeneous assemblages, results from fish weight vs. fish number analyses differ only slightly. Thus, the present analyses concentrated on analyses of catch weights standardized to kg km^{-1} .

The JETINDOFISH Project was extensive in scope, involving several taxonomists identifying species from a poorly known region. Data were copied from form to form and key-punched into mainframe computers, and later re-entered into a microcomputer manually after copying from printout summaries (after archived tapes became potentially unreadable). The resulting list of 9,723 species catch records contained 1,286 "species", including some identified only by a code and others known to family, genus or species level. The latter species names involved some misspellings. However, most of the names needed taxonomic updating. This was done for finfish using an automatic synonymy program (see Froese




Fig. 1. Map showing the trawl areas for the JETINDOFISH survey. Areas 1 and 2 were not surveyed. [Gambar 1. Peta yang menunjukkan daerah trawl untuk survei JETINDOFISH. Tidak ada survei yang dilakukan di daerah 1 dan 2.]

Table 1. Sample distribution chart. The numbers represent the number of hauls averaged within each area - depth class to give the average catch rates per species for each "site". Blank boxes are those for which no data were available.

[Tabel 1. Peta distribusi contoh. Angka menunjukkan jumlah tarikan rata-rata di tiap daerah dan kelompok kedalaman untuk memperoleh rata-rata hasil tangkapan per jam berdasarkan spesies di setiap daerah. Kotak yang tidak ada angkanya menunjukkan ketidaktersediaan data.]

Depth (m)	Area							
	A3	A4	A5	A6	A7	A8	A9	
10	-	3	-	-	1	2	-	
20	6	1	3	3	2	13	1	
30	8	7	10	10	5	16	7	
40	9	7	8	10	15	32	8	
50	4	4	4	17	12	26	1	
60	10	5	5	6	8	19	2	
70	8	5	3	5	14	9	1	
80	6	9	2	5	6	5	-	
90	2	-	2	5	2	-	-	
100	1	-	-	13	1	1	-	
110	-	1	1	10	2	-	-	
120	1	-	1	2	3	-	-	
130	-	-	1	-	1	2	-	
140	-	-	1	-	3	4	-	
150	-	-	-	1	3	1	-	
160	-	1	-	-	1	-	-	
170	-	1	2	2	2	-	-	
180	-	2	-	3	4	-	-	
190	1	3	2	2	2	-	-	
200	-	2	-	1	5	-	-	
210	-	-	2	-	2	-	-	
220	-	2	-	-	2	-	-	
230	1	1	-	-	4	1	-	
240	-	1	-	2	2	1	-	
250	-	-	-	-	4	-	-	
260	1	-	-	1	2	-	-	
270	-	-	1	-	3	-	-	
280	-	1	-	1	5	-	-	
290	-	-	-	1	1	-	-	
300	-	1	-	-	-	-	-	
320	-	-	-	1	-	-	-	
330	1	-	-	-	-	-	-	

et al., this vol.) which matched up the survey names with the list of over 60,000 synonymies in FishBase 96 and suggested corrections. The resulting corrected list of names contained many within-survey synonymies, reducing the total species list in the survey (including species not fully identified, as well as harvested cephalopods and crustaceans) to 703 species. The availability of this procedure greatly enhanced the accuracy of this study, and may be considered a significant step toward enabling more thorough comparative analyses of research trawl data from around the world in the future (see also Pauly, this vol.).

The data were classified into species-site groups using two-way indicator species analysis (TWIA - see Box 1 and Bianchi, this vol.). For this purpose, a microcomputer version of the original TWINSpan program (Hill 1979) was recompiled to be able to handle 400 species (with 119 sites), the maximum possible on the existing microcomputer system available. Thus, the data matrix was converted from the Microsoft Access database format to Microsoft Excel spreadsheet format, and reduced to the 400 most abundant species by weight before conversion (using a new macro in Visual Basic for Microsoft Excel) to the compressed data format required by TWINSpan.

Box 1. Two-way indicator species analysis (TWIA).

[Boks 1. Analisis spesies berdasarkan indikator dua arah (TWIA).]

TWIA is a method for rearranging tables of species abundances by site and dividing the table into distinct sets of similar species (based on where they are found in what abundance) and similar sites (based on what species are found in them in what abundance). A successful classification yields data arranged in clear blocks within the table. In the early days of plant community ecology ("phytosociology"), such tables were created by tediously rearranging raw data tables by hand, with repetitive cutting and pasting of columns and rows. The TWIA approach, embodied in the TWINSpan program of M.O. Hill (1979) revolutionized the field by automating the procedure.

TWIA begins by turning a table of abundances into a table of presence-absence values. This is accomplished by requiring an initial coding of all data into single digit integer values, such as a 1-5 semiquantitative scale. A species x with a maximum value of 3 in a row, becomes species x_1 , x_2 and x_3 , each with a maximum value of 1 in any column. The program then arranges the sites (columns) of a table based on the scores of each site on the first axis of a reciprocal averaging (basically correspondence analysis) ordination axis. This procedure groups similar sites together and lines them up along the axis representing the greatest variance among both sites and species taken simultaneously. The program then attempts to divide the table into left and right portions from the center of the columns. It analyzes the effect of this split in terms of how well the abundances of each species are clumped onto one side or the other of the division. It tries several such divisions, cutting left and right of the center, each time tallying up a score based on how well the species abundances are divided up. It then chooses the best division, and this becomes the highest level division of the classification. The program now repeats the ordination for only those sites (columns) on the left of the division, to find a good arrangement and division of this subset of columns. Then it does the same on the right of the first division. Now there are four classes of sites, as indicated by two levels of division. A third level of division would yield eight classes of sites, and so on. This is the basic procedure, although the program contains corrections and adjustments for rare species weightings, etc., which can be manipulated in the analysis.

Next, the program classifies the species (rows) in a manner similar to that of the sites (columns). Once the species classification has been completed, the program switches around the species (keeping the species groups intact) in such a way that the table is in an optimal "block" form, with blocks of high species abundances arranged along one or the other diagonal of the table. The table is then converted back to the integer values, and printed along with indications of the hierarchy of site and species divisions. Additionally, the program indicates what species are particularly good indicators of any site division, and helps to identify "border" species of uncertain position in a division of sites.

The method has the advantage over most agglomerative classifications that it is very easy in the end to explain why a particular set of sites or species were grouped together. This is generally obvious from studying the output table. As a divisive "top-down" approach, the method also avoids the problem of "chaining", i.e. the tendency for a single class in agglomerative "bottom-up" clustering to grow like a snowball, gathering new characteristics as it does, and thus obscuring the inherent structure of the data set. However, TWINSpan does not always provide clear one-to-one matching between site and species groups, e.g., there may be eight site groups but only six species groups due to failed divisions (based on various criteria for failure that the program uses). Even equal numbers of site and species groups may not match up well, such that the third group of sites may not correspond to the third group of species. This is because the method does not perform the two-way analyses simultaneously. Finally, there is no way to determine the probability that a given division could have been achieved by the mere fact that the matrix was sparse (containing lots of zero values) and patchy to begin with. Nonetheless, TWINSpan is the most widely used program of its type currently available.

The top 400 species represented 98% of the total weight of fish (after standardization to kg km^{-1}). The effort to include as many species as possible was necessary because many of the deeper water samples were dominated by species which were ranked very low in total catch weight in the overall study.

Box 2. Multidimensional scaling analysis (MDA) and other ordination methods.

[Boks 2. Analisis skala multidimensi (MDA) dan cara ordinasi yang lain.]

MDA is a method for ordinating multidimensional data, i.e., displaying the data (e.g., sites based on similarities among species abundances) on a few (often 1, 2 or 3) axes in a graph. The fundamental problem is that one cannot do so without distorting the data. For example, one cannot project the Earth onto a two-dimensional map without altering the appearances of the continents — this is why there are so many different kinds of map projections available. The most common ordination method, principal components analysis (PCA), works by rotating an axis into such a direction in the multidimensional cloud of points (e.g., sites), until the projection of those sites onto the axis explains the greatest possible variance (“spreads out” the points the best). This becomes the principal axis. The second axis is limited to being at right angles to the first, but rotated through the dimensions until it explains most of the residual variance. A third, fourth and fifth axis can be constructed accordingly, but this is difficult to see in one’s mind’s eye. The data cloud then projected onto those axes and plotted on a piece of paper then represents the most of the variance in the data cloud possible in a two-dimensional picture.

Correspondence analysis (CA) and the nearly identical reciprocal averaging (RA) work similarly to PCA. However, the final axes are not rotated to account simply for the variance of the sites or species, but rather to account for each relative to the other. The final graph can include both species and sites on the same plane, something which is not possible in simple PCA.

MDS differs from these methods in that the objective is more clearly to display the data in a given set of dimensions with as little distortion as possible. Thus, there is no rotation of axes. Rather, the data are shifted into place in an algorithm that minimizes the distortion in the distances between the points as it proceeds. Mathematically, this involves the creation of a distance matrix, and in this study the data were transformed to $\ln(x+1)$ to minimize effects of spurious abundances, and then converted to a Euclidean distance matrix for analysis. As with PCA, a major portion of the variance tends to be associated with the first axis (although not in such an “intentional manner”), and thus it is reasonable to plot the sites based on the first axis coordinates against environmental parameter values such as depth.

PCA is particularly problematic when dealing with nonlinearly related points, such as sites related by species abundances. Both PCA and MDS tend to form arches where one would expect straight lines reflecting known gradients. CA also arches, but less so, and avoids problems such as inversion of points at ends of a gradient. The arch can even be minimized through “detrending” (see Bianchi, this vol.). Thus CA or detrended CA (DCA) is preferred by many ecologists. Other ecologists prefer MDS because of its philosophical orientation (minimizing distortion). Still others prefer to use both, so as to check that results are robust, i.e., not an artifact of a particular method. In the end, the choice may be simply one depending on the availability of working, reliable software, as was the case in this study.

due to the relatively much higher biomasses encountered in the species-rich shallow waters. The data were analyzed using the default settings of the program except for the levels of hierarchical analysis, which were set to three. The default cut values (upper limits of 1, 2, 5, 10 and 20+ kg/km scaled to values 1 - 5, respectively) were used as they produced a reasonable distribution of values in the output table.

A graphical analysis of the relationships among sites was accomplished using multidimensional scaling (Norkis 1993, and see Box 2). The SPSS for Windows Version 5.0 software imposed a limit of 100 objects on the analysis. Thus, it was necessary to remove the least abundant 19 sites from the analysis. The top 100 sites used represented 98% of the total catch-weight. Data (x) were converted to $\ln(x+1)$ before MDS analysis to minimize the effects of spuriously high abundances from chance encounters with large schools. The sites were

identified as to groupings from TWIA (group H, consisting of a single site, was removed in the processing). The primary MDS axis was plotted against depth to highlight the relationships between the TWIA groupings and depth.

Finally, the mean kg km^{-1} values of each species by depth were tabulated for the top 50 species. These were converted to $\ln(x+1)$ values and then sorted by “center of gravity” (mean depth after weighting by abundance). The values were then plotted in stacked histograms representing the gradual shift in species structure as one proceeds from shallow to deeper waters.

Results

The TWIA output is shown in Table 2. Each column represents a “site” (average standardized haul within an area and a 10 m depth group - designated such that “20” means 10+ to 20 m). The area (3-9) is indicated at the top of each column. The site group identified by TWIA follows, as indicated by a dash or letter. The depth group is indicated as two or three digits arranged vertically (the first column is “140”). Each row of the table represents a species, either by name or code. Species identified by TWIA as indicators of a final site group (A-F) are marked accordingly (only lower hierarchical divisions are considered for simplicity). In the table, a dash indicates a zero value. The values 1-5 represent upper class limits of 1, 2, 5, 10 and 20+ kg km^{-1} , respectively. Only the top 100 species are displayed, although the analysis involved 400 species.

The table shows a very clear division, wherein groups E, F, G and H include most of the sites below 100 m. Groups A and H appear to contain sites at intermediate depths around 100 m. The MDS graph shows that most of the deep sites cluster tightly relative to the shallow sites, an indication of relatively little variability in the deeper assemblages (Fig. 2). The plot of the primary MDS axis against depth clearly shows the restriction of groups B and C to waters above 100 m (Fig. 3). Groups B and C differ only in relative dominances by fairly similar species, as do groups E and F. The hierarchical relationships are shown in Fig. 4, and support these relationships. The top ten species of each group are listed in Table 3. The TWIA table shows that the shallow sites included most of the species which dominated the deep sites. However, these species were reduced to lower significance in the shallow sites by the presence of large biomasses of the shallow preferential species. Thus, the community was apparently a combination of an ubiquitous assemblage of lizardfish (e.g. *Saurida undosquamis*), bigeyes (e.g., *Priacanthus macracanthus*) and others, overlain by a shallow-water restricted assemblage of leiognathids, lutjanids and many others. This is supported further by the histograms of individual species against depth (Fig. 5).

An inspection of the site groups indicates that Group B, dominated by *Dasyatis* sp., *Leiognathus equulus* and others, is a shallow-water assemblage characteristic of areas 8 and 9, and to a lesser extent, 3. Group C, dominated by *Sardinella*




Fig. 7. Some oceanographic features of the study area (after Wyrki 1961): A. Depth (in m) of the homogeneous layer. B. Depth (in m) of the core layer of the subtropical lower water. C. Oxygen profile (in ml.l⁻¹) from the Indian Ocean near Java. [Gambar 7. Berbagai sifat oseanografis dari daerah studi (menurut Wyrki 1961): A: Kedalaman (m) dari lapisan yang homogen. B: Kedalaman (m) dari lapisan pokok yang berasal dari perairan subtropik bagian bawah. C: Profil oksigen (ml.l⁻¹) dari perairan Samudra Hindia dekat Jawa.]

area experiences seasonal or otherwise periodic or sporadic oxygen depletions with depth, or that the fish are behaviorally adapted to avoiding water masses which have such depletions elsewhere or earlier in geological history, and that they use the water temperature boundary (or accompanying rapid increases in pressure) to delimit their activity accordingly.

Other possible explanations include changes in light levels, productivity and sediment composition. The light limitation would be expected to cause species with differential light requirements to drop away one by one more gradually than is seen. The depth extent of high primary and secondary productivity would be rapidly curtailed below the homogeneous zone, but one might expect the abundances of species to fall off rapidly while still finding these species below this layer. The sediments may indeed change in composition as one progresses from the shelf to the slope at approximately 100 m. However, this sediment change might be expected to affect the synodontids and others as well. Each of these physical explanations remains a possibility for future elucidation. However, at this point it seems likely that behavioral adaptations have sharpened the effect of whatever physical limitation is

most influential.

An immediate conclusion from this study would be that commercial trawl fishing below 100 m would not interfere with small-scale fishing in shallower waters, as it would not affect the vast number of species restricted to waters above this depth. While that may be true, such a restriction is unlikely to be practical. As shown in Fig. 8, there is very little biomass to catch below 100 m. This confirms earlier analyses for Southeast Asian waters (Pauly 1987), indicating the futility of planning future expansions into deeper-water fisheries.

The shallow-water assemblages appear to be geographically important rather than being related to depth. Species such as *Leiognathus equulus* and *L. bindus* are desirable target species, and the differences could be related to fishing pressure. Equally, there could be average sediment differences between the lesser Sunda Islands (areas 8 and 9) and the Java to Sumatra region, although the shelves appear to be very heterogeneous (Shepard et al. 1949). The high similarity in species composition between these site groups may be a result of this heterogeneity of bottom types.

Finally, it is noteworthy that the 40-m zonation found by




Fig. 8. Depth distribution of catch weights summarized for all sites, reexpressed as standing stock ($t \cdot km^{-1}$).

[Gambar 8. Sebaran kedalaman dari berat hasil tangkapan yang disarikan dari seluruh daerah pengamatan, dimana satuannya diseragamkan menjadi $t \cdot km^{-1}$.]

Bianchi (this vol.) in northwestern Sumatra, and by McManus (1985b, 1986) in Samar Sea was not apparent in the present analysis. This could be simply a problem of scale. The area considered in this study was very large, and covered a much greater range of habitat variability relative to the others mentioned. The multivariate methods used are based on the analysis of the total variance of the data matrix used. Thus, the more heterogeneous the data analyzed, the more that finer scale, more subtle patterns are obscured.

Acknowledgements

I would like to thank Dr. Rainer Froese for spending substantial amount of time and effort in helping correct the species names for this study. Ms. Luningning Malumay typed in

nearly 10,000 catch records to make this analysis possible. The German Agency for Technical Assistance (GTZ) funded both the original survey and the current analysis. Finally, I would like to thank Dr. Daniel Pauly for helpful discussions and for providing the impetus for this study.

References

- Beddington, J. 1984. The response of multispecies systems to perturbations, p. 191-207. *In* R.M. May (ed.) *Exploitation of marine communities*. Springer-Verlag, New York. 367 p.
- Bianchi, G. 1992. Demersal assemblages of tropical continental shelves. University of Bergen, Norway. 217 p. Ph.D. dissertation.
- Fager, E.W. and A.R. Longhurst. 1968. Recurrent group analysis of species assemblages of demersal fish in the Gulf of Guinea. *J. Fish. Res. Bd. Can.* 25: 1405-1421.
- Federizon, R.R. 1992. Description of the subareas of Ragay Gulf, Philippines, and their fish assemblages by exploratory data analysis. *Aust. J. Mar. Freshwat. Res.* 43:379-391.
- Hill, M.O. 1979. TWINSpan - a FORTRAN program for arranging multivariate data in an ordered two-way table by classification of individuals and attributes. Cornell University, Ithaca, New York.
- Longhurst, A.R. and D. Pauly. 1987. *Ecology of tropical oceans*. Academic Press, New York.
- McManus, J.W. 1985a. Descriptive community dynamics: background and an application to tropical fisheries management. Ph.D. thesis, University of Rhode Island, Kingston.
- McManus, J.W. 1985b. Marine speciation, tectonics, and sea-level changes in Southeast Asia. *Proc. Fifth Int. Coral Reef Congress* 4:133-138.
- McManus, J.W. 1986. Depth zonation in a demersal fishery in the Samar Sea, Philippines, p. 483-486. *In* J.L. Maclean, L.B. Dizon and L.V. Hosillos (eds.) *The First Asian Fisheries Forum*. Asian Fisheries Society, Manila, Philippines.
- McManus, J.W. 1989. Zonation among demersal fishes of Southeast Asia: the southwest shelf of Indonesia, p. 1011-1022. *In* *Proceedings of the Sixth Symposium on Coastal and Ocean Management/ASCE*, 11-14 July 1989, Charleston, South Carolina.
- McManus, J.W. 1993. The Spratly Islands: a marine park? *Ambio* 23(3):181-186.
- Norkis, M.J. 1993. SPSS for Windows professional statistics release 6.0. SPSS Inc., Chicago.
- Pauly D. 1987. Theory and practice of overfishing: a Southeast Asian perspective, p.146-163. *In* *Papers presented at the Symposium on the Exploitation and Management of Marine Fishery Resources in Southeast Asia*, 16-17 February 1987, Darwin, Australia. RAPA Rep. 1987/10 p.
- Ralston, S. and J.J. Polovina. 1982. A multispecies analysis of the commercial deep-sea handline fishery in Hawaii. *Fish. Bull.* 80(3):435-448.
- Shepard, F.P., K.D. Emery and H.R. Gould. 1949. Distribution of sediments on East Asiatic Continental shelf. *Allan Hancock Foundation Publ., Occas. Pap.* 9. University of Southern California Press, Los Angeles, California.
- Tyler, A.V., W.L. Gabriel and W.J. Overholtz. 1982. Adaptive management based on structure of fish assemblages of northern continental shelves, p. 149-156. *In* M.C. Mercer (ed.) *Multispecies approaches to fisheries management advice*. *Can. Spec. Publ. Fish. Aquat. Sci.* 59:169 p.
- Wyrski, K. 1961. *Physical oceanography of the Southeast Asian waters*. Naga Rep. Vol. 2. Scripps Institution of Oceanography, La Jolla, California. 195 p.

Results of Surveys for Pelagic Resources in Indonesian Waters with the *R/V Lemuru*, December 1972 to May 1976

SIEBREN C. VENEMA

FAO of the United Nations
Viale delle Terme di Caracalla
00100 Rome, Italy

VENEMA, S.C. 1996. Results of surveys for pelagic resources in Indonesian waters with the *R/V Lemuru*, December 1972 to May 1976 [*Hasil survei sumberdaya ikan pelagis di perairan Indonesia dengan kapal penelitian R/V Lemuru pada bulan Desember 1972 hingga Mei 1976*], p. 102-122. In D. Pauly and P. Martosubroto (eds.) Baseline studies of biodiversity: the fish resources of Western Indonesia. ICLARM Stud. Rev. 23, 312 p.

Abstract

Results are presented of 48 surveys executed from December 1972 to May 1976 with the purse seiner/trawler *R/V Lemuru* in the Java Sea, Sunda Strait, Bali Strait and coastal areas of the Indian Ocean off Java and the lesser Sunda Islands. Most of this large area was surveyed with a sonar in sweeping mode, looking for concentrations of pelagic fish which were fished by purse seine, often after light attraction. Echosounder records, occasional hauls with bottom and mid-water trawl and catches by trolling lines are also analyzed.

The largely qualitative results of these surveys are compared with the known migration patterns of some important resource species, and with fishing activities at the time of the surveys. The survey results are then related to the subsequent development of pelagic fisheries in the Bali Strait and the Java Sea.

Abstrak

Tulisan ini melaporkan hasil 48 kali survei yang dilaksanakan dengan kapal penelitian multi-guna (purse seine [pukat cincin] dan trawl) *Lemuru* di perairan Laut Jawa, Selat Sunda, Selat Bali dan perairan Samudra Hindia di selatan pulau Jawa dan kepulauan Sunda Kecil dalam bulan Desember 1972 hingga bulan Mei 1976. Sebagian besar dari daerah yang luas ini disurvei dengan alat bantu sonar guna mencari konsentrasi ikan yang bergerombol yang kemudian ditangkap dengan jaring purse seine (dan kerap kali dengan kombinasi lampu). Selain itu, ditampilkan pula hasil analisis deteksi echosounders, operasi penangkapan dengan jaring trawl (tengah dan dasar) dan hasil tangkapan pancing tonda

Hasil survei yang sangat kualitatif ini dibandingkan dengan pola migrasi beberapa spesies ikan penting yang sudah diketahui serta dengan aktivitas penangkapan ikan yang terlihat pada saat survei dilaksanakan. Selanjutnya hasil survei ini dikaitkan dengan perkembangan perikanan pelagis di Selat Bali dan di Laut Jawa.

Introduction

This paper is an abstract of the "Report on the operations and results of the UNDP/FAO vessel *Lemuru* in Indonesian waters, a full account of all fishing operations, echo recordings, observations, cruise tracks and oceanographic data", issued in a limited number of mimeographed copies by the Fisheries Development and Management Project (Venema 1976). It covers data collected from December 1972 to May 1976, during which time *R/V Lemuru* was deployed in eight areas, defined in Fig. 1 (see also Table 1).

In December 1972 *R/V Lemuru* (ex *Sagar Sandhani*) was transferred from Bangladesh, where a war had made its operation impossible, to Indonesia where it was first assigned to a UNDP/FAO Fisheries Development and Training Project

based in Tegal, Central Java, and devoted to exploratory fishing and training of crew, then, from April 1973, to the Jakarta-based UNDP/FAO Fisheries Development and Management Project for carrying out resource surveys, led by an FAO biologist (Jones 1976).

From December 1972 until the end of July 1974, *R/V Lemuru* operated from Bena, Bali, and covered the Bali Strait (Area 7 in Fig. 1), Nusa Tenggara (Area 8), the Indian Ocean south of Java (Area 6), parts of the Java Sea (Areas 1 and 2) and the Sunda Strait (Area 5).

Of this large area, the Bali Strait received most attention, because fishers based at Muncar, East Java, had complained about large fluctuations in the availability of the Bali sardinella (*Sardinella lemuru*)^a, locally called *lemuru*. Data collected during the International Indian Ocean Expedition suggested that an upwelling occurred in the Indian Ocean outside Bali Strait and it was therefore thought likely that the Bali sardinella would form part of a larger resource migrating between the Indian

^aThis was previously known as *Sardinella longiceps* (see Pauly et al., this vol.).




Fig. 1. Map of Western Indonesia, showing the areas surveyed by R/V Lemuru from December 1972 to May 1976. [Gambar 1. Peta Indonesia bagian barat yang menunjukkan daerah yang disurvei oleh kapal penelitian Lemuru, Desember 1972 hingga Mei 1976.]

Table 1. Overview of survey cruises of R/V Lemuru in Indonesia, 1972-1976 (total days at sea: 588; see Fig. 1 for definitions of areas covered).

[Tabel 1. Rangkuman pelayaran survei kapal penelitian Lemuru di Indonesia, 1972-1976 (total 588 hari pelayaran, lihat Gambar 1 untuk keterangan yang diliput).]

Cruise no.	Dates		Days at sea	Departure	Ports		Areas covered
	Start	Finish			Arrival		
7201	3/12	11/12	9	Tanjung Priok	Benoa		2,5,6,7
7202	15/12	20/12	6	Benoa	Benoa		7
7301	29/1	2/2	5	Benoa	Benoa		7
7302	19/2	24/2	6	Benoa	Benoa		7
7303	7/3	12/3	6	Benoa	Benoa		7,8
7304	23/5	4/6	13	Benoa	Benoa		1,7
7305	18/6	2/7	15	Benoa	Tanjung Priok		1,2,7
7306	22/7	31/7	10	Tanjung Priok	Benoa		2,5,6,7
7307	10/8	25/8	16	Benoa	Benoa		6,7
7308	19/9	26/9	8	Benoa	Benoa		7
7309	12/10	1/11	21	Benoa	Benoa		8
7310	8/11	14/11	7	Benoa	Benoa		7
7311	22/11	3/12	12	Benoa	Benoa		6
7312	14/12	21/12	8	Benoa	Benoa		7
7401a	6/1	12/1	7	Benoa	Cilacap		2,5,6
7401b	16/1	20/1	5	Cilacap	Singapore		
7402	10/2	20/2	11	Singapore	Benoa		2,5,6,7
7403	23/2	27/2	5	Benoa	Benoa		7
7404	6/3	1/4	27	Benoa	Benoa		1,7,8
7405	23/4	3/5	11	Benoa	Cilacap		1,6,7
7406	9/5	13/5	5	Cilacap	Tanjung Priok		2,5,6
7407	27/5	12/6	17	Tanjung Priok	Benoa		1,2,5,6,7
7408	22/6	25/6	4	Benoa	Benoa		7
7409	27/6	17/7	21	Benoa	Benoa		8
7410	28/7	2/8	6	Benoa	Cilacap		6,7
7411	6/8	16/8	11	Cilacap	Cilacap		2,5,6
7412	26/8	6/9	12	Cilacap	Tanjung Priok		1,2,6,7
7413	18/9	25/9	8	Tanjung Priok	Semarang		1
7414	30/9	9/10	10	Semarang	Semarang		1
7415	21/10	28/10	8	Semarang	Semarang		1
7416	31/10	10/11	11	Semarang	Semarang		1
7417	17/11	23/11	7	Semarang	Singapore		2
7418	24/12	26/12	3	Singapore	Tanjung Priok		2
7501	8/1	28/1	21	Tanjung Priok	Belawan		2,4
7502	10/2	28/2	19	Belawan	Tanjung Priok		2,3,4
7503	22/3	4/4	14	Tanjung Priok	Semarang		2
7504	17/4	3/5	17	Semarang	Semarang		2,5
7505	17/5	31/5	15	Semarang	Semarang		1
7506	11/6	28/6	18	Semarang	Benoa		1
7507	11/7	27/7	17	Benoa	Semarang		1,7
7508	6/8	22/8	17	Semarang	Semarang		2
7509	3/9	1/10	18	Semarang	Semarang		2,5
7510	10/10	23/10	15	Semarang	Semarang		1
7601a	9/1	13/1	5	Semarang	Semarang		1
7601b	26/1	8/2	14	Semarang	Semarang		
7602	18/2	3/3	15	Semarang	Semarang		2,5
7603	15/3	31/3	17	Semarang	Benoa		1,7
7604	7/4	14/4	8	Benoa	Semarang		1,7
7605	26/4	11/5	16	Tegal	Semarang		2
7606	21/5	31/5	11	Semarang	Semarang		2,5

Ocean and Bali Strait (see also Lohmeyer, this vol.).

The surveys in Nusa Tenggara were intended to provide data needed for fisheries development plans for that area; also, in mid-1974 there was an urgent need for information on the pelagic resources of the Java Sea. Thus, *R/V Lemuru* was transferred to Semarang, Central Java and a series of exploratory fishing surveys were carried out, with emphasis on fish detection and catching aspects.

The arrival of the second FAO biologist (S.C. Venema) in December 1974 coincided with another change in plans: *R/V Lemuru* was to cover the same areas as *R/V Mutiara 4*, viz., the Java Sea (Areas 1 and 2 in Fig. 1), the southern tip of the South China Sea (Area 3), and the Malacca Strait (Area 4) (see Pauly et al., this vol.). However, after only one coverage of the Malacca Straits (see Martosubroto et al., this vol.) and the South China Sea, it was decided to restrict *R/V Lemuru's* area of operation to the Java Sea, with only occasional

incursions to the adjacent straits. The cruises were effected in a more systematic way than before, and the acoustic equipment and fishing gear were upgraded to adapt the vessel to its more scientific role. This program was executed successfully until the end of May 1976, when a shortage of funds made it necessary to transfer *R/V Lemuru* to a regional project in the Persian Gulf.

The Vessel and the Surveys

R/V Lemuru was designed as a purse seiner, but trawl gallows were available and bottom trawl hauls were made very frequently, also for food supply on long cruises. Trolling lines were practically always out during daytime. Light attraction was done frequently in suitable areas, mainly during nights with little or no moonlight. Some oceanographic equipment was also

Box 1. Specifications of *R/V Lemuru*, including equipment and gear.

[Boks 1. Spesifikasi kapal Lemuru, termasuk peralatannya.]

1. Principal dimensions (steel construction)

Length over all	29.35	m
Breadth (moulded)	7.10	m
Depth (moulded)	3.20	m
Designed full load draft	2.55	m
Gross tonnage	165.36	T
Refrigerated fish hold	88	m ³

2. Engines and deck machinery

Main engine Caterpillar D 379 - TA 510 Ps Diesel
Auxiliary Cat. D 320 T, 40 KW
Auxiliary Cat. D 330 T, 75 KW
Main winch, hydraulic, high pressure 50 kg/cm²,
approx. 20 tonnes
Hydrographic winch, hydraulic
Boom winch, hydraulic
Boom swinger, hydraulic
Power bloc, hydraulic
Net sounder winch, manual (as of 1976)
Windlass, hydraulic

3. Electronic equipment

Navigation:

- Koden Direction Finder, KS-500
- Radar 1972-1975 Furuno FR/5/B
- Radar 1976 Furuno FRM 60 range 64 n.m.
- Auto pilot Tokyo Keico
- Compass, Magnetic, Tokyo-Keiki-Seizosho Co. Ltd. (new Dec. 1975)
- Log, one set taffrail log, without bridge repeater

Radios:

- Kelvin Hughes Pentland Bravo, Marine Telephone (1974)
- Radio telephone "Sailor", type 56D
- Public Addressor, Japan Radio Co. Ltd.
- Dymar, Marine VHF FM Radiotelephone Type 801A
- Transistorized Communication Receiver "Eddystone" Model EC - 10 MK. II

Acoustic:

- Koden Multistylus Fishfinder, removed Dec. 1975
- Sanken Sonar, Echosounder, Televigraph, model NTLB 3000 A, 20 and 197 kHz, removed in December 1975

- Sanken Fishfinder New Supergraph NST 300 A, 197 KHz, not in operation during 1976
- Simrad Skipper Sonar SK 3 (transducer shaft bent during cruise 7307. Repaired and dome installed during overhaul, Jan. 1974)
- Simrad EQ, 49 kHz, transducer 62 P, installed December 1975
- Atlas Monograph 58 AN 658, Recorder for net sounder + net sounder installed Dec. 1975
- Portable echosounder, Furuno, used on skiff

Oceanographic and meteorological equipment

- 5 Nansen bottles + reversing thermometers
- 1 Secchi Disc
- Electronic Thermometer, MS-2, Murayama - Denki Ltd.
- Compensated Aeolid Barometer. Not calibrated
- Marine Barometer, Sestrel. Not calibrated
- Anemometer, Thies Göttingen, 0-30 m/s
- Forel Scale, Sea color scale

4. Deck arrangements

- Prior to 1975 overhaul: Purse seine covering full aft of deck. Skiff almost at deck level on starboard. Only one (portside) gallow installed.
- In 1976: Purse seine mainly at starboard side. 2 meters space left at portside. Skiff raised on platform on portside. Second gallow installed, both further forward than before. Midwater trawl operated over purse seine and passage on portside. Much larger working space on deck, especially on starboard side. Sampling conditions much improved.

5. Fishing gear

- Purse seine, 1972, modified 1975 for shallower waters
- AKRA trawl, 1973?
- Semi balloon, shrimp and fish trawl, July 1975
- Semi balloon shrimp trawl
- Midwater trawl Engel, 308 meshes circ. by 800 mm stretched mesh. February 1976
- Skiff, 7 x 2 meters, fiberglass with 2 outboard motors, 40 hp. Johnson
- Underwater lamps (4) - 1000 watt bulbs
Overwater lamps (2) - 500 watt bulbs
Generator for skiff, 2 KW, portable. Honda
- Trolling lines with lures
- Gillnets (hardly used and many not recovered).

available and used.

The most important instrument was a sonar (Simrad SK3), which was used constantly in sweeping mode. Furthermore, the vessel was equipped with two rather primitive echosounders, which were eventually replaced by an echosounder suitable for more refined detection of fish traces (Simrad EQ 50). Further details of the vessel, equipment and gear are presented in Box 1.

The first FAO biologist trained his Indonesian colleagues in the use of oceanographic equipment, sampling and, above all, data recording. During all surveys a well-designed logbook was kept, which did not only contain a record of all major biological events, but also observations on fishing activities along the cruise tracks. The latter were copied on tracing paper at the same scale as the original charts. All data, charts, echosounder paper and sonar paper were taken to the Marine Fisheries Research Institute in Jakarta for processing. Unfortunately, a proper system of data processing and reporting was lacking. After a few attempts, biological cruise reports failed to appear. Although the FAO masterfisher partly compensated for the lack of reports from the biologist by issuing a rather elaborate report of his own (Bjarnason 1977), the overall situation was very unsatisfactory and the data were piling up at an alarming rate. At the beginning of 1975, material collected during more than 300 sea days (30 cruises) had accumulated, while new data were being collected almost continuously.

In early 1975, the highest priority was thus to reduce the cruise tracks to a size where the charts would become manageable, so that observations could be plotted along the tracks. It took about 6 months before this work was completed and then another year of intensive work to plot the data on the charts and to gather all related observations. The (wet!) sonar paper was kept in sealed tins, but by the time it was processed, it had dried out and the echograms were barely visible. Had it not been for the very good logbook records, it would have been impossible to use this material.

Although the bulk of the available data were "saved" in the mimeographed report (Venema 1976), it appears that very few scientists, in Indonesia and elsewhere, have been aware of the existence of this report and hence little use has been made of the data.

Surveys with pelagic gear in tropical seas are notoriously difficult and the number of successful fishing operations with the purse seine and midwater trawl was rather limited. Hence, the amount of biological data on pelagic species was very much lower than the hundreds of samples collected on demersal species during the trawl surveys documented elsewhere in this volume.

An additional drawback for data collection before 1975 is that the number of fishing operations was kept to a minimum in order "not to take fish away from the artisanal fishers". This singular philosophical position of the first FAO biologist resulted in the amount of information available from the lightly fished

pelagic resource of Bali sardinella being very limited. This resource is now overfished (see Pauly et al., this vol.) and data from the early period would be most useful for stock assessment purposes.

Nowadays it is unthinkable that surveys, sampling and data processing should be conducted this way. Indeed, many of the lessons learned from *R/V Lemuru* surveys have been incorporated in FAO manuals and guidelines for managers of fishery research vessels.

The more significant results of the surveys have been plotted on charts. This was done in a rather primitive way, by sticking a variety of symbols along a hand-drawn cruise track, with subsequent reduction by a photocopier.^b Since the fisheries and the data collection based on these have developed very much since the 1976 data report was produced, it was not considered worthwhile to reproduce here all of the data sets in this paper, and only a selection of (redrawn) maps is presented. Interested researchers may get access to the original data via the Marine Fisheries Research Institute in Jakarta, FAO's Fisheries Library in Rome, or the ICLARM Library in Manila, which all hold copies of Venema (1976).

Most of the data in the data report, and hence in this paper, were collected by the biologists Dr. Gede Sedana Merta (1972-76), Dr. Subhat Nurhakim (1973-75) and Mr. Edi Amin (1974-76) and by the assistant biologists, Mr. Isom Hadisubroto (1973-74), Mr. Dadang Karyana (1975) and Mr. Sudjianto (1975-76). Dr. Subhat, assisted by Messrs. Karyana and Sudjianto managed to reconstruct all cruise charts. For all involved in the work with *R/V Lemuru*, the cruises were an unforgettable experience, which generated profound knowledge of major Indonesian sea areas and their resources. It is hoped that this paper will lead to a recognition of the work done by the above Indonesian scientists, during 3.5 years on 48 cruises with a total of 588 seadays and an additional estimated three person-years for data processing.

Environmental Aspects and Migrations of Pelagic Resources

Most fisheries in Indonesia and, in particular, the pelagic fisheries in the Bali Strait, Java Sea and adjacent areas of the South China Sea are seasonal (Bailey et al. 1987). The migration patterns of the most important pelagic resource of the Java Sea (*Decapterus* spp.) and their relation with the monsoons were described by Hardenberg (1937, 1938) and by Potier and Boëly (1990) (Fig. 2 and Box 2).

The influence of oceanic waters is much stronger in the eastern part of the Java Sea, and this results in higher abundances of pelagic fish, in particular layang scads (*Decapterus* spp.). Soemarto (1958), also based on Hardenberg (1937), described their migratory behavior and distinguished three groups of layang, viz.,

1. *East layang* enters the Java Sea with the oceanic water from the Flores Sea during the dry northeast monsoon (May to September). This group may spawn near Bawean Island in June;

^bThe original charts have been lost; Figs. 3-11 were redrawn from photocopies.




Fig. 2. Water masses and pelagic fish migrations in the Java Sea, by season: A) wet monsoon (November to March); B) dry monsoon (May to September). (Modified from Potier and Boëly 1990, and Hardenberg 1937, 1938).

[Gambar 2. Massa air dan migrasi ikan pelagis di Laut Jawa, berdasarkan musim: A) musim hujan (November hingga Maret); B) musim kemarau (Mei sampai September). (Modifikasi dari Potier dan Boëly 1990, dan Hardenberg 1937, 1938).]

2. *West layang* enters the Java Sea with oceanic waters from the Indian Ocean through Sunda Strait during the northwest monsoon (November to March);
3. *North layang* enters the Java Sea with oceanic waters from the South China Sea through the Straits of Gaspar and Karimata, also during the northwest monsoon (November to March).

[The West and North layang tend to avoid a triangular area whose base is formed by the southeastern coast of Sumatra and the island of Bangka; see below].

Based on this, one should expect (1) generally higher abundance of pelagic fish in the eastern Java Sea from May to September, and low abundances towards the end of the wet monsoon in March, and (2) a higher abundance of pelagic fish in the Sunda Strait, along the adjacent North Java Coast and off southwestern Kalimantan from November to March (see

Box 2. Seasonality of pelagic fish abundance in the Java Sea.

[Boks 2. Kelimpahan ikan pelagis di Laut Jawa sesuai dengan musim.]

A. *Dry monsoon* (winds from N.E.) - May to September:

Currents in the Java Sea from East to West, strong influx of oceanic waters with high salinity, in particular in the eastern part of the Java Sea (Area 1).
At first (May) very low abundance of pelagic fish but then rapidly increasing.

B. *Intermonsoon* - October

Weak current, strong mixing.

Highest abundance of pelagic fish.

C. *Wet monsoon* (winds from N.W.) - November to March

Currents in the Java Sea from West to East, heavy rains, strong outflow from rivers, in particular from Kalimantan and lower salinities. Pelagic fish move to the other seas and the central part of the Java Sea. A sharp decline in catch rates starts in December and continues to reach a deep low in March.

D. *Intermonsoon* - April

Currents weaken, waters get mixed. A small peak in abundance can be inferred from catch rates.

also Box 2). However, a complete model of the life cycles of the various *Decapterus* species has still to be developed. It is quite likely, for example, that *Decapterus* spp. do not spawn in the Java Sea, but outside in deep waters, where no fishery takes place. There is no information on the distribution of eggs and larvae, while spent fish has only been encountered a few times in the Java Sea.

It is possible and even likely, that the Java Sea is mainly a nursery area for *Decapterus* spp. A full understanding of the life cycle is, of course, essential for a complete assessment of the stocks of these commercially important species.

The survey results in Areas 1, 2 and 5 as described below may be seen as a contribution to the resolution of this and related issues.

Relationships between environmental conditions and the abundance of Bali sardinella have also been noted (Soerjodinoto 1960; Dwiponggo 1974). For spawning, *S. lemuru*

comes inshore, especially in the northern part of the Strait where salinities are low, in particular during the rainy season. The growing fish move towards the shelf of Bali closer to the Indian Ocean and out of reach of sailing boats operating from Muncar and other harbors in the northern area. So, in the past, the fishery concentrated on juveniles in the northern area, while mature fish were rarely caught. This was still the case during the *R/V Lemuru* surveys from 1972 to 1974.

The very high concentration of densely packed schools of *S. lemuru* is a phenomenon that rarely occurs elsewhere in tropical waters and it is therefore logical to assume that it is

associated with upwelling, as in the case of the Indian "oil sardine" (*S. longiceps*). Previously it was assumed that the abundance of *S. lemuru* was associated with the (weak) upwelling in the open Indian Ocean waters off Java and Sumbawa, identified by Wyrтки (1962). Nowadays it is considered more likely that the occurrence of dense schools of *S. lemuru* is due to the strong upwelling within the Bali Strait. This would also explain why concentrations of *S. lemuru* are rarely found in the adjacent Indian Ocean. Ghofar and Mathews (in Pauly et al., this vol.) discuss the fluctuations on *S. lemuru*

Table 2. Catch by trolling of *R/V Lemuru* by species (A) and time of the day (B), 1972-1976, by areas.

[Tabel 2. Hasil tangkapan tonda menurut daerah oleh kapal penelitian Lemuru berdasarkan spesies (A) dan waktu penangkapan (B), 1972-1976 dan menurut daerah.]

A. Numbers caught, by area									
Species	Area								All (%)
	1	2	3	4	5	6	7	8	
<i>Atule mate</i>	-	-	-	-	-	-	1	-	0.3
<i>Caranx</i> sp.	-	-	-	-	-	-	2	-	0.6
<i>Elagatis bipinnulatus</i>	1	-	-	-	1	7	3	5	4.7
<i>Megalaspsis cordyla</i>	-	-	-	-	-	-	4	4	2.2
<i>Scomberoides commersonianus</i>	-	-	-	2	-	-	-	-	0.6
<i>Chirocentrus dorab</i>	-	1	-	-	-	-	-	-	0.3
<i>Coryphaena hippurus</i>	2	-	1	-	6	2	1	3	4.2
<i>Acanthocybium solandri</i>	-	-	-	-	1	-	3	2	1.7
<i>Auxis thazard</i>	-	-	-	-	-	1	1	3	1.4
<i>Euthynnus affinis</i>	25	9	2	-	-	14	44	19	31.3
<i>Katsuwonus pelamis</i>	1	-	-	2	11	9	2	26	14.1
<i>Sarda orientalis</i>	-	-	-	-	-	-	8	3	3.0
<i>Scomberomorus commerson</i>	4	11	2	-	4	2	7	6	10.0
<i>Scomberomorus guttatus</i>	1	-	-	-	-	-	-	-	0.3
<i>Thunnus alalunga</i> ^{a)}	5	-	-	-	-	4	-	-	2.5
<i>Thunnus albacares</i>	10	-	-	-	-	6	6	15	10.2
<i>Thunnus tonggol</i>	8	6	22	-	-	-	-	-	10.0
<i>Trichiurus lepturus</i>	1	-	-	-	-	-	-	-	0.3
Sphyraenidae	3	-	-	-	1	3	-	2	2.5
Total	61	27	27	4	24	48	82	88	100

B. Percentage of total catch, by hour of the day.									
	1	2	3	4	5	6	7	8	All (%)
5 + 6	8	7	3	0	4	2	6	5	5
7 + 8	18	48	3	25	29	2	16	14	16
9 + 10	41	3	7	50	33	10	23	13	20
11 + 12	8	11	56	0	21	6	21	10	16
13 + 14	13	3	22	0	13	27	10	23	16
15 + 16	10	15	3	0	0	33	18	26	18
17 + 18	2	11	3	25	0	19	6	10	8

^{a)}These may have been misidentified young bigeye tuna (*Thunnus obesus*), as young *T. alalunga* cannot tolerate temperatures > 26°C (Gary Sharp, pers. comm.).

Table 3. *R/V Lemuru* purse seine catches by areas, 1972-1976.

[Tabel 3. Hasil tangkapan purse seine kapal Lemuru berdasarkan daerah, 1972-1976.]

Area	Number of sets		Total catch (kg)					
	Total	Valid	0	1-49	50-199	200-999	1,000-2,999	≤3,000
1	53	49	9	13	5	10	4	8
2	21	18	1	12	1	3		1
3	3	2		1	1			
4	-	-						
5	6	6	1	2		1	1	1
6	4	3	1	1	1			
7	18	17	3	5	2	2	2	3
8	1	1	-	-	-	-	1	-
All	106	96	15	34	10	16	8	13

Table 4. Frequency of occurrence of pelagic species in *R/V Lemuru* purse seine catches over 100 kg.

[Tabel 4. Frekuensi keberadaan ikan pelagis pada hasil tangkapan kapal *Lemuru* yang beratnya lebih dari 100 kg.]

	Area						Total
	1	2	5	6	7	8	
Number of sets	28	5	3	1	9	1	47
Sharks	7	-	-	-	-	-	7
Ariidae	6	-	-	-	1	-	7
<i>Alepes</i> spp.	3	-	-	-	-	1	4
<i>Caranx</i> spp.	2	1	3	-	-	1	7
<i>Decapterus</i> spp.	13	-	-	-	1	1	15
<i>Elagatis bipinnulatus</i>	-	-	1	-	-	-	1
<i>Megalaspis cordyla</i>	4	-	1	-	1	-	6
<i>Selar</i> spp.	7	-	-	-	3	-	10
<i>Seriolina nigrofasciata</i>	1	-	-	-	-	-	1
<i>Scomberoides commersonianus</i>	2	-	-	-	-	-	2
<i>Chirocentrus dorab</i>	2	-	-	-	-	-	2
<i>Dussumieria acuta</i>	2	-	-	-	-	-	2
<i>Sardinella lemuru</i>	5	-	-	-	6	-	11
<i>Sardinella brachysoma</i>	1	-	-	-	-	-	1
<i>Sardinella gibbosa</i>	5	3	-	-	-	-	8
<i>Amblygaster sirm</i>	12	1	-	-	1	-	14
<i>Stolephorus</i> spp.	3	-	-	-	-	-	3
<i>Caesio cuning</i>	4	-	-	-	-	-	4
Lutjanidae	2	-	-	-	-	-	2
<i>Auxis thazard</i>	-	-	1	-	2	-	3
<i>Euthynnus affinis</i>	5	1	1	-	1	-	8
<i>Rastrelliger brachysoma</i>	2	-	-	-	-	-	2
<i>Rastrelliger kanagurta</i>	16	-	1	-	1	-	18
<i>Scomberomorus guttatus</i>	14	-	-	-	-	-	14
Sphyraenidae	5	-	-	-	1	-	6
Diodontidae	1	-	-	-	-	-	1

Table 5. *R/V Lemuru* bottom trawl catch rates, by areas, 1972-1976.

[Tabel 5. Laju tangkapan jaring trawl kapal *Lemuru* berdasarkan daerah, 1972-1976.]

Area	Number of hauls		Duration (hours)	Total catch (kg)	Density (kg-hour ⁻¹)
	Total	Valid			
1	45	39	44	5,593	127
2	37	36	37	1,743	48
3	1	1	1	27	(27)
4	12	12	17	493	41
5	9	9	10.5	1,014	97
6	57	56	81	18,252	225
7	27	21	22	4,624	210
8	30	28	27.5	4,244	154
All	217	201	240	35,990	150

catches from 1950 to 1993.

In the following sections, the *R/V Lemuru* survey results are described by (sub)area, except for the single coverage of the Malacca Strait (partly documented in Martosubroto et al., this vol.) and of the South China Sea (for which Venema 1976 must be consulted).

Prior to this, however, a few tables are presented with summary data for Areas 1-8, for trolling (Table 2), purse seining (Tables 3 and 4) and bottom trawling (Tables 5 and 6). Only few explicit references will be made to these tables, which, however, should be consulted when reading the area-specific accounts below.

Eastern Java Sea, Area 1

Sixteen cruises were made in Area 1; this area is large and therefore the results are discussed by subareas: (a) Madura Strait, (b) North coast of Java and Madura (c) South coast of Kalimantan, and (d) Makassar Strait and the eastern edge of Sunda Shelf. This is followed by general remarks pertaining to the whole of Area 1.

a. The *Madura Strait* is a rather small enclosed area, very shallow on the western side with a channel leading towards Surabaya, and very deep on the eastern side. The shallow part may be a spawning area for Indian mackerel (*Rastrelliger* sp.) while the deep waters are a fishing area for tuna (trolling). A large number of fishers lived on the East Java coast (e.g., Probolinggo, Kembang Island and South Madura). There were always a large number of artisanal crafts (*praus*), especially on the eastern side, where liftnets (*bagans*) were also plentiful. Small handliners were active on the east side of Madura (Raas Strait). The Madura Strait was surveyed only in April-July, thus seasonal patterns could not be determined. Quite a few schools were detected, especially on the shallow side, during cruises 7304 (May-June), 7405 (April-May), 7507 (July, sonar count 130 t) and 7604 (April, sonar count 205 t). The schools were usually small to medium size and not always easy to catch.

b. *North coast of Java and Madura*: Madura is a flat island, with no large harbors on the north coast, which in fact consists of one long beach. The north coast of East and Central Java is mainly low marshland where salt is panned. There are many fishing harbors with, at that time, mainly sailing *praus* fishing payang nets and handlines. Just east of Semarang behind a mountain lies Jepara, a populous fishing harbor, with many fixed liftnets. Near Semarang there were always a large number of active small *praus* as well as a fleet of approximately 50 "chungking" type shrimp trawlers.

The coastal waters are shallow and trawlable in most parts. The islands of Bawean, Masalembu, Arends and the Karimun Java archipelago were also centers of fishing activities, especially Bawean. Concentrations of boats were also found off North Madura, and halfway between Semarang and Surabaya (112°E). There were a large number of *bagans* near Gresik (Surabaya), off North Madura and near Semarang.

The area immediately north of Madura was quite rich in demersal fish (Fig. 3A). Many pelagic schools were also found there in July 1975 (Fig. 3B). The area near Surabaya was rich as well, as was the stretch just north of Semarang. Good concentrations of fish were found frequently near all the island groups (Karimun Java, Bawean, Masalembu). Fish densities

Table 6. Frequency of occurrence (in %) of various taxa in bottom trawl catches of *R/V Lemuru*^a.
 [Tabel 6. Frekuensi (dalam %) tertangkapnya berbagai taksa ikan oleh jaring trawl dasar pada kapal Lemuru^a.]

	Area							
	1	2	4 ^b	5	6	7	8	All
Valid hauls	40	37	12	9	55	23	25	202
Ariidae	70	24	67	33	55	43	16	46
<i>Decapterus</i> spp.	5	5	0	0	2	4	0	6
<i>Megalaspis cordyla</i>	15	3	0	44	2	4	0	6
<i>Selar</i> spp.	18	54	0	22	29	0	32	26
<i>Anadontostoma chacunda</i>	33	16	0	0	4	0	4	11
<i>Dussumieria acuta</i>	40	16	0	33	0	0	0	12
<i>Sardinella lemuru</i>	3	0	0	0	2	13	0	2
<i>Sardinella gibbosa</i>	8	5	0	0	0	0	0	2
Engraulidae	20	16	25	33	16	26	40	22
Formionidae	8	19	8	55	13	17	40	40
Gerreidae	60	57	83	55	9	4	4	11
Harpadontidae	0	3	8	0	9	0	0	3
Leicgnathidae	90	68	83	55	13	17	40	40
Lutjanidae	30	5	58	22	65	0	0	29
Mullidae	90	95	92	88	35	65	60	69
Nemipteridae	43	65	92	88	20	35	52	46
Polynemidae	15	8	8	22	65	0	0	29
Pomadasyidae	53	35	83	22	18	30	32	36
Priacanthidae	45	49	50	66	45	35	24	43
Sciaenidae	60	38	33	44	20	61	44	41
<i>Rastrelliger brachysoma</i>	10	8	0	33	0	0	8	6
<i>Rastrelliger kanagurta</i>	23	11	0	33	0	0	12	9
<i>Scomberomorus</i> spp.	38	19	17	33	15	0	8	18
Sphyraenidae	48	24	17	44	24	17	44	31
Stromateidae	25	5	17	44	0	10	0	9
Synodontidae	85	76	92	88	60	61	64	72
Terapontidae	40	49	50	44	49	61	64	72
Trichiuridae	68	41	50	22	56	30	32	48
Apogonidae	38	16	50	33	9	35	28	25
Diodontidae	8	8	0	0	2	0	0	3
<i>Fistularia</i> spp.	43	38	50	55	5	82	4	24
Platycephalidae	20	35	58	33	9	26	0	21
Tetrodontidae	33	24	0	44	0	0	0	13
Penaeid shrimps	78	76	66	44	71	48	60	73
Squids	70	81	92	77	49	26	40	45
Cuttlefish	50	65	83	33	0	0	24	32

^aExcluding Area 3 (with only one haul).

^bDetails on these 12 hauls may be found in Martosubroto (this vol.) and in Torres et al. (this voi.)




Fig. 3. *R/V Lemuru* survey tracks off northeastern Java and Madura, Area 1 (Cruise 7507): A: demersal schools; B: pelagic schools.
 [Gambar 3. Pelayaran survei kapal penelitian Lemuru di sekitar timur laut pulau Jawa dan Madura, Daerah 1 (Pelayaran 7507): A: gerombolan ikan demersal; B: nerombolan ikan pelanis.]

in the central part of the Java Sea were generally low, except during cruise 7507 (July), when high densities were found almost everywhere (Fig. 3). The same area was virtually devoid of fish during March 1976 (7603).

c. *The south coast of Kalimantan* consists of low lying marshland covered by a dense jungle. There are many shallow bays in which several huge rivers reach the sea (notably the Sampit River, and the Barito River). Near the mouth of the Sampit River the Department of Fisheries had cold storage facilities and an ice plant; however, the only fishing settlements were found at the banks of estuaries, and fishing activities were very low. Fishers from Java were reported to operate there,

but not many were observed from *R/V Lemuru*. A group of three small islands, the Laurot Islands or Pulau Laut Kecil (4-5°N and 115-116°E) seemed to be the best fishing area in the Java Sea, but only a few local fishers were active there.

The south coast of Kalimantan proper was sparsely covered by the survey. A first complete coverage was done during cruises 7505 (May) and 7506 (Fig. 4A) while the whole area was surveyed once more in January 1976 (Fig. 4B).

The coastal area near Banjarmasin, near the mouth of the Barito River, had large fish concentrations in June/July 1975 (7506) (Fig. 4A). The echosounder did




Fig. 4. *R/V Lemuru* survey tracks off southeastern Kalimantan, Area 1: A. Cruise 7506: density of sonar contacts; B. Cruise 7601: density of pelagic schools. [Gambar 4. Pelayaran survei kapal penelitian Lemuru di sekitar tenggara pulau Kalimantan, Daerah 1: A. Pelayaran 7506: kepadatan sonar kontak; B. Pelayaran 7601: kepadatan gerombolan ikan pelagis.]




Fig. 5. *R/V Lemuru* survey tracks around the Laurot Islands, Area 1 (Cruise 7416): A. pelagic schools; B. demersal schools. [Gambar 5. Pelayaran survei kapal penelitian Lemuru di sekitar kepulauan Laurot, Daerah 1 (Pelayaran 7416): A. gerombolan ikan pelagis; B. gerombolan ikan demersal.]

not "confirm" all sonar contacts, but several successful purse seine sets did. A very good catch of shrimp (200 kg·hour⁻¹) was also made south of Banjarmasin.

A concentrated survey was made in Sampit Bay; a large number of small schools were located at the edge of a small trench, the fish consisting of engraulids, clupeids and carangids. The Laurot Islands area (Fig. 5) was surveyed for the first time in October 1974, when many schools were detected. More schools were found during the next cruises in October and November 1974 (Fig. 5). In May 1975 (7505), only a short period of time was spent there, and only medium-size schools were observed, while during Cruise 7506 (June) the area was virtually devoid of fish. In October 1975 (7510), fish was found east and west of the islands, while in March 1976 (7603) several medium-size schools were encountered.

d. *Makassar Strait and the eastern edge of Sunda Shelf.*

The Java Sea ends abruptly where the Sunda Shelf drops off towards the Flores Sea and Makassar Strait. The edge is very rocky with many coral patches and small islands, the largest of which are the Kangean group. The Laut Island is separated from Kalimantan by a narrow, muddy channel, but east of 117°E there are again plenty of rocks and corals, with a steep drop off towards the central Makassar Strait.

The coast of South Sulawesi borders deep waters in the north (Mandar coast) and a shallow shelf area full of small islands and corals south of Barru. Some more coral reefs and islands lie in the middle of the Strait.

This area was surveyed as part of 5 cruises (7416, 7505, 7506, 7510 and 7603). During all cruises results were poor to very poor. Some small to medium-size schools were detected at the edge of the shelf in cruise 7416 (November) (Fig. 5B). Quite a few pelagic schools were detected during cruise 7505 (May), but these consisted mainly of pufferfish and flyingfish. A few bottom schools were detected above bottom outcrops. In June (7506), quite a few sonar contacts were made north of Laut Island and along the edge of the shelf (Fig. 4A), but echosounder recordings were less positive on the latter part.

The number of demersal schools was particularly low. Very few schools were detected on the echosounder in October 1975 (7510), but quite a few sonar contacts were made. In March 1976 (7603), several medium-size bottom schools were recorded on the new echosounder and the area north of Laut was found to be quite rich. Sonar contacts coincided with those recordings.

A few boats were present near small islands and larger numbers near Laut Island (Kotabaru) where there were also many bagans, and at the Mandar coast, especially near Pare Pare and outside Ujung Pandang.

Sightings of flying fish were very frequent in Makassar Strait, as might be expected given the existence of a fishery

on *Cypselurus* spp. in the area. Also observed were tuna schools, porpoises and surface schools; they occurred as well near the edge of the Sunda Shelf and between Bali and Madura.

Light attraction was tried several times just north of Bali, without success. During the four commercial type cruises, light attraction near the Laurot Island was reasonably successful and was followed several times by purse seine sets. In 1975 and 1976, light attraction was less successful, possibly because only a short time was spent in areas with known concentrations.

Out of a total of 52 purse seine sets, the average of the 30 successful sets (> 100 kg) was 2.4 t, with a maximum of nearly 19 t (cruise 7510), consisting mainly of catfish (*Arius* spp.). Some good catches were recorded in Madura Strait (7305, June, and 7604, April). However, fish were not easy to catch in this shallow area.

Bottom trawl hauls were made frequently, especially in the Makassar Strait area. The bottom was found suitable for trawling in most areas except near the drop off in Makassar Strait and Flores Sea. The highest yield was obtained in June, cruise 7506 (669 kg·hour⁻¹, of which 74% were leiognathids), while in one haul over 200 kg of shrimp were caught. The average catch rate of all 39 valid hauls made in Area 1 was only 127 kg·hour⁻¹.

Trolling catches were made during all cruises, and consisted of many different species, with yellowfin tuna and longtail tuna dominating (Table 2).

Midwater trawl was used three times near the Laurot Islands, once yielding a high catch of sharks and black pomfret and five times in Makassar Strait, where three attempts were made to sample the deep scattering layer. This yielded myctophids, small red shrimps and small squids (see also Lohmeyer, this vol.). The same deep sea species were obtained from a similar set in Madura Strait (Cruise 7604).

Overall, the fishing operations in Area 1 did not always match the areas of highest abundance of fish. It appears that there were potentially large fishing grounds south of Kalimantan and near the Laurot Islands which were, in the mid-1970s, practically untouched. Seasonal fluctuations in fish abundance were apparent, although not as obvious as in Area 2 (see below).

High concentrations of fish were found in several parts of Area 1 from April to November, while the South Kalimantan coast was also quite rich in fish in January 1976, under exceptionally bad weather conditions. The Makassar Strait seemed to be poor in pelagic fish. High concentrations of demersal fish were found in areas where bottom trawling was impossible or difficult (two nets lost). Development of demersal fishery at the edge of the Sunda Shelf might be feasible, but not based on bottom trawling.

In 1976, development of a fishery around the Laurot Islands to operate with purse seines or similar gear appears recommendable; it was also recommended to survey the shallow waters not covered by *R/V Lemuru*, especially along the South Kalimantan coast.




Fig. 10. *R/V Lemuru* survey track off southeastern Java, Area 6 (Cruise 7311, 22 November - 3 December 1973).

[Gambar 10. Pelayaran survei kapal penelitian Lemuru di sebelah tenggara pulau Jawa, daerah 6 (Pelayaran 7311, 22 November - 3 Desember 1973).]

smaller fishing villages, of which Pelabuhan Ratu, being relatively close to Jakarta, is the most important, are situated at small bays and near river mouths.

Most fishing activities occurred at a few places: Pelabuhan Ratu, Cilacap, Pacitan (see Figs. 9 and 10), Nusa Barung and Grajagan Bay. A fleet of approximately 80 trawlers fished for shrimp off Cilacap and a few smaller fishing grounds further east.

Schools of porpoises were sighted frequently, as well as jumping tuna, flying fish, etc. A few whale sharks (*Rhincodon typus*) were seen as well. However, the quantities of fish observed with sonar and on the echosounders were not large. In December 1972, July 1973, January 1974, February 1974, and June 1974, when the whole area was "run through", very little fish was recorded. Quite a few schools were observed in August 1973 (7307) along the entire coast between Cilacap and Benoa. In November of that year, a lot of fish was detected in small bays, but none in the deeper waters and in the areas lying between. The charts of Cruises 7411 and 7311 (Fig. 10) show where the fish were concentrated.

Thirty-four light fishing stations were made in this area but only a few were successful. Species attracted included squid, halfbeaks, sardines, Indian mackerel and scad. Most successful were the stations where an anchorage was sought inshore.

Only four purse seine sets were made, of which the first one was sheer disaster, as it took ten hours to haul the net on board. Only one set was successful with a catch of 114 kg of frigate tuna (*Auxis* sp.).

Bottom trawl catches were surprisingly good in this area,

the average catch rate of 225 kg·hour⁻¹ (n = 57) being the highest achieved among all areas (Table 5). Indeed, a single haul of more than 1.5 t·hour⁻¹ was achieved (on Cruise 7311), which, considering the small net as *R/V Lemuru's*, must be considered very good.

The trawlfish species composition did not differ much from that of other areas, although the high occurrence of snappers was noteworthy. Small croakers (Sciaenidae) were also abundant. Leiognathids were also a dominant species here, both in terms of volume and rate of occurrence (Table 5).

Trolling catches were lower than might be expected, especially in 1973. Yellowfin tuna, little tuna and skipjack were caught, as well as blue runners, dolphinfish, Spanish mackerel and barracudas.

The conclusion (drawn in 1976) for this area was that pelagic resources availability was low overall. The results of the bottom trawl hauls were better, although the total stocks may be small. It would seem worthwhile to exploit this resource because it is so close to markets where it is most needed.

Nusa Tenggara, Area 8

The Nusa Tenggara area consists of a chain of islands east of Bali, between the Indian Ocean and the Flores Sea (see also the report of surveys by *Bawal Putih 2* by Martosubroto, this vol.) and includes the sea area between Sumba, Flores and Timor (the "Sawu Sea"). There are a large number of straits between the islands, e.g., the Lombok and Alas Straits, which connect the Pacific with the Indian Ocean. Due to differences in water level between the two oceans, there

is always a strong current towards the Indian Ocean, which hinders navigation, even of vessels like *R/V Lemuru*, and precludes the use of certain fishing gears, such as the purse seine.

The area was visited in four cruises (7303, 7309, 7404 and 7409), which reached as far east as 125°E. Most of the tracks were made over the narrow continental shelves of the various islands, nearly always on the Indian Ocean side, except in cruise 7404 when the north coast of Sumbawa (Flores Sea) was covered as well.

The presence of an FAO-TF Fisheries Project in Lomblen Island stimulated visits to this area in particular, also because *R/V Lemuru* was carrying supplies for the Project. The authorities in Kupang always showed great interest in the vessel's activities in this area, and demonstration fishing was thus carried out a few times.

Local fishing activities were restricted to a few places, as this area is arid and not densely populated. Some coastal concentrations of praus were found in the Alas Strait, around the eastern part of Flores, between Lombok and Sumbawa, and near Lomblen Island.

Sightings of surface schools of flyingfish and tunas were numerous. Also, a large number of whales were seen between Flores and Timor during Cruise 7309 (October-November), while a few whales were also seen off Lombok and Sumbawa in July 1974 (Cruise 7409).

The number of schools picked up by the sonar could be determined from the logbooks only for the 1974 cruises. In March/April, only a few contacts were made, always inshore. In June-July more schools were picked up, also when crossing

deep waters.

The echosounder recorded good schools in Alas Strait and south of Lombok during cruise 7303, but in November only one school was found, and none in March 1974. In a small bay off east Sumbawa, near Kupang (Timor) and in the Lomblen area, several, mostly small schools, were picked up.

A total of 52 light attraction stations was made, 46 times with *R/V Lemuru* anchored in shallow coastal waters and six times while drifting over deep to very deep waters. Squid and sardine-like fish were quite frequently observed near the lamps. Catches were made of those species, as well as of tunas, hardtail scad (*Megalaspis cordyla*) and layang scad (*Decapterus* sp.).

Some very large snappers were caught on handlines, e.g., one *Lutjanus argentimaculatus* of 11.7 kg near Komodo Island. The light attraction stations of Cruise 7409 had very little success, probably due to the full moon, combined with a nearly cloudless sky.

Only one purse seine set was made after light attraction, which yielded a mixture of jacks and squids, of which a large part escaped. The purse seine would have been shot more often if suitable schools and conditions had been encountered.

Twenty-eight valid and two invalid bottom trawl hauls were made in the area, with an average catch of 154 kg·hour⁻¹. One haul in Cempi Bay (8°44', 118°48') yielded 706 kg of mixed small fish (especially Leiognathidae, which were also the dominant family in other good catches). Large to medium-size fish were not encountered, and squid (max 0.2 kg·hour⁻¹) and shrimp (max. 3 kg·hour⁻¹) only in low quantities. Quite a few




Fig. 11. Major fishing grounds of the Java Sea purse seine fishing (redrawn from Boëly et al. 1987); the inset shows a time series of catches from that fishery (smoothed over three months; from Potier et al. 1989).

[Gambar 11. Daerah penangkapan utama purse seine (pukat cincin) di Laut Jawa (digambar kembali dari Boëly et al. 1987); sisipan menunjukkan gambar hasil tangkapan tahunan (data gabungan kuartalan; diambil dari Potier et al. 1989).]

ments of purse seining techniques that simultaneously developed in Bali Strait saved this plan and brought it to fruition.

The development of the pelagic fisheries in the Java Sea is now closely monitored by a project funded by the European Union, and operated by ORSTOM (e.g., Boëly et al. 1987; Potier et al. 1989; see Box 4). The locations of the main fishing grounds of the purse seiners in the Java Sea, with layang scads (*Decapterus* spp.) as the main target species cover the entire Java Sea, always near island groups, and also parts of Makassar Strait and Karimata Strait (see Fig. 11; the match with the results of *R/V Lemuru* is very high). Indeed, the so-called "triangle", the area off southeastern Sumatra reported to be avoided by *layang* (Hardenberg 1937, 1938) is also avoided by the vessels. The seasonalities noted by Hardenberg (1937, 1938) also appear clearly in the purse seine fisheries: peak landings occur in October, while the lowest landings occur around May (Fig. 11, insert).

Thus, while the surveys of *R/V Lemuru* may not have had much influence on the actual development of the pelagic fisheries in Indonesia, they show clearly that it was possible to obtain a good idea of the distribution and relative abundance of the pelagic resources even with the equipment then available. Acoustic surveys with modern echo integrators should thus be, nowadays, even more appropriate to monitor these important resources.

Acknowledgements

This paper came into being thanks to the tireless efforts of Daniel Pauly in documenting research work done in tropical areas. I am very grateful for the opportunity he has offered me to bring to light the results of a major research effort - a fitting expression of our friendship, which, incidentally, began at the Marine Fisheries Research Institute, Jakarta, in mid-1975.

Due to time constraints and distance, this paper was put together by a single author, yet it is based mainly on the work of a number of Indonesian fishery biologists, mentioned above, who are still pursuing the task of assessing the pelagic stocks of Indonesia. I am sure that Bjørn Bjarnason and José Almenar Sansaloni, respectively Captain and Chief Engineer of *R/V*

Lemuru, will also be happy to see some results of their efforts in print.

References

- Bailey, C., A. Dwiponggo and F. Marahudin. 1987. Indonesian marine capture fisheries. ICLARM Stud. Rev. 10, 196 p.
- Bjarnason, B. 1977. Indonesia. Operations of the UNDP/FAO vessel *Lemuru* - 1973-76. A report prepared for the Fisheries Development and Management Project. FI:DP INS/72/064/3, 26 p.
- Boëly, T., M. Potier and Subhat Nurhakim. 1987. Study on the big purse seiners fishery in the Java Sea. (III. The fishing method). J. Pen. Perikanan Laut 47:69-86.
- Dwiponggo, A. 1974. The fishery for and preliminary study on the growth rate of "lemuru" (oil sardine) at Muntjar, Bali Strait. Symposium on Coastal and High Sea Pelagic Resources. Proc. Indo-Pac. Fish. Council. 15 (Section III): 221-240.
- FAO/IOP. 1977. Report of the joint mission to plan development of the *Sardinella* fisheries in the Bali Strait. Tech. Rep. Indian Ocean Programme 15, 51 p.
- Hardenberg, J.D.F. 1937. Preliminary report on a migration of fish in the Java Sea. Treubia 16:295-300.
- Hardenberg, J.D.F. 1938. Theorie omtrent den trek van layang in de Java zee. Med. Inst. Zeevisserij, Batavia 2B:124-131.
- Jones, A. 1976. Development of the aquatic resources of Indonesia. A report prepared for the Fisheries Development and Management project. FI:DP INS/72/064/1, 13 p.
- Potier, M. and T. Boëly. 1990. Influence de paramètres de l'environnement sur la pêche à la senne tournante et coulissante en mer de Java. Aquat. Living Resour. 3:193-205.
- Potier, M., T. Boëly and Subhat Nurhakim. 1989. Study on the big purse seiners fishery in the Java Sea. (VII. Environment of the Java Sea). (Lingkungan Perairan Laut Jawa). J. Pen. Perikanan Laut 51:79-100.
- Smith, P.E. 1971. The horizontal dimensions and abundance of fish schools in the upper mixed layer as measured by sonar. p. 563-591. In G. Brooke Farquhar (ed.) Proceedings of the International Symposium on Biological Sound Scattering in the Ocean, Washington, DC, Maury Center for Ocean Science, Rep. MC-005.
- Soemarto. 1958. Fish behaviour with special reference to pelagic shoaling species: layang (*Decapterus* spp.). Proc. Indo-Pac. Fish. Council. 8(3):89-93.
- Soerjodinoto, R. 1960. Synopsis of the biological data on lemuru, *Sardinella longiceps*. FAO Fish. Synop. 1(15):313-328.
- Venema, S.C. 1976. Report of the operations and results of the UNDP/FAO vessel *Lemuru* in Indonesian waters. A full account of all fishing operations, echo-recordings, observations, cruise tracks and oceanographic data. INS/72/064, 10 parts. pag. var. Development and Management Project. (Mimeo).
- Wyrski, K. 1962. The upwelling in the region between Java and Australia during the southeast monsoon. Aust. J. Mar. Freshwat. Res. 13:217-225.

Demersal Fish Assemblages of Trawlable Grounds off Northwest Sumatra

GABRIELLA BIANCHI

Institute of Marine Research

Division for International Development Programmes

P.O. Box 1870

5024-Bergen, Norway

BIANCHI, G. 1996. Demersal fish assemblages of trawlable grounds off northwest Sumatra [*Kelompok ikan-ikan demersal di daerah penangkapan trawl di sekitar barat laut pulau Sumatra*], p. 123-130. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

Bottom trawl stations from the shelf and upper slope off northwest Sumatra (Indonesia) were analyzed by means of multivariate analysis techniques (detrended correspondence analysis and two-way indicator species analysis) in order to find patterns of zonation among the demersal megafauna assemblages. Apart from the obvious difference between the shelf and slope assemblages, evidence was found for the presence on the shelf of two main species groups, which could partly be explained by a depth gradient. Species composition, with a high percentage of snappers (Lutjanidae) and ponyfishes (Leiognathidae), suggests that, at the time of the survey, the communities in the area were lightly exploited.

Abstrak

Analisis dilakukan terhadap stasiun pengamatan trawl dasar di daerah paparan dan lereng bagian atas di perairan barat-laut Sumatra (Indonesia) dengan teknik analisis multivariate (detrended correspondence analysis dan analisis spesies indikator dua arah) dengan tujuan mencari pola zonasi diantara kelompok fauna besar demersal. Disamping adanya perbedaan nyata antara kelompok fauna di daerah paparan dan lereng, terbukti bahwa sebagian keberadaan dua grup spesies utama terkait dengan tingkat kedalaman. Komposisi spesies, dengan persentase tinggi antara lain ikan kakap merah (Lutjanidae) dan peperek (Leiognathidae), menunjukkan bahwa pada saat survei, tingkat eksploitasi komunitas di daerah survei hanya sedang-sedang saja.]

Introduction

As part of an investigation program of the marine resources of Southeast Asia, the Norwegian research vessel *Dr. Fridtjof Nansen* surveyed the north and northwest coasts of Sumatra between 6 and 30 August 1980. To the author's knowledge no investigation of this type had previously been performed in this area (see also Longhurst and Pauly 1987).

Basic survey results, including biomass estimates for demersal and pelagic stocks, as well as a description of the environmental conditions (hydrography and nutrients) were presented by Aglen et al. (1981). Based on the species composition recorded from the bottom trawl sampling and information on the environmental conditions at each station, an attempt is made here to identify patterns in the distribution of the various species and correlate these with the available environmental information. Because of the very limited coverage in time and space of this survey, the results should be considered as a low-resolution "snapshot" of the demersal assemblages in this area.

This study may be seen as a contribution to "landscape ecology", a relatively new discipline that is distinguished from traditional ecological studies, which usually assume that systems are spatially homogeneous. Landscape ecology stresses that widely used generalizations like "tropical seas" or "tropical communities" do not do justice to the variety of combinations of oceanographic conditions, bottom types and zoogeographic patterns that occur on the ground. Identification and typification of the communities living in well defined "ecological units" will provide a better basis both for understanding the underlying ecological processes and better operational units for management. This will probably involve the definition of broad-scale indices of "landscape" structure which will eventually allow the definition of an appropriate metric for monitoring regional ecological changes. Basic information required will involve definition of appropriate ecological units in terms of physical and biological characterization. The biological characterization should involve among other things species composition, size-spectra, and measures of natural and anthropogenic changes.




Fig. 1. Position of trawl hauls and hydrographic stations, August 1980. [Gambar 1. Posisi stasiun trawl dan hidrografi, Agustus 1980.]

The specific objective of this study was to detect the presence of large-scale trends in the occurrence of fish, cephalopods and other nektonic animals in the trawl catches and from this, to infer zonation among those groups.

Materials and Methods

The area covered by this study includes the north and west coasts of Sumatra, from its northern tip to about 1°S (Fig. 1).

In the north, the shelf is very narrow and steep, and thus trawling was very limited. The two large islands of Simeulue and Nias off the northwest coast also have narrow shelves and are, therefore, mostly untrawable. These islands are separated from the Sumatra shelf by deep basins, except for a narrow connection between Nias and the Sumatra shelf. The northwest coast of Sumatra has a wider shelf with trawable grounds and most of the samples analyzed here were taken in this area.

The west coast of Sumatra is subject to the monsoon gyre circulation system of the northern Indian Ocean (Wyrtki 1973; Roy, this vol.; Sharp, this vol.). During the northeast monsoon, from November to April, prevailing currents in the northern part of the Indian Ocean are from east to west, but

the drift appears to be rather shallow and to exert little influence below the thermocline. During the southwest monsoon (May to October), the main circulation is opposite, and the general movement is eastward. On the other side of the northern Indian Ocean, off Somalia, the waters flow northward, resulting in a well known, strong upwelling. Off Sumatra, the monsoon current flows southward, crosses the Equator, then turns into the South Equatorial Current, thus closing wind-driven gyre of the Equatorial Indian Ocean. Further, during the southwest monsoon, low-salinity water from the Bay of Bengal flows to the southeast along the west coast of Sumatra where the salinity is further reduced by high rainfalls.

Hydrographic sections carried out in the course of the *Dr. Fridtjof Nansen* survey (Aglen et al. 1981) show the presence of a pronounced thermocline between 100 and 125 m. Above the thermocline the mixed layer is rather homogeneous, with temperatures of 28-29°C throughout. Salinity is low, about 33 ppt, which can be explained by the circulation pattern described above. Surface salinity increases southwards to values above 34 ppt off Siberut. Oxygen levels are rather high throughout the shelf and values below 1 ml·l⁻¹ are found only below 200 m depths (Fig. 2).

In the course of the same survey, nutrient analyses (phosphate, nitrate and silicate) were also performed. These showed that, above the thermocline, the water masses were almost depleted of nutrients, possibly due to the lack of exchange with deeper and richer water layers.

A total of 49 trawl hauls were performed with a bottom trawl whose codend had 2-cm meshes (stretched), but whose effective mesh size was smaller, because it was double-lined. Fig. 1 shows the position of sampled trawl and hydrographic stations, while Table 1 shows the number of stations by depth stratum and the area of each stratum. The uneven sampling distribution is due to varying bottom conditions in the above strata.

As a first step, all trawl data, covering 337 species, were entered in the NAN-SIS software package of Strømme (1992), which can be used, among other things, for creating files for analysis using various multivariate methods (see Torres et al., this vol.)

These methods included two-way indicator analysis (TWIA, Hill 1979) and an ordination technique, detrended correspondence analysis or DCA (Hill and Gauch 1980), implemented by the program DECORANA. These methods, originally developed for floristic studies, have proven to be very useful also in marine faunistic studies (McManus 1985, 1989; Bianchi 1991, 1992; Federizon 1992).

Table 1. Number of sampled bottom trawl stations. [Tabel 1. Jumlah stasiun trawl dasar yang diambil contohnya.]

Depth range (m)	No. of hauls	Area (nm ²)
10-25	8	7,350
26-50	18	9,800
51-75	13	4,900
76-99	9	2,450
100-199	4	-
≥ 200	-	-




Fig. 2. Hydrographic profiles of temperature, salinity and oxygen at selected places: a) Breueh, 13 August; b) Kokos, 14-15 August; c) Siberut, 20 August.

[Gambar 2. Profil hidrografi dari suhu, salinitas dan oksigen di tempat-tempat tertentu: A) Breueh, 13 Agustus; B) Kokos, 14-15 Agustus; C) Siberut, 20 Agustus.]

In this study biomass (wet weight) was used as a measure of abundance. Biomass seems to be ecologically appropriate and can be more relevant for practical applications, as for example, for management-orientated studies of fisheries resources. Also, as shown in Bianchi and Høisæter (1992), overall ecological typification is not affected by the abundance measure used by DCA and TWIA when analyzing data covering long gradients.

Each weight (\bar{x}) was converted to $\ln(\bar{x}+1)$ before analysis with DCA. This transformation minimizes the dominant effect of anomalous catches, while the addition of 1 unit is necessary to avoid problems related to the log transformation (required by the presence of zero records). No transformation is necessary in the case of TWIA, where abundances are converted to numbers corresponding to different abundance classes ("pseudospecies"). In this study, five "pseudospecies"

were used, corresponding to classes with lower limits set at 0, 0.5, 5, 50 and 100. These cutoff levels are lower than those used in previous studies (Bianchi 1991, 1992) but they are appropriate, due to the lower fish abundances of the south-west Indian Ocean as compared with the areas covered in the earlier studies.

Depth and bottom type were used in the analysis to study the main gradients along which community changes take place. Temperature, oxygen and salinity were not included. In fact,

as the bottom trawl stations performed in the course of the *Dr. Fridtjof Nansen* survey were all shallower than 100 m (except for a few carried out on the slope), the physical oceanographic parameters were nearly constant, i.e., did not provide environmental gradients in the sampled area.

Bottom type information was derived from the echo-traces along the cruise tracks and classified, according to its suitability for bottom trawling, into three categories: even/flat, uneven and rough bottom. This is a gross classification but it was found

Table 2. Two-way station by *Taxon* table resulting from the program TWINSpan. Values denote abundance categories: 1: $W < 0.5$ kg; 2: $0.5 \leq W < 5$ kg; 3: $5 \leq W < 50$ kg; 4: $50 \leq W < 100$ kg; 5: $W \leq 100$ kg. [Tabel 2. Stasiun (two-way) berdasarkan tabel *Taxon* diturunkan dari program TWINSpan. Angka menunjukkan kategori kelimpahan: 1: $W < 0.5$ kg; 2: $0.5 < W < 5$ kg; 3: $5 < W < 50$ kg; 4: $50 < W < 100$ kg; 5: $W < 100$ kg.]

<i>Taxon</i>	1	2	3	4
<i>Abalistes stellatus</i>	—1—13-2—2211—	41-2-2	—————	—
<i>Alepes djedaba</i>	—————	—————	-11-2—111-14-1—	—
<i>Arius thalassinus</i>	—————2—	3-21—	—14-13—	—
<i>Carangoides armatus</i>	—————1—1—	11—3	—11—31-1	—
<i>Carangoides malabaricus</i>	—————1-2-23-25-1-1-	412355	-1-5-12—1—1	—
<i>Carcharhinus sealei</i>	—————	—332	-33—4—12—	—
<i>Centrophorus</i> sp.	—————	—————	—————	5534
<i>Chlorophthalmus agassizi</i>	—————	—————	—————	5123
<i>Dactyloptena orientalis</i>	-1—121—12-11—	-1—	—————	—
<i>Diagramma pictum</i>	-2—3—	312—2	1—	—
<i>Fistularia</i> sp.	—————112-1-21-1—	-1113-	—————	—
<i>Gazza minuta</i>	—————	-21-45	-13—4—1—1-1—	—
<i>Gymnocranius grandoculis</i>	22—5—1—3—	4—1	—————	—
<i>Harpadon nehereus</i>	—————	—2—	-1—21—531	—1
Jellyfish	—————	—————	4—5555—	—
<i>Lactarius lactarius</i>	—————	1-1-51	1-35113—1-11331-1	—
<i>Leiognathus bindus</i>	—————1—	—1555	4115—4111115-231-	—
<i>Leiognathus equulus</i>	—————	-43-52	5-35-221—15-21—	—
<i>Leiognathus fasciatus</i>	—————	-5—44	-1—	—
<i>Leiognathus leuciscus</i>	—————1—	—1-42	—1—1—1-2—	—
<i>Leiognathus splendens</i>	—————1—	-14-52	-1145431-1—1-5—	—
<i>Loligo</i> sp.	—————1121-1-11-111—1	-111—	—————111—12—	1-1-
<i>Lutjanus erythropterus</i>	—————2-15—534-4-2—	413533	—————4-	—
<i>Lutjanus malabaricus</i>	—————51—3—24—	—————	—————	—
Myctophidae	—————	—————	—————	1113
<i>Opisthopterus tardoore</i>	—————	—————	2-1-2—1—2—2-1	—
<i>Parastromateus niger</i>	—————	—————	-14—3—11—1-1—	—
<i>Pellona ditchela</i>	—————	3-1-52	1-15—3—1—1—	—
<i>Pentaprion longimanus</i>	—————431-12-23-1—	14-151	-1-1—1—1—	—
<i>Peristedion</i> sp.	—————	—————	—————	3111
<i>Polynemus sextarius</i>	—————1—	211-5-	-121-2—	—
<i>Pomadasys argyreus</i>	—————1—	-13-5-	51552432—1-3-3	—
<i>Pomadasys maculatus</i>	—————1-1—1—	413-4-	-14—1—	—
<i>Priacanthus tayenus</i>	—————31—1—32-2—	-3—	-1—	—
<i>Pristipomoides typus</i>	—————53-34—3—35—	—————	—————	—
<i>Rastrelliger brachysoma</i>	—————1—	—1—	-1-1—15—13—	—
<i>Saurida tumbil</i>	—————1-3—1-23-1—	1514—	2341245—1—1—	—
<i>Saurida undosquamis</i>	—————11112-1111111—	—————	—————1—	-1—
<i>Scomberomorus guttatus</i>	—————	—4	4-252-43-2—522	—
<i>Secutor insidiator</i>	—————	—1—2	-1111-112-1—3—	—
<i>Selar crumenophthalmus</i>	-1—2—	—3-53	1—5—21—1—	—
<i>Sphyræna barracuda</i>	-2—1—	-51—	—————411—	—
<i>Sphyræna obtusata</i>	—————	-14-5-	-15—5—1—	—
<i>Terapon theraps</i>	—————111—1—	11115-	21322-1111—1—21-	—
<i>Thenus orientalis</i>	-11—1—1—1—	13-1—	—————1—	—
<i>Thyrsitoides</i> sp.	—————	—————	—————	2112
<i>Trichiurus lepturus</i>	—————1—1—	215-5-	4-541-55-1-1-22411	1—
<i>Upeneus moluccensis</i>	—————1—11—1—22-1—	21-1-1	-1—	—
<i>Upeneus sulphureus</i>	—————1—	-24351	24252141-1—1—1—	—

useful and used in the assemblage-environment correlation analysis. As bottom type is a nominal variable, this had to be transformed in order to allow statistical analysis. This was done by coding each class as a separate variable as suggested in Ter Braak (1987).

Catch rates by depth stratum and average weight were calculated using NAN-SIS (Strømme 1992).




Fig. 3. DCA of bottom-trawl stations on the shelf. The corresponding TWIA Groups (1 to 3) can be recognized by different symbols.
[Gambar 3. DCA dari stasiun trawl dasar di daerah paparan. Kelompok TWIA (1 hingga 3) ditampilkan dengan simbol yang berbeda.]

Table 3. Pearson product-moment correlation coefficient between sample scores on DCA (detrended correspondence analysis) Axes 1 and 2 and environmental variables for shelf stations, Sumatra, August 1980 survey. Values with asterisk indicate significant correlation ($p < 0.05$, $df: 42$).

[Tabel 3. Koefisien korelasi product-moment Pearson antar nilai contoh pada DCA sumbu 1 dan 2 serta variabel lingkungan untuk stasiun-stasiun di paparan, Sumatra, survei bulan Agustus 1980. Nilai dengan tanda bintang menunjukkan korelasi nyata ($p < 0.05$, $df: 42$).]

Variable	Axis 1	Axis 2
Depth	-0.77*	-0.39*
Even	0.39*	0.15
Uneven	-0.23	-0.15
Rough	-0.28	-0.04

Table 4. Average values and standard deviation (s.d. in brackets) for environmental variables used in the analysis, for each of the groups identified by TWIA (n.a. = not available).

[Tabel 4. Nilai rata-rata dan simpangan baku (s.d.) untuk variabel lingkungan yang digunakan didalam analisis, untuk setiap grup berdasarkan TWIA (n.a. = tidak ada data).]

	Group 1 (n=19)	Group 2 (n=6)	Group 3 (n=18)	Group 4 (n=4)
Depth (m)	67 (19)	47 (4)	30 (14)	298 (30)
Even bottom (%)	0.66 (0.47)	0.36 (0.48)	0.89 (0.3)	n.a.
Uneven bottom (%)	0.33 (0.47)	0.26 (0.44)	0.11 (0.3)	n.a.
Rough bottom (%)	0	0.37 (0.48)	0	n.a.

Results

Table 2 presents a summary of the TWIA output, where the trawl stations (represented by columns) are grouped according to their species composition. (All species caught were included in the analysis but only the 50 commonest are presented in Table 2; see Torres et al. (this vol.) for details on a file with the complete list). Four major groups of stations are evident, each with a characteristic species composition. The dendrogram at the bottom of the table shows the hierarchical relationship between these groups, based on species composition. The first division separates the four slope stations from the shelf stations. None of the typical slope species are found on the shelf. Indicator species are the greeneyes (*Chlorophthalmus agassizi*) and the deepwater shark (*Centrophorus* sp.). The second division marks two main groups, one characterized by the lizardfish (*Saurida undosquamis*) and several snapper species (Group 1) and the other by a number of ponyfishes, Family Leiognathidae, the false trevally, *Lactarius lactarius*, the hairtail *Trichiurus lepturus* (Groups 2 and 3). Group 2 represents an intermediate association, with species of Groups 1 and 3. A few taxa appear to be ubiquitous, i.e., *Loligo* spp., *Saurida tumbil* and *Carangoides malabaricus*. It was not considered meaningful




Fig. 4. Plot of station scores on DCA, Axis 1 against depth. Corresponding TWIA Groups (1 to 3) can be recognized by different symbols.
[Gambar 4. Plot nilai stasiun pada DCA, Sumbu 1 terhadap kedalaman. Kelompok TWIA (1 hingga 3) ditampilkan dengan simbol yang berbeda.]

to analyze further divisions due to the small number of available stations.

Fig. 3 shows the results from the analysis using DCA. The groups identified by TWIA are also shown, with exception of the four slope stations, which were excluded to allow better resolution of the plot for the shelf stations. Table 3 shows the results from the correlation between the ordination axes and the values of the environmental variables available for each station. The highest correlation was found between Axis 1 and depth ($r = 0.77$). A positive and significant correlation is also found between Axis 1 and even bottoms, suggesting that the bottom was more even in the shallow than in the deeper part of the shelf. Stations of Group 1 show a great spread along Axis 2 indicating that one or more sources of residual variation are present. Table 4 shows the average values for the bottom characteristics, for each of the groups, confirming the decreasing trend in depth from Groups 1 to 3 (67, 47 and 30, respectively). Highest percentage of bottom type classified as even is found in Group 3, the shallowest, while rough bottoms were




Fig. 6. Box and whisker plot for CPUEs in each depth stratum: 1: 10-30 m; 2: 30-50 m; 3: 50-100 m; 4: (100-300 m). Small square: mean CPUE; box: ± 1.00 SD; whisker: ± 1.96 SD.

[Gambar 6. Boks dan whisker plot dari CPUE di setiap strata kedalaman: 1: 10-30 m; 2: 30-50 m; 3: 50-100 m; 4: 100-300 m. Kotak kecil: CPUE rata-rata; boks: ± 1.00 SD; whisker: ± 1.96 SD.]




Fig. 5. Map showing position of trawl stations after having been assigned to the different groups. Divisions based on TWIA divisive clustering.

[Gambar 5. Peta posisi stasiun trawl setelah dilakukan pengelompokan berdasarkan grup yang berbeda. Pembagian grup dilakukan berdasarkan teknik TWIA divisive clustering.]

classified only in stations of Group 2.

Fig. 4 shows a plot of DCA Axis 1 against depth, the thick line representing the first TWIA division of the shelf stations. Although the correlation with depth is evident, there is a wide spread in depth values especially within Group 1 and Group 3. Unfortunately, detailed environmental information is not available which could be used to interpret these results. In particular, the classification of bottom type (even, uneven, rough) is probably too coarse. For example, the category "even" includes bottoms ranging from sandy to muddy and, i.e., bottom types usually associated with different taxa.

Fig. 5 shows the trawl stations represented by different symbols depending on which group they belong to. The leiognathid assemblages (2 and 3) are the most coastal ones, being located mainly on the shallower part of the shelf. The snapper/lizardfish assemblage is found in the deeper part of the northern shelf and in the southern part of the sampled area.

Table 5 shows the relative abundances of the dominant species in each of the groups. In the deepest group (Group 1), snappers represent 32% of total biomass, while in Groups 2 and 3 ponyfishes represent 53 and 21%, respectively, of the total biomass.

Fig. 6 shows the relative abundances by depth stratum. Highest abundances on the shelf were found between 30 and 50 m, and the lowest were between 50 and 100 m.

Discussion

Apart from the very obvious difference between the shelf and slope assemblages, there is evidence for the presence on the shelf of two main assemblages, which could partly be explained by a depth gradient. Thus a shallow, leiognathid assemblage could be defined, as well as a deeper snapper-lizardfish group (*S. undosquamis*). The two overlap at about

Table 5. Total catch, catch/effort and contribution to the catches for main species in station groups 1 to 4. Catch/effort = total weight caught/number of station in each group.

Tabel 5. Hasil tangkapan total, hasil tangkapan per upaya dan kontribusi terhadap hasil tangkapan untuk spesies utama dalam grup 1 hingga 4. Hasil tangkapan per upaya = jumlah berat total hasil tangkapan dibagi jumlah stasiun di setiap grup.]

Species	Total (catch) (kg)	Catch/effort (kg-hour ⁻¹)	Proportion (%)	(Σ%)
Group 1 (19 stations)				
<i>Abalistes stellatus</i>	19.6	1.0	2.3	50.0
<i>Carangoides malabaricus</i>	40.9	2.1	4.7	42.0
<i>Gymnocranius grandoculis</i>	30.0	1.6	3.4	45.4
<i>Loligo</i> sp.	4.9	0.3	0.5	54.5
<i>Lutjanus malabaricus</i>	73.1	3.8	8.4	32.3
<i>Lutjanus sanguineus</i>	120.6	6.3	13.8	13.8
<i>Nemipterus tambuloides</i>	7.6	0.4	0.9	54.0
<i>Pentaprion longimanus</i>	43.4	2.3	5.0	37.3
<i>Priacanthus tayenus</i>	17.4	0.9	2.0	52.0
<i>Pristipomoides typus</i>	88.6	4.7	10.1	23.9
<i>Saurida tumbil</i>	19.9	1.4	2.3	47.7
<i>Saurida undosquamis</i>	10.0	0.5	1.1	53.1
Others (comprising 124 taxa)	398.2	19.9	45.5	100.0
Total	874.2	46.0	100.0	
Group 2 (6 stations)				
<i>Carangoides malabaricus</i>	109.8	18.3	2.6	67.7
<i>Gazza minuta</i>	122.3	20.4	2.9	62.5
<i>Leiognathus bindus</i>	692.6	115.4	16.3	46.5
<i>Leiognathus equulus</i>	112.7	18.8	2.6	65.1
<i>Leiognathus fasciatus</i>	50.0	8.3	1.2	72.7
<i>Leiognathus splendens</i>	1,286.0	214.3	30.2	30.2
<i>Lutjanus malabaricus</i> (non <i>sanguineus</i>)	74.8	12.5	1.7	71.5
<i>Pellona ditchela</i>	31.5	5.3	0.8	74.3
<i>Pentaprion longimanus</i>	91.0	15.2	2.1	69.8
<i>Pomadasys argyreus</i>	138.9	23.2	3.3	59.6
<i>Pomadasys maculatum</i>	32.5	5.4	0.8	73.5
<i>Trichiurus lepturus</i>	196.9	32.8	4.6	56.3
<i>Upeneus sulphureus</i>	221.2	36.9	5.2	51.7
Others (comprising 117 taxa)	1,096.1	182.7	25.7	100.0
Total	4,256.3	709.4	100.0	
Group 3 (18 stations)				
<i>Lactarius lactarius</i>	85.4	4.7	2.7	40.5
<i>Leiognathus bindus</i>	192.6	10.7	6.1	26.2
<i>Leiognathus equulus</i>	348.7	19.4	11.1	11.1
<i>Leiognathus splendens</i>	101.9	5.7	3.2	37.8
<i>Pomadasys argyreus</i>	155.7	8.6	5.0	31.2
<i>Saurida tumbil</i>	72.1	4.0	2.4	42.9
<i>Scomberomorus guttatus</i>	106.6	5.9	3.4	34.6
<i>Terapon theraps</i>	26.8	1.5	0.9	45.4
<i>Trichiurus lepturus</i>	282.2	15.7	9.0	20.1
<i>Upeneus sulphureus</i>	47.0	2.6	1.6	44.5
Others (comprising 134 taxa)	1,710.3	95.0	54.6	100.0
Total	3,129.3	173.9	100.0	
Group 4 (4 stations)				
<i>Centrophorus</i> sp.	109.0	27.3	20.0	20.0
<i>Chlorophthalmus agassizi</i>	45.0	11.2	8.2	44.8
<i>Peristedion</i> sp.	7.6	1.9	1.4	47.6
Shrimp	90.7	22.7	16.6	36.6
<i>Thyrsitoides</i> sp.	8.0	2.0	1.4	46.2
Others (comprising 70 taxa)	286.8	71.7	52.4	100.0
Total	547.1	136.8	100.0	

40 m. However, depth does not seem a satisfactory variable when considered alone. In fact, as shown in Fig. 4, all three groups have representatives in comparable depth ranges, at least for depths less than about 70 m. Each group also includes stations that do not conform with the general trends. It is thus possible that our trawl stations sampled fish from specific biotypes that were overall not well represented, because of the gear used and/or the sampling design. Bottom type usually is the next important variable, in uniform waterbodies, but no detailed information was available. Federizon (1992) found, for example, that in the central Philippines, two assemblages associated with coralline and sandy substrates of the same depth could be separated.

However, the assemblages defined above appear to be meaningful. Previous studies of this type in Southeast Asia show in fact comparable results. McManus (1985, 1986) reported on a study on the assemblages of the Samar Sea, based on 28 stations at depths of about 10 to 100 m and thus comparable with the shelf stations of the present study. He found a main depth boundary at 40 m, with a shallower community dominated by two leiognathids, while the deeper (> 40 m) community was characterized by *S. undosquamis*, as also found here. However, while his accompanying species are similar, their relative abundances were different. In particular, the snappers are an important element of assemblages 1 and 2 in our study, but do not appear in the deeper assemblage from the Samar Sea, where in the late 1970s trawl fishing had been banned because of overexploitation (Saeger 1981). Thus it is easy to speculate that the scarcity in the Samar Sea, of these highly-priced, long-lived species might be due not to ecological differences, but to the effects of high fishing pressure on community composition. Conversely, this would suggest that in 1980 the fish communities in the *R/V Dr. F. Nansen* survey area were not heavily exploited. A similar suggestion is derived by comparing our results with those of Pauly (1979) and Suvapepun (1991) from the Gulf of Thailand. In this area the dominant group in the trawl catches in the 1960s were leiognathids, followed by a number of groups similar to those found off Sumatra. By the end of the 1970s, the leiognathids, the false trevally (*Lactarius lactarius*) and snappers had disappeared in the Gulf of Thailand from the list of the top 20 species, while squid (*Loligo* sp.) had become very abundant. The species composition off the northwest Sumatra shelf in 1980 suggests that exploitation in this area had not reached those levels. This is also confirmed by the biomass estimate of demersal fish derived from the *Dr. Fridtjof Nansen* survey, of 90×10^3 t (Aglen et al. 1981), which is rather large relative to the estimated catches for that region, for the period 1976-1977, i.e., only about 13×10^3 t·year⁻¹.

Acknowledgements

I wish to thank Daniel Pauly for improving the draft version of this contribution and Mr. Albert Contemprate, ICLARM artist, for drawing the figures.

References

- Aglen, A., L. Fjøn, O.R. Godø, S. Myklevoll and O.J. Østvedt. 1981. A survey of the marine resources of the north and west coasts of Sumatra, August 1980. Reports on surveys with the *R/V Dr. Fridtjof Nansen*. Institute of Marine Research, Bergen, Norway.
- Bianchi, G. 1991. Demersal assemblages of the continental shelf and slope edge between the Gulf of Tehuantepec (Mexico) and the Gulf of Papagayo (Costa Rica). *Mar. Ecol. (Prog. Ser.)* 73: 121-140.
- Bianchi, G. 1992. Study of the demersal assemblages of the continental shelf and upper slope of Angola. *Mar. Ecol. (Prog. Ser.)* 81: 101-120.
- Bianchi, G. and T. Høisæter. 1992. The relative merits of using biomass and numbers in fish community studies. *Mar. Ecol. (Prog. Ser.)* 85: 25-33.
- Federizon, R.R. 1992. Description of the subareas of Ragay Gulf, Philippines, and their fish assemblages by exploratory data analysis. *Aust. J. Mar. Freshwat. Res.* 43:379-391.
- Hill, M.O. 1979. TWINSpan - a Fortran program for arranging multivariate data in an ordered two-way table by classification of individuals and attributes. Cornell University, Ithaca, New York.
- Hill, M.O. and H.G. Gauch. 1980. Detrended correspondence analysis, an improved ordination technique. *Vegetatio* 42:47-58.
- Longhurst, A.R. and D. Pauly. 1987. *Ecology of tropical oceans*. Academic Press, San Diego, California.
- McManus, J.W. 1985. Descriptive community dynamics: background and an application to a tropical fishery. University of Rhode Island. PhD thesis.
- McManus, J.W. 1986. Depth zonation in a demersal fishery in the Samar Sea, Philippines, p. 483-486. In J.L. MacLean, L.B. Dizon and L.V. Hosillos (eds.) *The First Asian Fisheries Forum*. Asian Fisheries Society, Manila, Philippines.
- McManus, J.W. 1989. Zonation among demersal fishes of Southeast Asia: the southwest shelf of Indonesia, p. 1011-1022. In *Proceedings of the Sixth Symposium on Coastal and Ocean Management/ASCE*, 11-14 July 1989, Charleston, South Carolina.
- Pauly, D. 1979. Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries. *ICLARM Stud. Rev.* 1, 35 p.
- Saeger, J. 1981. Do trawling bans work in tropical waters? *ICLARM Newsl.* 4(1): 3-4.
- Strømme, T. 1992. NAN-SIS: software for fishery survey data logging and analysis. User's manual. *FAO Comput. Info. Ser. (Fish.)* No. 4, 103 p. FAO, Rome.
- Suvapepun, S. 1991. Long term ecological changes in the Gulf of Thailand. *Mar. Pollut. Bull.* 23:213-217.
- Ter Braak, C.J.F. 1987. CANOCO - a FORTRAN program for canonical community ordination by (partial) (detrended) (canonical) correspondence analysis, principal components analysis and redundancy analysis (version 3.10). ITI TNO, Wageningen, The Netherlands.
- Wyrčki, K. 1973. Physical oceanography of the Indian Ocean, p. 18-36. In B. Zeitzschel (ed.) *Ecological studies, analysis and synthesis*. Vol. 3. Springer-Verlag, Berlin.

Narrative and Major Results of the Indonesian Module (I) of the JETINDOFISH Project, November 1980 to October 1981

PURWITO MARTOSUBROTO

*Marine Resources Service
Fishery Resources Division
FAO of the United Nations
Viale delle Terme di Caracalla
00100 Rome, Italy*

MARTOSUBROTO, P. 1996. Narrative and major results of the Indonesian Module (I) of the JETINDOFISH Project, November 1980 to October 1981 [*Sekilas kegiatan dan hasil utama Module (I) Proyek JETINDOFISH, November 1980 sampai Oktober 1981*], p. 131-134. In Pauly, D. and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

An account is presented of Indonesian trawl survey cruises conducted from November 1980 to October 1981 in the framework of the multiagency "Joint Eastern Indian Ocean Fisheries Survey" (JETINDOFISH). The area surveyed - mainly through demersal trawling - ranged from east of 115°E to 120°E, and covered the shelf off Lombok, Sumbawa, Sumba, Flores and Timor.

Abstrak

Sekilas disajikan hasil pelayaran survei trawl yang dilaksanakan oleh Indonesia dalam rangka kerjasama Proyek JETINDOFISH sejak November 1980 hingga Oktober 1981. Daerah yang disurvei - khususnya dengan trawl dasar - mencakup wilayah perairan dari sebelah timur pada 115°BT hingga 120°BT, dan meliputi daerah paparan di sekitar Lombok, Sumbawa, Sumba, Flores dan Timor.

Introduction

The paper summarizes the trawl surveys conducted as part of "Module I" of the "Joint Eastern Indian Ocean Fisheries Survey" (JETINDOFISH). The surveys of Module I were conducted in the southeastern part of Indian Ocean, i.e., in the vicinity of Lombok, Sumbawa, Sumba, Flores and Timor (Fig. 1). These surveys were conducted in the context of a project of the Government of Indonesia (through the Directorate General of Fisheries), the Federal Republic of Germany (through the Deutsche Gesellschaft für Technische Zusammenarbeit [GTZ]) and the Commonwealth Government of Australia (through the Common Scientific and Industrial Research Organisation [CSIRO]), under the coordination of FAO. GTZ provided financial assistance and experts for the survey of Module II which covered the area from west of Sumatra to south of Bali, while Australia conducted a survey (Module III) in the northwestern shelf of Australian continent (see Lohmeyer, this vol.).




Fig. 1. Area covered by the *Bawal Putih 2* surveys, and consisting of subarea 8 (South of Lombok and Sumbawa, and west of Sumba) and (south of Flores), details on each haul one given in the database documents by Torres et al. (this vol.)

[Gambar 1. Daerah yang disurvei kapal *Bawal Putih 2* yang meliputi subarea 8 (bagian selatan Lombok dan Sumbawa serta bagian barat Sumba) dan subarea 9 (bagian selatan Flores), rincian tiap tarikan trawl terdapat dalam dokumen Torres et al. (dalam buku ini).]

Box 1. Vessel specifications of *R/V Bawal Putih 2*.
 [Box 1. Spesifikasi kapal Bawal Putih 2.]

Type: stern trawler

Main dimensions:

Length O.A.	36.0 m
Length B.P.P.	30.0 m
Breadth MLD	9.4 m
Tonnage:	
GRT	370.3 t
NRT	115.5 t

Engine specifications:

1. Main engine (1)	: Caterpillar D 399
No. of cylinder	: 16
High idle engine RPM	: 1,345
Pull load engine RPM	: 1,200
HP at sea	: 1,125
2. Auxillary engine (2)	: Yanmar D 6 KFL
HP	: 170




Fig. 2. Technical features of the trawl used during the *Bawal Putih 2* surveys.

[Gambar 2. Gambaran teknis dari jaring trawl yang digunakan dalam survei dengan kapal Bawal Putih 2.]

The Surveys

The surveys of Module I were conducted by the *R/V Bawal Putih 2*, a stern trawler of 350 GRT owned by the Directorate General of Fisheries (Box 1). The scientific teams and the crew were Indonesian and surveys were conducted independently of those in Modules II and III. However, most activities and sampling protocols were standardized by an Advisory and Technical Committee composed of members of all agencies involved in the JETINDOFISH project, while FAO provided a project coordinating officer at the project office in Denpasar, Bali.

The survey was conducted during the period from November 1980 to October 1981. Owing to limitations of its rigging, the trawling operations of *Bawal Putih 2* were conducted only at depths of less than 100 m. To compensate for this limitation, a small number of hauls at depths of more than 100 m were done by *R/V Jurong* of Module II (see Lohmeyer, this vol.) in July 1981; these covered the shelf of Lombok Island and its vicinity (subarea 8).

A total of eleven cruises was made during the survey period. Two or three biologists and three or four technicians were onboard in each cruise. A FAO associate professional officer (APO) from Denmark on fish taxonomy participated in four of the eleven cruises, to assist the biologists with fish identifications.

The Vessel

The *R/V Bawal Putih 2* is a steel stern trawler of 350 GRT (Box 1), equipped with an Engel-type high opening bottom trawl (Fig. 2). In the absence of acoustic equipment, echotracing for pelagic fish resources could not be performed (but see Venema, this vol. for an echoacoustic survey of the area in Fig. 1). Measurements of oceanographic parameters during

the survey were limited to temperature profiles, obtained through bathythermographs.

Sampling and Fish Species Identification

Sorting and sampling of the catches were done directly onboard after the catches had been dumped onto the rear deck. The sampling procedures were the same as for the other Modules of the JETINDOFISH project (see Lohmeyer, this vol.).

Shipborne identifications were based on FAO Species Identification Sheets (Fischer and Whitehead 1974) and "The Fishes of New Guinea" (Munro 1967).

Results and Discussion

Eleven cruises of *R/V Bawal Putih 2* completed during the survey period resulted in 121 trawl hauls in areas 8 and 9.

Surface temperatures during the survey period ranged from 26 to 30°C, but fell down to a range of 8.5 to 10°C at 300 m.

Table 1. Distribution of average catch rate (C/f; kg-hour⁻¹) and stock density (D, in t·km⁻²) by subarea (8 or 9), depth and monsoon (east or west monsoon)^a. [Tabel 1. Penyebaran rata-rata hasil tangkapan per upaya (C/f; kg/jam) dan kepadatan stok (D, dalam t·km⁻²) berdasarkan subarea (8 atau 9), kedalaman dan monsun (timur atau barat)^a.]

Sub-area	Monsoon	N ^a	0-50 m			50-99 m			100-149 m ^b			150-199 m ^b			>200 ^b			All depths	
			C/f	D	N	C/f	D	N	C/f	D	N	C/f	D	N	C/f	D	N	C/f	D
8	E	34	251	2.100	46	287	2.991	10	204	2.370	1	31	0.309	2	128	1.215	93	256	2.326
	W	30	264	1.987	16	191	1.778	-	-	-	-	-	-	-	-	-	46	231	1.892
	E	3	776	5.432	7	320	2.148	-	-	-	-	-	-	-	-	-	10	646	4.493
9	W	9	178	1.053	2	411	2.852	-	-	-	-	-	-	-	-	-	11	211	4.500

^aN = Number of hauls.

^bHauls by FRV Jurong.

Box 2. Small-Scale Fisheries Development Project in Lombok, Nusa Tenggara Barat, Indonesia.

[Boks 2. Proyek Pengembangan Perikanan Skala Kecil di Lombok, Nusa Tenggara Barat, Indonesia.]

In 1983, the DGF initiated the "Small-Scale Fisheries Development Project on Bali and Lombok" (SSFDP), implemented again with German technical assistance through the Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ). Because of the comparatively high development of the fisheries around Bali, the project since 1986 concentrated entirely on Lombok and adjacent waters. SSFDP responded directly to the DGF (Resources Management Division) in Jakarta, but was attached to the provincial fisheries office (Dinas Perikanan Tingkat I) in Mataram, Lombok.

In its 10 years of existence (1983-1993), SSFDP went through several phases of organization and implementation, all aimed at promoting employment in and income from small-scale fishing and related fields, while preserving the natural resources. The project staff therefore worked not only on fishing technology, resource monitoring and fish processing, but also considered local social and economic conditions. This entailed extensive training programs for the target fishermen and their wives in organizational, technical and economic aspects.

Central to the project implementation was the support and guidance given to fishing groups established by local fishers from Labuhan Lombok at the east coast of Lombok. Each group consisted of four members, operating a newly designed fishing boat equipped with tuna drift gillnets, acquired on a credit basis. Aside from the formation of 30 fishing groups, the important task was to identify and establish an appropriate organizational framework (a self-supporting or cooperative structure) in order to gain access to the credit program of the Regional Development Bank (Bank Pembangunan Daerah). Due to lack of collateral, credit facilities would not easily be available to individual fishers. Close cooperation was maintained between the fisheries office and the bank, but also with nongovernment organizations (NGOs), especially in the fields of extension and training.

Resource monitoring was the main task of the section of SSFDP devoted to fisheries biology. Target species of the developing tuna drift net fisheries were highly migratory fish, distributed over a much larger area than the actual fishing ground. No stock assessment of the pelagic resources of Alas Strait and adjacent waters was carried out, since this would have been beyond the scope of the project; the chosen approach was therefore to monitor the catch and effort of the developing fishery. Information on yields and catch rates could also immediately be used to formulate extension advice both in technical and economic terms. Dominant target species was skipjack (*Katsuwonus pelamis*) with 54%, followed by frigate tuna (*Auxis thazard*) and kawakawa (*Euthynnus affinis*) with 17 and 7% of the catch, respectively.

The recommended fishing gear was based on a series of fishing gear trials and comparisons with other drift net fisheries south of Java and Sumbawa. Various parameters such as mesh size, fishing (float-line) depth, hanging ratio, trimming, fishing times and increased net length were tested. Since personnel constraints had to be considered in this nonmechanized fisheries, the recommended drift net had a length of about 1,250 m, consisting of 25 single net pieces (nylon multifilament D12-D15) of 50 m each; net depth was around 12 m (140 meshes) with mesh sizes ranging from 10 to 15 cm. Nets drifted near-surface and were set during night time for approximately six hours at a distance of about 30-50 km from the coast. By introducing drift gill nets and motorized fishing boats, SSFDP avoided conflict between the traditional and the newly formed fishing groups. Fishing boats were designed by project staff and produced both by private boat yard and a project run

facility, which was later privatized. The boats had a length of 7.6 m, powered by 10-15 hp inboard diesel engines, and were built of fiberglass laminated marine plywood, incorporating a wash deck and insulated fish boxes into the design.

The section of SSFDP devoted to fish processing and marketing tested and introduced improvements in the formulation of traditional products like boiled salted fish (*pindang*), fish floss and fish cracker (*krupuk*). At the same time, attention was also focused on the hygienic aspects of fish handling and the economic aspects of marketing, in order to assure better income not only from increased amounts of fish marketed but also from higher fish prices. SSFDP assisted in the setting up of a functional quality control laboratory at the fisheries office in Mataram. Here, routine control tests such as organoleptic evaluation, chemical and microbiological quality analyses were carried out on fish, invertebrates and processed market products.

Advisory (extension) services to the fishing groups in group formation and management, as well as the development of a financial system for the acquisition of the fishing system was the central task of SSFDP. Group formation involved a stepwise process of application, interview and selection, involving project staff, village leaders and NGOs, as well as a bank representative. Intensive training both ashore and at sea was followed by a practice period, during which a group could use a project owned boat for free, but under supervision. This period served to assess the group's skills and chances of success and to accumulate own funds (collateral) for the downpayment of 10% of the boat price. The credit contract for boat and equipment was signed by all four members. The interest rate for the loan was 12% per year, with a loan period of up to five years. In order to assure a better administrative and legal basis, fishing groups joined together to form a pre-cooperative. Its functions were to solve any coordination problems, make joint purchases and to act as a channel for the credit repayment procedures.

Monitoring was not only extended to the resources, but also the economic and social situation of the individual fishing groups (impact monitoring) and the process of project implementation (activity monitoring). When necessary, training and extension input required was tailored according to needs. Apart from monitoring and evaluation, several other concept elements of SSFDP need to be underlined: activities were designed to stimulate the self-help potential of the target groups, long-term supply and operation of the boats were assured through privatization of boat yard and maintenance workshop, and the viability of the financing concept was established with the credit scheme provided by a local bank.

During the last years of implementation of SSFDP, independent boat owners operating at the east and south coast of Lombok started to equip their units with drift nets recommended by the project, adding 20-30 boats to the SSFDP-assisted fisheries.

Rudolf Hermes

Marine Biologist

Department of Fisheries and Marine Resources

Momase Coastal Fisheries Development Project

P.O. Box 4197

Lae, Papua New Guinea

Table 2. Estimate of potential yield by fish category^a and subarea.

[Tabel 2. Perkiraan potensi hasil tangkapan berdasarkan katagori^a grup ikan dan subarea.]

Sub-area	Area swept (km ²)	Potential yields (t·year ⁻¹)						Total
		A	B	C	D	E	F	
8	2,470	1,155	4,977	12,489	36,885	18,300	45,701	119,507
9	800	11,831	142	1,055	14,237	1,858	15,320	44,443
Total	3,270	12,986	5,119	13,544	51,122	20,158	61,021	163,950

^aA to F are fish categories (see text).

For analysis, the fish catches were grouped as follows (see also Table 1):

- A : sharks and rays;
- B : sardines (*Sardinella* spp.) and anchovies (Engraulidae);
- C : Carangidae;
- D : Lutjanidae, Sparidae, Lethrinidae;
- E : Nemipteridae, Polynemidae, Trichiuridae, Ariidae, Bothidae;
- F : Leiognathidae, Gerreidae, Mullidae, Synodontidae

Estimation of fish stock density was based on the swept area method, with an assumed escapement factor of 50%. The potential yield estimation followed the method of Gulland (1971) with an assumed average annual natural mortality of 0.5 year⁻¹. As the survey area is under the influence of monsoonal regime (Sharp, this vol.; Roy, this vol.; Venema, this vol.), the catch rates were grouped according to monsoon (east monsoon: April - October, west monsoon: November - March). Table 1 summarizes the distribution of catch rate and

of estimated fish stock densities according to monsoon and depth category, while Table 2 presents the estimate of potential yields by fish group and area. The abundance (potential) of group D, the most valuable group of species, is relatively high, as noted during the *R/V Lemuru* surveys (Venema, this vol.)

Overall, the survey did not indicate the area to show a sufficient potential for the development of trawl fisheries; this led to the initiation of a project on small-scale fisheries around Lombok island (Box 2).

Pauly et al. (this vol.) and Torres et al. (this vol.) present further analyses based on the data from the *R/V Bawal Putih 2* surveys, and details on the stations, respectively.

References

- Fischer, W. and P. Whitehead, Editors. 1974. FAO species identification sheets for fisheries purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). Vol. I-IV. FAO, Rome.
- Gulland, J.A. Editor. 1971. The fish resources of the ocean. Fishing News Books, Farnham, Surrey, UK.
- Munro, I.S.R. 1967. The fishes of New Guinea. Department of Agriculture, Stock and Fisheries, Port Moresby, New Guinea.

Fishery Biology of 40 Trawl-caught Teleosts of Western Indonesia^a

D. PAULY^b

A. CABANBAN^c

F.S.B. TORRES, Jr.

*International Center for Living Aquatic
Resources Management
MCPO Box 2631, 0718 Makati City
Philippines*

PAULY, D., A. CABANBAN and F.S.B. TORRES, Jr. 1996. Fishery biology of 40 trawl-caught teleosts of Western Indonesia [*Biologi perikanan dari 40 ikan teleostei yang tertangkap oleh jaring trawl di perairan Indonesia bagian barat*], p. 135-216. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

A review of the biology of 40 fish species abundant in bottom trawl catches in Western Indonesia is presented. This emphasizes geographic and depth distribution, based on surveys conducted from 1974 to 1981 by the research vessels *Jurong*, *Mutiara 4*, *Dr. Fridtjof Nansen*, *Lemuru* and *Bawal Putih 2*, and biological information (growth, length-weight relationships, food and feeding habits) estimated from the survey data and/or extracted from FishBase, the computerized encyclopedia of fish.

Abstrak

Tulisan ini menyajikan suatu tinjauan terhadap biologi dari 40 spesies ikan yang banyak terdapat dalam hasil tangkapan trawl di perairan Indonesia bagian barat. Tulisan ini menekankan penyebaran secara geografis dan kedalaman, berdasarkan survei yang dilaksanakan dari tahun 1974 hingga 1981 oleh kapal-kapal penelitian Jurong, Mutiara 4, Dr. Fridtjof Nansen, Lemuru dan Bawal Putih 2, serta informasi biologi (pertumbuhan, hubungan panjang-berat, makanan dan kebiasaan makan) yang diperoleh dari data survei dan/atau diambil dari FishBase, suatu ensiklopedia ikan dalam bentuk perangkat lunak komputer.

Introduction

The following review of the biology of 40 trawl-caught species of Western Indonesia was written for a number of interrelated purposes:

- 1) to serve as repository for selected information on commercially (or potentially) important fish resources, extracted from the trawl surveys documented elsewhere in this book;
- 2) to make available, in a single source document, key parameters on the biology of these important species for stock assessment and related purposes in Indonesia and other countries with similar ichthyofauna;
- 3) to refute for audiences elsewhere, the often-stated but increasingly untrue statement that "nothing is known on the biology of tropical fishes" and, last but not least;
- 4) to illustrate how information extracted from FishBase, the computerized encyclopedia of fishes

(see Froese et al., this vol.) can be combined with field data to characterize any species of fish.

Materials and Methods

The catch/effort data obtained during the trawl survey of *Jurong*, *Mutiara 4*, *Dr. Fridtjof Nansen*, *Lemuru* and *Bawal Putih 2*, documented in Lohmeyer (this vol.), Bianchi (this vol.), Martosubroto (this vol.), Pauly et al. (this vol.), Bianchi et al. (this vol.) and Torres et al. (this vol.) were used to identify 4 important teleosts species of Western Indonesia, listed on Table 1 in taxonomic order. For each species, the following information is presented, so far available:

- i) Valid scientific name (including author and date), and common names, in English and Indonesian when available (see Froese et al., this vol.);
- ii) A brief description of the distinctive characteristics of the species including meristic counts, adapted wherever possible from the appropriate FAO species catalogues. The graph illustrating each species was either scanned, or redrawn by Mr. Robbie Cada, of the FishBase project, based on various sources. Maximum lengths are given for each species, and may refer to total length (TL), fork length (FL) or standard length (SL); these codes are omitted when the length type could not be determined from the reference used. These

^aICLARM Contribution No. 1315.

^bAlso at Fisheries Centre, The University of British Columbia, 2204 Main Mall, Vancouver, B.C. Canada V6T 1Z4; e-mail: pauly@fisheries.com

^cPresent address: Marine Research Unit, Universiti Malaysia Sabah

- a) reported maximum length of any specimen of the species in question, from locations outside Indonesia (here coded L_{max1});
- b) maximum length in Indonesia, as observed during the surveys reported upon in this volume or related publications (here coded L_{max2});
- c) maximum length (and 95% confidence interval) that may be expected in Indonesia, based on the maxima of a series of length-frequency samples and on extreme value theory (Formacion et al. 1991). Such values are here coded L_{max3} , and are presented along with the graph through which they were estimated, themselves outputs of the FiSAT software (Gayanilo et al. 1996);
- iii) Geographic distribution: outside Indonesia through a brief text, and within Indonesia through a map generated by the MAPPER software (Coronado and Froese 1994) and showing the occurrences of each species at stations covered by the surveys documented in this volume;
- iv) Graphs illustrating the depth distribution of each species in a survey conducted in Western Indonesia;
- v) A brief account of the biology of the species. Emphasis herein is given to habitats, food and feeding habits and, so far available, to estimates of the von Bertalanffy (1951) growth function (VBGF) for the species in question, either in Indonesia or elsewhere. The VBGF has, for length, the form

$$L_t = L_{\infty} (1 - \exp(-K(t-t_0))) \quad \dots 1$$

where L_{∞} is the mean length the fish of the population would reach if they were to grow indefinitely (here always in cm), K is the rate at which L_{∞} is approached, and t_0 is the theoretical

Box 1. Estimating the parameters of length-weight relationships from length-frequency samples and their weights.

[Boks 1. Estimasi parameter hubungan panjang-berat dari contoh frekuensi-panjang dan berat.]

Length-weight relationships, in fisheries biology, usually take the form

$$W = a \cdot L^b \quad \dots 1$$

where W is the body weight (live or gutted) of the fish, a is a multiplicative factor, L a linear measure (e.g., total or fork length) of the fish body, and b is an exponent, usually close to 3 but which may range from 2.5 to 3.5 and exceptionally from 2 to 4.

Estimating the parameters of such relationships is usually straightforward, and is usually done by plotting the logarithms of the available individual weights against the logarithms of the corresponding lengths, i.e.,

$$\log(W) = \log(a) + b \log(L) \quad \dots 2$$

and using a Type I (or predictive) linear regression to estimate $\log(a)$ and b . Variants of this approach exist, but this need not concern us here, as we deal below with cases where the available data do not consist of L-W data pairs.

During the demersal trawl surveys described in this volume, there was often not enough time for fully analyzing the catch of one station before the catch of the next station was hauled in; such cases resulted in aggregated data, i.e., samples of fish that had been *measured* individually, leading to length-frequency samples (L/F), but not *weighted* individually. Thus, only the bulk weights of the L/F are available (accurate shipborn weighting of small fishes was usually not possible anyway).

We present here a new method to estimate a and b in length-weight relationships using such data; this requires the computation of "pseudoweights", i.e., of sample weights obtained using estimates of the parameters a and b of a length-weight relationship.

Estimating the pseudoweight of samples requires an accurate estimator of the mean weight (\bar{w}_i) of the fish within a given length class (i), which is not equal to the weight corresponding to the midpoint of that length class, or midlength. For this, we use

$$\bar{w}_i = (1 / L_{i+1} - L_i) \cdot (a / b + 1) \cdot (L_{i+1}^{b+1} - L_i^{b+1}) \quad \dots 3$$

where a and b are as defined in Equation (1).

The pseudoweight (W'_j) of a given sample (j) is then estimated from

$$W'_j = \sum_{i=1}^{n_j} (\bar{w}_i \cdot f_{i,j}) \quad \dots 4$$

where:

\bar{w}_i is the mean weight of class i (Equation 3);

$f_{i,j}$ is the frequency of class i in sample j ; and

n_j is the number of classes in sample j .

When a number (≥ 3) of length-frequency samples and their bulk weights are available, a and b can be estimated iteratively, using arbitrary seed values of a and b (e.g., $a = 0.01$ and $b = 3$), and using a nonlinear least squares procedure (here: Marquardt's compromise algorithm) which minimizes the sum of the squared differences (SSE) between the sample weights (W_j) and the pseudoweights (W'_j), both previously log-transformed to stabilize the variance, or

$$SSE = \sum [\log(W_j) - \log(W'_j)]^2 \quad \dots 5$$

The results of the final iteration can be shown by plotting the sample pseudoweights against the observed sample weights; this leads to graphs such as shown in this contribution, which can be used to identify outliers.

These steps are all quickly performed by a new software, ABee, available from ICLARM, and which includes a version of Marquardt's algorithm that provides standard errors for all parameter estimates.

Daniel Pauly and Felimon Gayanilo, Jr.

age the fish would have had at length zero if they had always grown according to the VBGF. This parameter, difficult to estimate in the absence of *absolute* age data (i.e., when L_{∞} and K are estimated from length-frequency data, as is also the case for Indonesian fishes) is not given here. Few stock assessment models require t_0 , in any case.

A length-weight relationship of the form

$$W = a \cdot L^b \quad \dots 2)$$

is also given for each species. The parameters *a* and *b* of equation (2) are usually estimated from the intercept and slope, respectively of a linear regression i.e.,

$$\log(W) = \log(a) + b \log(L) \quad \dots 3)$$

A new method was developed, while compiling this review, to estimate the parameters *a* and *b* of such relationship from length-frequency samples and their bulk weights (Box 1). This

method was applied wherever suitable data were available.

vi) Finally, for each species, we give the (FishBase) numbers of the references documenting the sources of data in (i) to (v) (see Appendix I for full references).

Results

The 40 species considered in this review are listed in Table 1, in the sequence also used for presentation of results on a per-species basis.

Table 1. Classification (from Eschmeyer 1990; see also Froese et al., this vol.) of 40 trawl-caught teleosts of Western Indonesia. [Tabel 1. Klasifikasi (menurut Eschmeyer 1990; lihat juga Froese et al., dalam buku ini) dari 40 spesies ikan demersal penting di Indonesia bagian barat.]

Clupeiformes		
Clupeidae	14 <i>Parastromateus niger</i>	Mullidae
1 <i>Amblygaster sirm</i>	15 <i>Selar crumenophthalmus</i>	29 <i>Upeneus moluccensis</i>
2 <i>Dussumieria acuta</i>	Gerreidae	30 <i>Upeneus sulphureus</i>
3 <i>Pellona ditchela</i>	16 <i>Pentaprion longimanus</i>	Nemipteridae
4 <i>Sardinella gibbosa</i>	Haemulidae	31 <i>Nemipterus thosaporni</i>
5 <i>Sardinella lemuru</i>	17 <i>Diagramma pictum</i>	Priacanthidae
Siluriformes	18 <i>Pomadasys argenteus</i>	32 <i>Priacanthus macracanthus</i>
Ariidae	19 <i>Pomadasys maculatus</i>	Scombridae
6 <i>Netuma thalassina</i>	Lactariidae	33 <i>Rastrelliger kanagurta</i>
Aulopiformes	20 <i>Lactarius lactarius</i>	34 <i>Scomberomorus commerson</i>
Synodontidae	Leiognathidae	35 <i>Scomberomorus guttatus</i>
7 <i>Saurida micropectoralis</i>	21 <i>Leiognathus splendens</i>	Sphyraenidae
8 <i>Saurida undosquamis</i>	22 <i>Leiognathus bindus</i>	36 <i>Sphyraena obtusata</i>
Perciformes	23 <i>Leiognathus equulus</i>	Stromateidae
Carangidae	24 <i>Leiognathus leuciscus</i>	37 <i>Pampus argenteus</i>
9 <i>Carangoides malabaricus</i>	25 <i>Gazza minuta</i>	Terapontidae
10 <i>Caranx ignobilis</i>	Lethrinidae	38 <i>Terapon jarbua</i>
11 <i>Caranx tille</i>	26 <i>Gymnocranius grandoculis</i>	Trichiuridae
12 <i>Decapterus macrosoma</i>	Lutjanidae	39 <i>Trichiurus lepturus</i>
13 <i>Decapterus russelli</i>	27 <i>Aprion virescens</i>	Tetraodontiformes
	28 <i>Pristipomoides typus</i>	Balistidae
		40 <i>Abalistes stellatus</i>

Amblygaster sirm (Walbaum, 1792)

Spotted sardinella (English); sardin (Indonesian).

Scutes not prominent. Distinguished from *A. leiogaster* and *A. clupeioides* by the presence of a series of 10 to 20 gold (in life) or black (on preservation) spots down the flank (but sometimes missing) and more lower gillrakers, and from *Sardinella* species by its fewer pelvic finrays and lower gillrakers. Dorsal spines: 0-0; soft rays: 13-21; anal spines: 0-0; soft rays: 12-23. $L_{max1} = 26$ cm (Sudan, Red Sea); $L_{max2} = 20$ cm; $L_{max3} = 22.7$ cm TL (Fig. 1A). See Fig. 1B and Table 2 for length-weight relationship.

Indo-West Pacific: coasts of Africa, including Red Sea and Madagascar to Southeast Asia (Fig. 2). Extending northeastward to Taiwan, and Okinawa (Japan), and southeastward to New Guinea, the northern coasts of Australia and Fiji.

A schooling species occurring in coastal waters. Depth range: 10-75 m (Fig. 3). Feeds mainly on small crustaceans and their larvae, larval bivalves and gastropods, as well as phytoplankton (e.g., *Peridinium*, *Ceratium*). Table 3 presents three sets of growth parameters from Indonesia.

References: 171, 188, 312, 762, 823, 1263, 1314, 1439, 1442, 1443, 1444, 1447, 1488, 1602, 1911, 2178, 2857, 3785, 4615, 5213, 5525, 5542, 5730, 5736, 5756, 5763, 6313

Table 2. Length-weight (g/[TL;cm]) relationship of spotted sardinella, *Amblygaster sirm*, in Indonesia. [Tabel 2. Hubungan panjang-berat (g/[TL;cm]) dari ikan sardin, *Amblygaster sirm*, di Indonesia.]

Parameter	Estimate
a	0.1177
s.e.(a)	0.1265
b	2.0748
s.e.(b)	0.3688
r ²	0.9933

Table 3. Growth parameters of spotted sardinella, *Amblygaster sirm*.

[Tabel 3. Parameter pertumbuhan ikan sardin, *Amblygaster sirm*.]

Parameter	A	B	C
L _∞ (TL, cm)	25.2	25.8	24.3
K (year ⁻¹)	1.175	1.150	0.586

A. "Java Sea" (Ref. 1447)

B. Off Pekalongan, North/Central Java (Ref. 1314)

C. Thousand Islands, Java (Ref. 823)




Fig. 1. (A) Extreme value plot for spotted sardinella, *Amblygaster sirm*, in Indonesia based on data from *R/V Dr. Fridtjof Nansen*, showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 22.7 \pm 2.2$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 4 length-frequency samples of *Amblygaster sirm* from northern Borneo based on data from *R/V Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 2). Open dot represents an outlier not used for analysis.

[Gambar 1. (A) Gambaran nilai ekstrim untuk ikan sardin, *Amblygaster sirm*, di Indonesia berdasarkan data dari kapal penelitian Dr. Fridtjof Nansen, yang menunjukkan nilai maksimum dari 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 22.7 \pm 2.2$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 4 contoh frekuensi panjang ikan sardin, *Amblygaster sirm*, dari Kalimantan berdasarkan data Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 2). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]




Fig. 2. Distribution of spotted sardinella, *Amblygaster sirm*, in Western Indonesia based on records of the surveys of *R/V Dr. Fridtjof Nansen*. [Gambar 2. Penyebaran ikan sardin, *Amblygaster sirm*, di Indonesia bagian barat berdasarkan laporan survei kapal penelitian Dr. Fridtjof Nansen.]




Fig. 3. Depth distribution of spotted sardinella, *Amblygaster sirm*, in Western Indonesia based on surveys of R/V Dr. Fridtjof Nansen.

[Gambar 3. Penyebaran kedalaman ikan sardin, *Amblygaster sirm*, di Indonesia bagian barat berdasarkan survei kapal penelitian Dr. Fridtjof Nansen.]

Dussumieria acuta (Valenciennes, 1847)

Rainbow sardine (English); Djapuh (Indonesian); Ajapu, Djapuh (West Java, Jakarta); Tjapo (Madura); Tembang djawa (South Sulawesi, Makassar); Tembang rakapeng (South Sulawesi, Bugis); Bete kalo (South Sulawesi, Badjo).

Brachiostegal rays fewer (12 to 15) and posterior part of scales marked with numerous tiny radiating striae. Color is iridescent blue with a shiny gold/brass line below (quickly fading after death). W-shaped pelvic scute; isthmus tapering evenly forward; more anal finrays. Dorsal spines: 0-0; soft rays: -; anal spines: 0-0; soft rays: 14-18. $L_{max1} = 20$ cm SL; $L_{max2} = n.a.$; $L_{max3} = 20.9$ cm TL (Fig. 4A). See Fig. 4B and Table 4 for length-weight relationship.

Warmer waters of the Indo-Pacific, from the Persian Gulf (and perhaps south to Somalia), along the coasts of Pakistan, India and Malaysia to Indonesia (Fig. 5) and the Philippines.

Earlier records included *D. elopsoides*.

Mainly an inshore species. Depth range: 10-120 m (Fig. 6). Earlier studies on the habitat and biology may have equally referred to *D. elopsoides* which this species closely resembles.

References: 171, 188, 280, 1449, 2178, 2857, 2860, 4789, 5193, 5381, 5525, 5541, 5579, 5730, 5736, 5756, 6313, 6328, 6365

Table 4. Length-weight (g/[TL;cm]) relationship of rainbow sardine, *Dussumieria acuta*, in Indonesia.
[Tabel 4. Hubungan panjang-berat [g/(TL; cm)] ikan japuh, *Dussumieria acuta*, di Indonesia.]

Parameter	Estimate
a	0.0056
s.e.(a)	0.00402
b	3.1462
s.e.(b)	0.25642
r^2	0.9692




Fig. 4. (A) Extreme value plot for rainbow sardine, *Dussumieria acuta*, in Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen, showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 20.9 \pm 1.5$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 10 length-frequency samples of rainbow sardine, *Dussumieria acuta*, from Western Indonesia based on data from R/Vs Dr. Fridtjof Nansen, Mutiara 4 and Jurong as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 4).
 [Gambar 4. (A) Gambaran nilai ekstrim dari ikan japuh, *Dussumieria acuta*, di Indonesia berdasarkan data kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen, yang menunjukkan nilai maksimum dari 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 20.9 \pm 1.5$ cm TL. (B) Berat prediksi terhadap berat-berat observasi (dalam g berat basah) dari 10 contoh frekuensi-panjang ikan japuh, *Dussumieria acuta*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4 dan Jurong sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 4).]




Fig. 5. Distribution of rainbow sardine, *Dussumieria acuta*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Mutiara 4, Jurong and Bawal Putih 2.
 [Gambar 5. Penyebaran ikan japuh, *Dussumieria acuta*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4, Jurong dan Bawal Putih 2.]




Fig. 6. Depth distribution of rainbow sardine, *Dussumieria acuta*, in Western Indonesia based on records of the surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Mutiara 4 and (C) Jurong.

[Gambar 6. Penyebaran kedalaman ikan japuh, *Dussumieria acuta*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian: (A) Dr. Fridtjof Nansen, (B) Mutiara 4, dan (C) Jurong.]

Pellona ditchela (Valenciennes, 1847)

Indian pellona (English); Dero (Indonesian); Dero, Longlong mata (Java); Puput (West Java, Jakarta).

Belly with usually 18 or 19 + 8 or 9, total 26 to 28 scutes, strongly keeled. Eye large, lower jaw projecting. Dorsal fin origin near midpoint of body. Scales with upper and lower vertical striae slightly overlapping each other at center of scales. Dorsal spines: 0-0; soft rays: 0-0; anal spines: 0-0; soft rays: 34-42. $L_{max1} = 16$ cm SL; $L_{max2} = n.a.$; $L_{max3} = 17.7$ cm TL (Fig. 7A). See Fig. 7B and Table 5 for length-weight relationship.

Indian Ocean: Madagascar, and from Durban, South Africa to the Gulf of Oman and the coasts of India. From the Andaman Sea to Indonesia (Fig. 8) and the Philippines; southeast to Papua New Guinea and Northern and Western Australia.

Occurs in coastal areas, entering mangrove swamps and penetrating estuaries and freshwater. Depth range: 10-55 m (Fig. 9).

References: 171, 188, 1455, 2857, 3225, 3509, 4749, 4789, 4959, 4967, 5193, 5213, 5284, 5339, 6313, 6365, 6567, 6822

Table 5. Length-weight (g/[TL;cm]) relationship of Indian pellona, *Pellona ditchela*, in Indonesia. [Tabel 5. Hubungan panjang-berat (g/[TL; cm]) ikan puput, *Pellona ditchela*, di Indonesia.]

Parameter	Estimate
a	0.0018
s.e.(a)	0.00357
b	3.6209
s.e.(b)	0.62929
r ²	0.9988




Fig. 7. (A) Extreme value plot for Indian pellona, *Pellona ditchela*, in Indonesia based on data from R/V Dr. Fridtjof Nansen showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 17.7 \pm 0.6$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 4 length-frequency samples of Indian pellona, *Pellona ditchela*, from northern Borneo based on data from R/V Dr. Fridtjof Nansen as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 5).

[Gambar 7. (A) Gambaran nilai ekstrim dari ikan puput, *Pellona ditchela*, di Indonesia berdasarkan data dari kapal penelitian Dr. Fridtjof Nansen yang menunjukkan 4 contoh frekuensi panjang dan angka perkiraan $L_{max3} = 17.7 \pm 0.6$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 4 contoh frekuensi panjang ikan puput, *Pellona ditchela*, dari Kalimantan Utara berdasarkan data dari kapal penelitian Dr. Fridtjof Nansen]




Fig. 8. Distribution of Indian pellona, *Pellona ditchela*, in Western Indonesia based on records of the surveys of R/V Dr. Fridtjof Nansen.
 [Gambar 8. Penyebaran ikan puput, *Pellona ditchela*, di perairan Indonesia bagian barat berdasarkan laporan survei dari kapal penelitian Dr. Fridtjof Nansen.]




Fig. 9. Depth distribution of Indian pellona, *Pellona ditchela*, in Western Indonesia based on surveys of R/V Dr. Fridtjof Nansen.
 [Gambar 9. Penyebaran kedalaman ikan puput, *Pellona ditchela*, di Indonesia bagian barat berdasarkan survei dengan kapal penelitian Dr. Fridtjof Nansen.]

Sardinella gibbosa (Bleeker, 1849)

Goldstriped sardinella (English); Tembang (Indonesian); Djuwi djuwi, Mursiah, Tjiro (Java); Tamban sisik, Tembang, Tembang djuwi (West Java, Jakarta); Maos, Tandjan (West Java, Bandung); Sintring (Madura); Djurung (East Sumatra); Tamban (South Borneo); Tembang lakara (South Sulawesi, Bugis); Totata (South Sulawesi, Badjo); Mengida (Bali).

Total number of scutes: 32 to 34. A golden mid-lateral line down flank; dorsal and caudal fin margins dusky; a dark spot at dorsal fin origin. Lower gillrakers: 45 to 59 (at 6 to 17 cm SL, not increasing with size of fish after 6 cm SL). Dorsal spines: 0-0; soft rays: 13-21; anal spines: 0-0 soft rays: 12-23. $L_{max1} = 17$ cm SL. $L_{max2} = n.a.$; $L_{max3} = 20.2$ cm TL (Fig. 10A). See Fig. 10B and Table 6 for length-weight relationship.

Indo-West Pacific: from the East African coasts (but not the Red Sea) and Madagascar eastward to the Persian Gulf and Indonesia (Fig. 11), north to Taiwan and Korea; south to Northern Australia. In India, often confused with *S. fimbriata*.

Forms schools in coastal waters. Depth range: 10-70 m (Fig. 12). Feeds on phytoplankton and zooplankton (crustacean and molluscan larvae). Table 7 presents a set of growth parameters from Indonesia.

References: 171, 188, 280, 811, 1314, 1443, 1444, 1488, 1504, 1529, 1751, 2178, 2857, 2948, 3560, 3605, 4330, 4331, 5213, 5284, 5730, 5736, 5756, 6313, 6365

Table 6. Length-weight (g/[TL;cm]) relationship of goldstriped sardinella, *Sardinella gibbosa*, in Indonesia.

[Tabel 6. Hubungan panjang-berat (g/[TL; cm]) dari ikan tembang, *Sardinella gibbosa*, di Indonesia.]

Parameter	Estimate
a	0.0158
s.e.(a)	0.0136
b	2.7837
s.e.(b)	0.3073
r^2	0.9778

Table 7. Growth parameters of goldstriped sardinella, *Sardinella gibbosa*.

[Tabel 7. Parameter pertumbuhan ikan tembang, *Sardinella gibbosa*.]

Parameter	A
L_{∞} (TL, cm)	19.5
K (year ⁻¹)	1.20

A. Riau waters (Ref. 1314)




Fig. 10. (A) Extreme value plot for goldstriped sardinella, *Sardinella gibbosa*, in Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen, showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 20.2 \pm 0.6$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 9 length-frequency samples of goldstriped sardinella, *Sardinella gibbosa*, from Western Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 6).

[Gambar 10. (A) Gambaran nilai ekstrim dari ikan tembang, *Sardinella gibbosa*, di Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen, yang menunjukkan nilai maksimum 6 contoh frekuensi panjang dan nilai perkiraan maksimum $L_{max3} = 20.2 \pm 0.6$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 9 contoh frekuensi panjang ikan tembang, *Sardinella gibbosa*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1) dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 6).




Fig. 11. Distribution of goldstriped sardinella, *Sardinella gibbosa*, in Western Indonesia based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.
 [Gambar 11. Penyebaran ikan tembang, *Sardinella gibbosa*, di Indonesia bagian barat berdasarkan laporan survei dari kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 12. Depth distribution of goldstriped sardinella, *Sardinella gibbosa*, in Western Indonesia based on surveys of R/V Dr. Fridtjof Nansen.
 [Gambar 12. Penyebaran kedalaman ikan tembang, *Sardinella gibbosa*, di Indonesia bagian barat berdasarkan survei kapal penelitian Dr. Fridtjof Nansen.]

Sardinella lemuru (Bleeker, 1853)

Bali sardinella (English); Tembang montjong (Indonesian); Lemuru (Java); Lemuru, Tembang mata kutjing, Tembang montjong (West Java, Jakarta); Soroi (Madura); Temban montjo (South Sulawesi, Makassar); Bete laki (South Sulawesi, Bugis).

A faint golden spot behind gill opening, followed by a faint golden mid-lateral line; a distinct black spot at hind border of gill cover. Body elongate, subcylindrical. Distinguished from all other clupeids in the eastern Indian Ocean and western Pacific by its pelvic finray count of 8; from *S. longiceps* by its shorter head length and fewer lower gillrakers. Dorsal spines: 0-0; soft rays: 13-21; anal spines: 0-0; soft rays: 12-23. $L_{max1} = 23$ cm SL. $L_{max2} = 21$ cm TL; $L_{max3} = 19.9$ cm TL (Fig. 13A). See Fig. 13B and Table 8 for length-weight relationship.

Eastern Indian Ocean: Phuket, Thailand and southern coasts of East Java and Bali (Fig. 14); also in Western Australia, Western Pacific: Java Sea north to the Philippines, Hong Kong, Taiwan Island to southern Japan.

Forms large schools in coastal waters. Depth range: 15-100 m (Fig. 15). Feeds on phytoplankton and zooplankton (chiefly copepods). Ghofar and Mathews (Box 2) discuss the fluctuations of the Bali Straits lemuru fishery from 1976 to 1993. Table 9 presents six sets of growth parameters from Indonesia.

References: 171, 188, 280, 818 819, 1263, 1314, 1392, 1449, 1511, 1830, 2178, 2858, 3268, 3557, 3605, 3784, 5381.

Table 8. Length-weight (g/[TL;cm]) relationship of Bali sardinella, *Sardinella lemuru*, in Indonesia.

[Tabel 8. Hubungan panjang-berat (g/[TL;cm]) dari ikan lemuru, *Sardinella lemuru*, di Indonesia.]

Parameter	A	B
a	0.0012	0.0299
s.e.(a)	0.0012	n.a.
b	3.7515	2.671
s.e.(b)	0.3087	n.a.
r^2	0.9641	n.a.

A. This study

B. Bali Strait (Ref. 3268), L in SL.

Table 9. Growth parameters of Bali sardinella, *Sardinella lemuru*.

[Tabel 9. Parameter pertumbuhan dari ikan lemuru, *Sardinella lemuru*.]

Parameter	A	B	C	D	E	F
L_{∞} (TL, cm)	21.6	21.1	22.3	22.5	23.2	23.8
K (year ⁻¹)	0.95	0.8	0.85	1.0	1.28	0.505

A. Bali Strait (Ref. 3268), L originally in SL.

B. Bali Strait, 1981 (Ref. 1314)

C. Bali Strait, 1977 (Ref. 1314)

D. Bali Strait, 1980 (Ref. 1314)

E. Bali Strait, 1979 (Ref. 1314)

F. Muntjar, Bali Strait (Ref. 819)




Fig. 13. (A) Extreme value plot for Bali sardinella, *Sardinella lemuru*, in Indonesia based on data from *R/V Jurong*, showing maxima of 7 length-frequency samples, and estimate of $L_{max3} = 19.9 \pm 0.5$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 12 length-frequency samples of Bali sardinella, *Sardinella lemuru*, from Western Indonesia based on data from *R/V Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 8). Open dots represent outliers, not used for analysis.

[Gambar 13. (A) Gambaran nilai ekstrim untuk ikan lemuru, *Sardinella lemuru*, di Indonesia berdasarkan data dari kapal penelitian *Jurong*, yang menunjukkan nilai maksimum dari 7 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 19.9 \pm 0.5$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 12 contoh frekuensi-panjang ikan lemuru, *Sardinella lemuru*, dari Indonesia bagian barat berdasarkan data dari kapal penelitian *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1) dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 8). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]

Box 2. The Bali Straits lemuru fishery.
 [Boks 2. Perikanan lemuru Selat Bali]

The Bali Straits lemuru (*Sardinella lemuru*) fishery relied until 1975 mainly on small (<5 GT) sail powered boats with a range of up to about 20 nm offshore. In the early fishery four kinds of hand operated gear were used: *payang* (a non-closing surface seine net); *jala buang* (throw net); *serok* (dip net); and *bagan cancap* (lift net), which attract lemuru with lights at night to a bamboo platform: the fish are caught by lifting a large suspended net. Starting around the early 1970s, the ca. 9 m boats were motorized, and by the mid-1970s, the older gear was being replaced by (often larger) mechanized purse seiners (*pukat cincin*) which are now the dominant type, with over 85% of the catch; this is taken mainly from October to April, with a strong peak in December and January.

Two data sets on catch and effort are available: 1950-1973 and 1976-1993 (Fig. 1). Both series cover the same stock and area, with catches overwhelmingly (<90%) of lemuru, but have been gathered using different methodologies. Still, increasing mechanization, and natural variations similar to those observed in earlier and later years could account for the jump in landings from ca. 5,000 in 1973 to about 28,000 t·year⁻¹ in 1976 recorded at the end of the first, and the beginning of the second, respectively, of the data sets.

Catches show an overall increase, but with marked fluctuations (Fig. 1). Data for 1994 are not yet available but the decline that started in 1993 appears to have continued. Effort was low prior to 1976, but quantitative data are not available. Observed effort changes (Fig. 1) cannot account for the low landings in 1977-1980 and 1985-1987.

To test if lemuru landings may be related to the El Niño/Southern Oscillation (ENSO) events, the catch series were converted into a single series of standardized anomalies (A) by fitting a five-year running mean to the data from 1950 to 1993. We defined $A = (C_o/C_m) - 1$, where C_o is the observed catch in a given year, and C_m is the running mean for the same year.

As might be seen from Fig. 1 (insert), high positive anomalies values are clearly related to ENSO events. Therefore, the increased effort directed towards the Bali Straits lemuru stock from 1950 to 1993, which produced an increase in landings for <5,000 t·year⁻¹ to >50,000 t·year⁻¹ is not the primary cause for the strong catch fluctuations which are probably due to large-scale oceanographic events (see also Sharp, this vol.).

Nevertheless, there is a suggestion that higher effort in the last decades had some effect on the stock: the amplitude of the positive anomalies declines from >1.5 to <0.5 over the time series (Fig. 1, insert), while the amplitude of the negative anomalies remains roughly constant over the whole time series. Perhaps the higher catches in recent years were taken from a stock that has become less resilient, with the fishery removing most of the surplus production. This would prevent the spawning stock from recovering during favorable periods ("E"), making it more vulnerable to fishing during less favorable periods ("N"). This suggests that further effort increases, while not increasing catches, will increase the risk of a collapse. Also, the quick recovery observed from 1986 to 1988 may not recur at higher effort levels. This issue needs further work; to support this, we have contributed our time series of catch and effort data to the database (Diskette 2) described by Torres et al. (this vol.).




Fig. 1. Major features of the Bali Straits lemuru fishery.
 [Gambar 1. Sifat-sifat utama perikanan lemuru Selat Bali.]
 Main panel: catches of 1950 to 1973 (from Ref. 3268); catch and effort for 1976-1993: original data.
 Insert: catch anomalies (1950 to 1993) and ENSO events ("E") and their opposite ("N") from Refs. 9577, 9578 and 9580.

A. Ghofar
 Department of
 Fisheries
 Diponegoro
 University
 Semarang -
 Indonesia
 and
C.P. Mathews
 Marine
 Resources Evaluation
 and Planning Project
 (MCESP)
 Central
 Research Institute for
 Fisheries, P.O. Box 50
 Slipi, Jakarta
 11401A, Indonesia




Fig. 14. Distribution of Bali sardinella, *Sardinella lemuru*, in Western Indonesia based on records of the surveys of R/Vs Jurong and Bawal Putih 2.

[Gambar 14. Penyebaran ikan lemuru, *Sardinella lemuru*, di Indonesia bagian barat berdasarkan laporan survei dari kapal-kapal penelitian Jurong dan Bawal Putih 2.]




Fig. 15. Depth distribution of Bali sardinella, *Sardinella lemuru*, in Western Indonesia based on surveys of R/V Jurong.

[Gambar 15. Penyebaran kedalaman ikan lemuru, *Sardinella lemuru*, di Indonesia bagian barat berdasarkan survei kapal penelitian Jurong.]

***Netuma thalassina* (Rüppell, 1837)**

Giant catfish (English); Manjung (Indonesian); Mangmung, Manjong (Java); Manjung kerbo (West Java); Duri padi, Manjung (West Java, Jakarta); Duri utek, Utik (West Java, Jakarta); Gaguk, Putih (South Sumatra); Duri padi, Duri utek (Riouw); Gugup, Gungut (West Borneo); Barukang (South Sulawesi, Makassar); Lampa (South Sulawesi, Badjo).

Head shield weakly striated and granulated, its surface nearly smooth. Three pairs of barbels around mouth. Supraoccipital process about 1.5 times as long as broad. Dorsal spines: 1-1; soft rays: 7-7; anal spines: - ; soft rays: 16-30. $L_{max1} = 100$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 83$ cm TL (Fig. 16A). See Fig. 16B and Table 10 for length-weight relationship.

Indian Ocean: known with certainty only from the Red Sea and northwest Indian Ocean. Malaysia, Indonesia (Fig. 17) and southeast to north Australia.

A marine species often found in estuaries, but rarely enters freshwater; depth range 10-195 m (Fig. 18). Feeds mainly on crabs, prawns, mantis shrimps (*Squilla* spp.) but also on fishes and molluscs. Table 11 presents two sets of growth parameters from Indonesia.

References: 1263, 1314, 2045, 2857, 2872, 3279, 3290, 3627, 3641, 4515, 4557, 4600, 4735, 4789, 4883, 4959, 5213, 5450, 5736, 5756, 6313, 6365, 6567

Table 10. Length-weight (g/[TL;cm]) relationship of giant catfish, *Netuma thalassina*, in Indonesia. [Tabel 10. Hubungan panjang-berat (g/[TL;cm]) ikan manyung, *Netuma thalassina*, di Indonesia.]

Parameter	Estimate
a	0.0097
s.e.(a)	0.0032
b	3.0404
s.e.(b)	0.0886
r ²	0.9787

Table 11. Growth parameters of giant catfish, *Netuma thalassina*. [Tabel 11. Parameter pertumbuhan ikan manyung, *Netuma thalassina*.]

Parameter	A	B
L_{∞} (TL, cm)	52.7	60.0
K (year ⁻¹)	0.27	0.65

A. Sampit Bay, Central Kalimantan (Ref. 4557)
 B. Java Sea (South Kalimantan) (Ref. 1314)




Fig. 16. (A) Extreme value plot for giant catfish, *Netuma thalassina*, in Indonesia based on data from R/Vs Mutiara 4 and Jurong, showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 83.0 \pm 11.0$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 26 length-frequency samples of giant catfish, *Netuma thalassina*, from Western Indonesia based on data from R/Vs Mutiara 4 and Jurong as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 10). Open dots represent outliers, not used for analysis.

[Gambar 16. (A) Gambaran nilai ekstrim ikan manyung, *Netuma thalassina*, di Indonesia berdasarkan data dari kapal-kapal penelitian Mutiara 4 dan Jurong menunjukkan nilai maksimum dari 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 83.0 \pm 11.0$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 26 contoh frekuensi-panjang ikan manyung, *Netuma thalassina*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4 dan Jurong sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 10). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 17. Distribution of giant catfish, *Netuma thalassina*, in Western Indonesia based on records of the surveys of *R/Vs* Mutiara 4, Bawal Putih 2, Dr. Fridtjof Nansen, and Jurong.
 [Gambar 17. Penyebaran ikan manyung, *Netuma thalassina*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Dr. Fridtjof Nansen dan Jurong.]




Fig. 18. Depth distribution of giant catfish, *Netuma thalassina*, in Western Indonesia based on surveys of *R/Vs* (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong and (D) Bawal Putih 2.
 [Gambar 18. Penyebaran kedalaman ikan manyung, *Netuma thalassina*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong dan (D) Bawal Putih 2.]

Saurida micropectoralis (Shindo & Yamada, 1972)

Shortfin lizardfish (English); Beloso sirip pendek (Indonesian).

Body elongate, cylindrical. Back and upper sides brown, lower sides and belly white. Nine to ten faint blotches along the lateral line, sometimes with traces of very indistinct cross-bars on the back. Dorsal spines: -; soft rays: -; anal spines: -; soft rays: -. $L_{max1} = 38$ cm; $L_{max2} = n.a.$; $L_{max3} = 49.7$ cm FL (Fig. 19A). See Fig. 19B and Table 12 for length-weight relationship.

Indo-West Pacific: Andaman and South China Sea, Indonesia (Fig. 20); south to northern Australia.

Occurs over muddy bottoms from 20 to 260 m (Fig. 21). Feeds on small bottom-dwelling invertebrates and fishes. Table 13 presents a set of growth parameters from Indonesia.

References: 393, 1314, 2117, 2857, 4749, 4789, 5756

Table 12. Length-weight (g/[FL;cm]) relationship of shortfin lizardfish, *Saurida micropectoralis*, in Indonesia.

[Tabel 12. Hubungan panjang-berat [g/(FL;cm)] ikan beloso sirip pendek, *Saurida micropectoralis*, di Indonesia.]

Parameter	Estimate
a	0.0050
s.e.(a)	0.0008
b	3.1959
s.e.(b)	0.0530
r ²	0.9988

Table 13. Growth parameters of shortfin lizardfish, *Saurida micropectoralis*.

[Tabel 13. Parameter pertumbuhan dari ikan beloso sirip pendek, *Saurida micropectoralis*.]

Parameter	A
L_{∞} (TL, cm)	42.0
K (year ⁻¹)	0.88

A. Java Sea (Central Java) (Ref. 1314)




Fig. 19. (A) Extreme value plot for shortfin lizardfish, *Saurida micropectoralis*, in Indonesia based on data from R/Vs *Mutiara 4* and *Jurong*, showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 49.7 \pm 12.8$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 7 length-frequency samples of shortfin lizardfish, *Saurida micropectoralis*, from Western Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 12). Open dot represents outlier, not used for analysis.

[Gambar 19. (A) Gambaran nilai ekstrim untuk ikan beloso sirip pendek, *Saurida micropectoralis*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong*, menunjukkan nilai maksimum dari 5 contoh frekuensi-panjang, dan nilai perkiraan $L_{max3} = 49.7 \pm 12.8$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 7 contoh frekuensi-panjang ikan beloso sirip pendek, *Saurida micropectoralis*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1), dan memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 12). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]




Fig. 20. Distribution of shortfin lizardfish, *Saurida micropectoralis*, in Western Indonesia based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2 and Jurong.

[Gambar 20. Penyebaran ikan beloso sirip pendek, *Saurida micropectoralis*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2 dan Jurong.]




Fig. 21. Depth distribution of shortfin lizardfish, *Saurida micropectoralis*, in Western Indonesia based on surveys of R/Vs (A) Mutiara 4, (B) Jurong and (C) Bawal Putih 2.

[Gambar 21. Penyebaran kedalaman dari ikan beloso sirip pendek, *Saurida micropectoralis*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Mutiara 4, (B) Jurong dan (C) Bawal Putih 2.]

Saurida undosquamis (Richardson, 1848)

Brushtooth lizardfish (English); Beloso (Indonesian).

Cigar-shaped, rounded or slightly compressed; the head pointed and depressed; the snout rounded. Color is brown-gray above and creamy below, with 8-10 indistinct darker spots along the middle of the sides. Dorsal spines: 0-0; soft rays: 11-12; anal spines: 0-0; soft rays: 11-12. $L_{max1} = 50$ cm. SL; $L_{max2} =$ n.a.; $L_{max3} = 41.45$ cm TL (Fig. 22A). See Fig. 22B and Table 14 for length-weight relationship.

Indo-West Pacific from South Africa, through Indonesia (Fig. 23) to Japan and Western Australia (Great Barrier Reef). Migrated from the Red Sea through the Suez Canal to the eastern Mediterranean.

Found over muddy substrates of coastal waters, from about 20-290 m (Fig. 24). Feeds on fishes, crustaceans, and other invertebrates. Table 15 presents a set of growth parameters from Indonesia.

References: 231, 312, 1139, 1263, 1288, 1289, 1314, 1449, 1474, 1486, 1488, 1498, 1524, 1532, 2178, 2857, 2877, 3397, 3557, 3626, 3670, 3674, 3675, 3676, 3678, 4055, 4595, 4789, 4964, 5193 5213, 5284, 5337, 5381, 5385, 5450, 5525, 5736, 5756, 5760, 5829, 6313, 6328, 6365, 6567

Table 14. Length-weight (g/[TL;cm]) relationship of brushtooth lizardfish, *Saurida undosquamis*, in Indonesia.

[Tabel 14. Hubungan panjang-berat (g/[TL;cm]) ikan beloso, *Saurida undosquamis*, di Indonesia.]

Parameter	Estimate
a	0.0027
s.e.(a)	0.0017
b	3.3200
s.e.(b)	0.1918
r ²	0.9601

Table 15. Growth parameters of brushtooth lizardfish, *Saurida undosquamis*.

[Tabel 15. Parameter pertumbuhan ikan beloso, *Saurida undosquamis*.]

Parameter	A
L_{∞} (TL, cm)	33.5
K (year ⁻¹)	0.95

A. Java Sea (Central Java) (Ref. 1314)




Fig. 22. (A) Extreme value plot for brushtooth lizardfish, *Saurida undosquamis*, in Indonesia based on data from R/Vs *Mutiara 4* and *Jurong*, showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 41.45 \pm 5.92$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 18 length-frequency samples of brushtooth lizardfish, *Saurida undosquamis*, from Western Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 14). Open dots represent outliers, not used for analysis. [Gambar 22. (A) Gambaran nilai ekstrim untuk ikan beloso, *Saurida undosquamis*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* menunjukkan nilai maksimum 6 contoh frekuensi-panjang, dan nilai perkiraan $L_{max3} = 41.45 \pm 5.92$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 18 contoh frekuensi-panjang ikan beloso, *Saurida undosquamis*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 14). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 23. Distribution of brushtooth lizardfish, *Saurida undosquamis*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Mutiara 4, Jurong and Bawal Putih 2.
 [Gambar 23. Penyebaran ikan beloso, *Saurida undosquamis*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4, Jurong dan Bawal Putih 2.]




Fig. 24. Depth distribution of brushtooth lizardfish, *Saurida undosquamis*, in Western Indonesia based on surveys of R/V Jurong.
 [Gambar 24. Penyebaran kedalaman ikan beloso, *Saurida undosquamis*, di Indonesia bagian barat berdasarkan survei kapal penelitian Jurong.]

Carangoides malabaricus (Bloch & Schneider, 1801)

Malabar trevally (English); Karang trevali, Kuwe (Indonesian).

Silvery, bluish gray dorsally. Opercle with a small, black spot. Lateral line with 19-36 weak scutes. Pectoral fins falcate; first dorsal lobe slightly falcate. No scales on breast to behind pelvic origin and laterally to pectoral base, including the small area anteriorly just above fin. Dorsal spines: 9 - 9; soft rays: 20-23; anal spines: 3-3; soft rays: 17-19. L_{max1} = 60 cm; L_{max2} = n.a.; L_{max3} = 29.18 cm TL (Fig. 25A). See Fig. 25B and Table 16 for length-weight relationship.

Ranges from the east coast of Africa (without verified records from the Red Sea) to Sri Lanka and farther eastward to the Gulf of Thailand and Indonesia (Fig. 26), north to Okinawa

(Japan) and south to Australia.

Found near rocks and coral reefs. Depth range: 20-110 m (Fig. 27). Juveniles inhabit sandy bays. Feeds on crustaceans, small squids and fish.

References: 280, 1449, 2334, 2857, 3280, 3287, 3605, 5213, 5450, 5736, 5756, 6313, 6365, 6567

Table 16. Length-weight [g/(TL;cm)] relationship of Malabar trevally, *Carangoides malabaricus*, in Indonesia. [Tabel 16. Hubungan panjang-berat [g/(TL;cm)] ikan karang trevali, *Carangoides malabaricus*, di Indonesia.]

Parameter	Estimate
a	0.0205
s.e.(a)	0.0120
b	2.8476
s.e.(b)	0.1953
r ²	0.9796




Fig. 25. (A) Extreme value plot for Malabar trevally, *Carangoides malabaricus*, in Indonesia based on data from R/Vs Dr. Fridtjof Nansen and Jurong, showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 29.18 \pm 0.945$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 10 length-frequency samples of Malabar trevally, *Carangoides malabaricus*, from Western Indonesia based on data from R/Vs Mutiara 4, Dr. Fridtjof Nansen and Jurong as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 16). Open dot represents outlier, not used for analysis.

[Gambar 25. (A) Gambaran nilai ekstrim dari ikan karang trevali, *Carangoides malabaricus*, di Indonesia berdasarkan data dari kapal-kapal penelitian Dr. Fridtjof Nansen dan Jurong, yang menunjukkan angka maksimum dari 5 contoh frekuensi-panjang, dan nilai perkiraan $L_{max3} = 29.18 \pm 0.945$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 10 contoh frekuensi-panjang ikan karang trevali, *Carangoides malabaricus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Dr. Fridtjof Nansen dan Jurong sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 16). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]




Fig. 26. Distribution of Malabar trevally, *Carangoides malabaricus*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Mutiara 4, Jurong and Bawal Putih 2.
 [Gambar 26. Penyebaran ikan karang trevali, *Carangoides malabaricus*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4, Jurong dan Bawal Putih 2.]




Fig. 27. Depth distribution of Malabar trevally, *Carangoides malabaricus*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Jurong and (C) Bawal Putih 2.
 [Gambar 27. Penyebaran kedalaman ikan karang trevali, *Carangoides malabaricus*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) [Fridtjof Nansen, (B) Jurong dan (C) Bawal Putih 2.]

Caranx ignobilis (Forsskål, 1775)

Giant trevally (English); Karang besar (Indonesian).

Head and body dusky golden dorsally, silver ventrally; fins usually pigmented gray to black. Opercular spot absent. Twenty-six to 38 strong scutes. Breast scaleless ventrally; a small patch of prepelvic scales. Pectoral fins falcate; anal fin with 2 detached spines. Dorsal spines: 9-9; soft rays: 17-22; anal spines: 3-3; soft rays: 15-17. $L_{max1} = 165$ cm FL; $L_{max2} =$ n.a.; $L_{max3} = 57.2$ cm FL (Fig. 28A). See Fig. 28B and Table 17 for length-weight relationship.

Widely distributed throughout most of the Indian Ocean, the Indonesian Archipelago (Fig. 29) and the Central Pacific, eastward to the Hawaiian and Marquesas Islands.

Juveniles are found in small schools over sandy inshore bottoms, adults usually solitary, over the reef. Depth range 20-100 m (Fig. 30). Usually feeds at night on fishes and

crustaceans such as crabs and spiny lobsters. Large individuals may be ciguatoxic.

References: 171, 583, 1602, 2334, 2857, 2872, 3280, 3287, 3605, 3626, 3678, 3804, 3807, 4332, 4362, 4390, 4560, 4699, 4735, 4795, 4821, 4887, 4917, 4959, 5213, 5450, 5525, 5736, 5756, 5970, 6026, 6057, 6273, 6306, 6313, 6365

Table 17. Length-weight (g/[FL;cm]) relationship of giant trevally, *Caranx ignobilis*, in Indonesia.

[Tabel 17. Hubungan panjang-berat (g/[FL;cm]) ikan karang besar, *Caranx ignobilis*, di Indonesia.]

Parameter	Estimate
a	0.0202
s.e.(a)	n.a.
b	3.0000
s.e.(b)	n.a.
r^2	0.0000




Fig. 28. (A) Extreme value plot for giant trevally, *Caranx ignobilis*, in Indonesia based on data from *R/V Jurong*, showing maxima of 2 length-frequency samples, and estimate of $L_{max3} = 57.2 \pm 8.2$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 2 length-frequency samples of giant trevally, *Caranx ignobilis*, from Western Indonesia based on data from *R/V Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 17).

[Gambar 28. (A) Penggambaran nilai ekstrim ikan karang besar, *Caranx ignobilis*, di Indonesia berdasarkan data dari kapal penelitian *Jurong*, yang menunjukkan nilai maksimum dari 2 contoh frekuensi-panjang, dan nilai perkiraan $L_{max3} = 57.2 \pm 8.2$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 2 contoh frekuensi-panjang ikan karang besar, *Caranx ignobilis*, dari Indonesia bagian barat berdasarkan data kapal penelitian *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 17).




Fig. 29. Distribution of giant trevally, *Caranx ignobilis*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2.
 [Gambar 29. Penyebaran ikan karang besar, *Caranx ignobilis*, di Indonesia bagian barat berdasarkan laporan dari survei kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2.]




Fig. 30. Depth distribution of giant trevally, *Caranx ignobilis*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) Jurong.
 [Gambar 30. Penyebaran kedalaman ikan karang besar, *Caranx ignobilis*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) Jurong.]

Caranx tille (Cuvier, 1833)

Tille trevally (English); Karang tile (Indonesian).

Body dark olive green to bluish gray dorsally, silvery white below; soft dorsal lobe olive gray to blackish. Upper part of opercle with a small blackish spot. Thirty-three to 42 strong scutes. Pectoral fins falcate. Two anal fin spines detached. Breast fully scaled. Dorsal spines: 9-9; soft rays: 20-22; anal spines: 3-3; soft rays: 16-18. $L_{max1} = 80$ cm; $L_{max2} = n.a.$; $L_{max3} = 54.5$ cm FL (Fig. 31A). See Fig. 31B and Table 18 for length-weight relationship.

Distribution in the Indian Ocean not well established; reported from Durban to Zanzibar; also recorded in Madagascar and Sri Lanka. Ranges from Indonesia (Fig. 32) to southern Japan (Okinawa), Australia and Fiji.

Inhabits coastal waters, near coral reefs and rocks. Depth range: 30 to 120 m (Fig. 33). Feeds on fish and crustaceans.

References: 171, 2334, 2857, 3197, 3280, 3807, 5193, 5213

Table 18. Length-weight (g/[FL;cm]) relationship of tille trevally, *Caranx tille*, in Indonesia.

Tabel 18. Hubungan panjang-berat (g/[FL;cm]) dari ikan karang tile, *Caranx tille*, di Indonesia.

Parameter	Estimate
a	0.0088
s.e.(a)	0.0092
b	3.1630
s.e.(b)	0.2859
r^2	0.9928




Fig. 31. (A) Extreme value plot for tille trevally, *Caranx tille*, in Indonesia based on data from *R/V Jurong*, showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 54.5 \pm 17.6$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 4 length-frequency samples of tille trevally, *Caranx tille*, from Western Indonesia based on data from *R/V Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 18).

[Gambar 31. (A) Penggambaran nilai ekstrim ikan karang tile, *Caranx tille*, di Indonesia berdasarkan data dari kapal penelitian Jurong, menunjukkan nilai maksimum dari 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 54.5 \pm 17.6$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 4 contoh frekuensi-panjang ikan karang tile, *Caranx tille*, dari Indonesia bagian barat berdasarkan data kapal penelitian Jurong sebagai output perangkat lunak ABee (lihat Boks 1), yang memungkinkan estimasi hubungan panjang-berat (lihat Tabel 18).




Fig. 32. Distribution of tille trevally, *Caranx tille*, in Western Indonesia based on records of the surveys of R/Vs *Jurong* and *Bawal Putih 2*.
 [Gambar 32. Penyebaran ikan karang tile, *Caranx tille*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian *Jurong* dan *Bawal Putih 2*.]




Fig. 33. Depth distribution of tille trevally, *Caranx tille*, in Western Indonesia based on surveys of R/Vs (A) *Jurong*, and (B) *Bawal Putih 2*.
 [Gambar 33. Penyebaran kedalaman ikan karang tile, *Caranx tille*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) *Jurong* dan (B) *Bawal Putih 2*.]

Decapterus macrosoma (Bleeker, 1851)

Shortfin scad (English); Lajang deles (Indonesian); Bengol deles, Deles, Lajang, Lajang deles, Lajang lidi, Luntju (Java); Lajang (West Java, Jakarta); Bulus blangseng, Kaban bulus, Kaban laes, Kaban padara, Kaban patek, Ladjeng lakek, Rentjek bulus, Rentjek kaban (Madura); Bulus (Bawean).

Metallic blue dorsally, silvery ventrally; fins hyaline. Opercle with a small black spot. Anal fin with 2 detached spines. Twenty-four to 40 scutes. Upper jaw reaching below front margin of eye. Dorsal spines: 9-9; soft rays: 33-38; anal spines: 3-3; soft rays: 27-30. $L_{max1} = 35$ cm; $L_{max2} = 20$ cm; $L_{max3} = 28.95$ cm TL (Fig. 34A). See Fig. 34B and Table 19 for length-weight relationship.

Pacific Ocean: from southern Japan to warm waters of the Western Pacific, including the Indonesian Archipelago (Fig. 35). Eastern Pacific: from the Gulf of California, Mexico to Peru, including the Galapagos Islands.

Forms schools. Depth range: 20-140 m (Fig. 36). Feeds on small invertebrate plankton. Table 20 presents six sets of growth parameters from Indonesia.

References: 171, 312, 559, 761, 1263, 1314, 1386, 1392, 1447, 1449, 1462, 1467, 1602, 2021, 2023, 2334, 2857, 3280, 3287,

3555, 3556, 3786, 3804, 3807, 4536, 4789, 4838, 5213, 5337, 5340, 5530, 5730, 5756, 6313, 6365

Table 19. Length-weight (g/[TL;cm]) relationship of shortfin scad, *Decapterus macrosoma*, in Indonesia. [Tabel 19. Hubungan panjang-berat (g/[TL;cm]) ikan layang deles, *Decapterus macrosoma*, di Indonesia.]

Parameter	A	B
a	0.0076	0.009
s.e.(a)	0.0125	n.a.
b	3.0051	3.01
s.e.(b)	0.5630	n.a.
r ²	0.8669	n.a.

A. This study.
B. Java Sea (Ref. 1386).

Table 20. Growth parameters of shortfin scad, *Decapterus macrosoma*. [Tabel 20. Parameter pertumbuhan ikan layang deles, *Decapterus macrosoma*.]

Parameter	A	B	C	D	E	F
L_{∞} (cm)	24.0	24.0	25.4	25.6	25.7	27.7
K (year ⁻¹)	1.15	1.00	0.98	1.05	0.90	1.20

A. Java Sea (Ref. 1386), L in FL.
B. Asahan, Sumatra (Ref. 1467).
C. Java Sea (Pekalongan) (Ref. 1314), L in TL.
D. Java Sea (Ref. 1447), L in TL.
E. Langsa, Sumatra (Ref. 1467).
F. Banda Aceh, Sumatra (Ref. 1467).




Fig. 34. (A) Extreme value plot for shortfin scad, *Decapterus macrosoma*, in Indonesia based on data from R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2, showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 28.95 \pm 5.24$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 8 length-frequency samples of shortfin scad, *Decapterus macrosoma*, from Western Indonesia based on data from R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2 as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 19). Open dot represents outlier, not used for analysis.

[Gambar 34. (A) Gambaran nilai ekstrim ikan layang deles, *Decapterus macrosoma*, di Indonesia berdasarkan data dari kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2, menunjukkan nilai maksimum dari 6 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 28.95 \pm 5.24$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 8 contoh frekuensi-panjang ikan layang deles, *Decapterus macrosoma*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2 sebagai output perangkat lunak ABee (lihat Box 1), dan yang memungkinkan suatu hubungan panjang-berat (lihat Tabel 19). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]




Fig. 35. Distribution of shortfin scad, *Decapterus macrosoma*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2.

[Gambar 35. Penyebaran ikan layang deles, *Decapterus macrosoma*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2.]




Fig. 36. Depth distribution of shortfin scad, *Decapterus macrosoma*, in Western Indonesia based on surveys of R/V Jurong.

[Gambar 36. Penyebaran kedalaman ikan layang deles, *Decapterus macrosoma*, di Indonesia bagian barat berdasarkan survei kapal penelitian Jurong.]

***Decapterus russelli* (Rüppell, 1830)**

Indian scad (English); Lajang (Indonesian); Bengol, Korok, Ladjeng, Lajang (Java); Lajang (West Java, Jakarta); Kaban padara, Kaban patek, Ladjang (Madura); Rentjek bulus, Rentjek kaban, Rentjek padara, Rentjek patek (Madura).

Lateral line curved below soft dorsal and with 30-44 strong scutes; bluish green above, silvery below; caudal fin hyaline to yellowish; dorsal fins hyaline basally, light dusky distally. Opercle with small, black spot; opercular membrane with smooth margin. Snout longer than eye diameter; squarish lower posterior edge of maxilla; upper jaw with small teeth anteriorly; soft dorsal and anal fins relatively low, not falcate; pectoral fin subfalcate. Dorsal spines: 9-9; soft rays: 28-31; anal spines: 3-3; soft rays: 25-28. $L_{max1} = 35$ cm FL; $L_{max2} = n.a.$; $L_{max3} = n.a.$ See Table 21 for length-weight relationship.

From East Africa via Southeast Asia and the Indonesian Archipelago (Fig. 37) to Japan and Australia (and possibly to New Caledonia).

Schooling in coastal waters and on open banks. Depth range: 40-275 m (Fig. 38). Feeds mainly on smaller planktonic invertebrates. Table 22 presents five sets of growth parameters from Indonesia.

References: 171, 312, 559, 761, 1263, 1314, 1384, 1385, 1454, 1455, 1632, 2021, 2334, 3131, 3197, 3287, 3555, 3556,

3807, 4537, 4546, 4591, 4838, 4883, 4931, 5213, 5284, 5337, 5339, 5406, 5417, 5418, 5432, 5433, 5434, 5440, 5441, 5443, 5444, 5446, 5525, 5730, 5736, 5756, 5885, 5970, 6026, 6365

Table 21. Length-weight (g/[TL; cm]) relationship of Indian scad, *Decapterus russelli*, in Indonesia.

[Tabel 21. Hubungan panjang-berat (g/[TL;cm]) ikan layang, *Decapterus russelli*, di Indonesia.]

Parameter	Estimates	
	A	B
a	0.0112	0.0104
b	2.970	3.000
r ²	n.a.	0.980

A. Tegal (Ref. 5441), Length type unspecified.
B. Java Sea (Ref. 1385).

Table 22. Growth parameters of Indian scad, *Decapterus russelli*.

[Tabel 22. Parameter pertumbuhan ikan layang, *Decapterus russelli*.]

Parameter	A	B	C	D	E
L_{∞} (cm)	26.0	26.6	27.0	27.0	28.4
K (year ⁻¹)	0.90	0.95	1.15	1.18	0.90

A. Idi, Malacca Strait (Ref. 5432), L in FL.
B. Java Sea (Seribu Island) (Ref. 1314), L in TL.
C. Jakarta Bay (Seribu Island), L in TL, 1973 (Ref. 1314).
D. Jakarta Bay (Seribu Island), L in TL, 1975 (Ref. 1314).
E. Java Sea (Ref. 1385), L in FL.




Fig. 37. Distribution of Indian scad, *Decapterus russelli*, in Western Indonesia based on records of the surveys of R/Vs *Jurong* and *Bawal Putih 2*.

[Gambar 37. Penyebaran ikan layang, *Decapterus russelli*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian *Jurong* dan *Bawal Putih 2*.]




Fig. 38. Depth distribution of Indian scad, *Decapterus russelli*, in Western Indonesia, based on surveys of *R/V Jurong*.

[Gambar 38. Penyebaran kedalaman ikan layang, *Decapterus russelli*, di Indonesia bagian barat berdasarkan survei kapal penelitian *Jurong*.]

***Parastromateus niger* (Bloch, 1795)**

Black pomfret (English); Bawal hitam (Indonesian); Gebel (Java); Bawal, Bawal hitam, Dorang, Dorang hitam (West Java, Jakarta); Dibas, Kandibas, Kapet, Kibas, Tjeplek (Madura); Bawal hitam (East Sumatra); Manriwasa lelung (South Sulawesi, Makasar); Peda-peda lotong (South Sulawesi, Bugis).

Deep-bodied and strongly compressed. Lateral line ends in weakly-developed scutes on the caudal peduncle. Pelvic fins lost in individuals over 9 cm. Color is brown above, silvery-white below. The anterior parts of the dorsal and anal fins bluish-gray, other fins yellowish. Dorsal spines: 2-6; soft rays: 41-46; anal spines: 2-2; soft rays: 35-40. $L_{max1} = 75$ cm; $L_{max2} = n.a.$; $L_{max3} = 38.4$ cm TL (Fig. 39A). See Fig. 39B and Table 23 for length-weight relationship.

From East Africa through the Indonesian Archipelago (Fig. 40) to southern Japan and Australia.

Forms large schools in coastal areas with muddy substrate. Depth range: 20-105 m (Fig. 41); near the bottom during daytime and near the water surface at night. Table 24 presents a set of growth parameters from Indonesia.

References: 171, 1314, 2334, 3287, 4789, 5213, 5284, 5736, 5756, 6365, 6567

Table 23. Length-weight (g/[TL;cm]) relationship of black pomfret, *Parastromateus niger*, in Indonesia. [Tabel 23. Hubungan panjang-berat [g/(TL;cm)] ikan bawal hitam, *Parastromateus niger*, di Indonesia.]

Parameter	Estimate
a	0.0073
s.e.(a)	0.0063
b	3.3189
s.e.(b)	0.2676
r ²	0.8901

Table 24. Growth parameters of black pomfret, *Parastromateus niger*. [Tabel 24. Parameter pertumbuhan ikan bawal hitam, *Parastromateus niger*.]

Parameter	A
L_{∞} (TL, cm)	29.5
K (year ⁻¹)	0.68

A. Java Sea (Central Java) (Ref. 1314)




Fig. 39. (A) Extreme value plot for black pomfret, *Parastromateus niger*, in Indonesia based on data from *R/Vs Mutiara 4*, *Dr. Fridtjof Nansen*, *Bawal Putih 2* and *Jurong*, showing maxima of 8 length-frequency samples, and estimate of $L_{max3} = 38.4 \pm 3.3$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 26 length-frequency samples of black pomfret, *Parastromateus niger*, from Western Indonesia based on data from *R/Vs Mutiara 4*, *Dr. Fridtjof Nansen*, *Bawal Putih 2* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 23). Open dots represent outliers, not used for analysis.

[Gambar 39. (A) Gambaran nilai ekstrim untuk ikan bawal hitam, *Parastromateus niger*, di Indonesia berdasarkan data dari kapal-kapal penelitian Mutiara 4, Dr. Fridtjof Nansen, Bawal Putih 2 dan Jurong, menunjukkan nilai maksimum untuk 8 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 38.4 \pm 3.3$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 26 contoh frekuensi-panjang ikan bawal hitam, *Parastromateus niger*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian Mutiara 4, Dr. Fridtjof Nansen, Bawal Putih 2 dan Jurong sebagai output perangkat lunak ABee (lihat Boks 1) yang memungkinkan estimasi hubungan panjang-berat (lihat Tabel 23). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 40. Distribution of black pomfret, *Parastromateus niger*, in Western Indonesia based on records of the surveys of *R/Vs Dr. Fridtjof Nansen*, *Mutiara 4*, *Jurong* and *Bawal Putih 2*.

[Gambar 40. Penyebaran ikan bawal hitam, *Parastromateus niger*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4, Jurong dan Bawal Putih 2.]




Fig. 41. Depth distribution of black pomfret, *Parastromateus niger*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) Jurong.

[Gambar 41. Penyebaran kedalaman ikan bawal hitam, *Parastromateus niger*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) Jurong.]

Selar crumenophthalmus (Bloch, 1793)

Bigeye scad (English); Bentong (Indonesian); Penteng, Pentong, Selar bentong (West Java, Jakarta); Bun bun, Tong gentong (Madura); Gintong (Central Sumatra).

Metallic blue to bluish green dorsally, shading to white ventrally; the lateral yellow stripe sometimes present. Lower margin of gill opening with a deep furrow, a large papilla immediately above it and a smaller one near upper edge. Operculum with black spot. Straight part of lateral line with 0-11 scales and 29-42 scutes. First two anal spines detached; pectoral fins falcate. Dorsal spines: 9-9; soft rays: 24-27; anal spines: 3-3; soft rays: 21-23. $L_{max1} = 60$ cm SL; $L_{max2} = n.a.$; $L_{max3} = 26.7$ cm FL (Fig. 42A). See Fig. 42B and Table 25 for length-weight relationship.

Circumtropical; Indo-Pacific: from southern Africa to Indonesia (Fig. 43); northeast to southern Japan and the Hawaiian Islands; south to New Caledonia and Rapa; east to Mexico to Peru and the Galapagos Islands, Western Atlantic: through the West Indies.

Forms small to large compact schools in inshore water and shallow reefs; mainly nocturnal; younger stages feed inshore on small shrimp and benthic invertebrates (including foraminiferans). The adults feed further offshore on zooplankton and fish larva, and range in depth from 10 to 170 m (Fig. 44). Table 26 presents two sets of growth parameters from Indonesia.

References: 171, 276, 1263, 1314, 1447, 1602, 2178, 2300, 2325, 2334, 2857, 3084, 3277, 3605, 3786, 3804, 3807, 4390, 4789, 4795, 4821, 4838, 4839, 4887, 4905, 5213, 5217, 5284, 5288, 5337, 5450, 5525, 5530, 5730, 5736, 5756, 5970, 6026, 6273, 6306, 6313, 6315, 6365, 6567, 6810

Table 25. Length-weight (g/[FL;cm]) relationship of bigeye scad, *Selar crumenophthalmus*, in Indonesia.

[Tabel 25. Hubungan panjang-berat (g/[FL;cm]) ikan selar bentong, *Selar crumenophthalmus*, di Indonesia.]

Parameter	Estimate
a	0.0176
s.e.(a)	0.0109
b	3.0039
s.e.(b)	0.2102
r^2	0.9737

Table 26. Growth parameters of bigeye scad, *Selar crumenophthalmus*.

[Tabel 26. Parameter pertumbuhan ikan selar bentong, *Selar crumenophthalmus*.]

Parameter	A	B
L_{∞} (TL; cm)	25.9	26.9
K (year ⁻¹)	1.25	1.35

A. Java Sea (Pekalongan) (Ref. 1386)
B. Java Sea (Ref. 1447)




Fig. 42. (A) Extreme value plot for bigeye scad, *Selar crumenophthalmus*, in Indonesia based on data from R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2, showing maxima of 7 length-frequency samples, and estimate of $L_{max3} = 26.7 \pm 0.86$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 28 length-frequency samples of bigeye scad, *Selar crumenophthalmus*, from Western Indonesia based on data from R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2 as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 25).

[Gambar 42. (A) Gambaran nilai ekstrim dari ikan selar bentong, *Selar crumenophthalmus*, di Indonesia berdasarkan data dari kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2 menunjukkan nilai maksimum dari 7 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 26.7 \pm 0.86$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 28 contoh frekuensi-panjang dari ikan selar bentong, *Selar crumenophthalmus*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2 sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 25).




Fig. 43. Distribution of bigeye scad, *Selar crumenophthalmus*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Jurong and Bawal Putih 2.

[Gambar 43. Penyebaran ikan selar bentong, *Selar crumenophthalmus*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2.]




Fig. 44. Depth distribution of bigeye scad, *Selar crumenophthalmus*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Jurong and (C) Bawal Putih 2.
 [Gambar 44. Penyebaran kedalaman ikan selar bentong, *Selar crumenophthalmus*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Jurong dan (C) Bawal Putih 2.]

***Pentaprion longimanus* (Cantor, 1850)**

Longfin mojarra (English); Lontong (Indonesian); Lontjong (Java); Hajam (West Java, Jakarta).

Body is slender, with weakly attached silvery scales. The spines of the dorsal and anal fins longer than the rays; the pectoral fins long and pointed, reaching beyond the anal fin spines; the anal fin is long; the caudal fin lobes rounded. Dorsal spines: 9-10; soft rays: 14-15; anal spines: 5-6; soft rays: 12-13. $L_{max1} = 15$ cm; $L_{max2} = n.a.$; $L_{max3} = 15.5$ cm TL (Fig. 45A). See Fig. 45B and Table 27 for length-weight relationship.

Indian Ocean: from the western and southern coasts of India and off Sri Lanka to Indonesia. Western Pacific: Indonesia (Fig. 46) to the Philippines and the Ryukyu Islands, and south to the northern part of Australia.

Forms large schools in coastal waters. Depth range: 20-220 m (Fig. 47). Feeds on small benthic animals. Table 28 presents six sets of growth parameters from Indonesia.

References: 393, 559, 1263, 1314, 1381, 1392, 1435, 1449, 1452, 1486, 1966, 2029, 2178, 2857, 2872, 2926, 3131, 3399, 3409, 3807, 4672, 4749, 5381, 5756, 6365, 6567

Table 27. Length-weight (g/[TL;cm]) relationship of longfin mojarra, *Pentaprion longimanus*, in Indonesia. [Tabel 27. Hubungan panjang-berat (g/[TL;cm]) ikan loncong, *Pentaprion longimanus*, di Indonesia.]

Parameter	Estimate
a	0.0169
s.e.(a)	0.0080
b	2.9173
s.e.(b)	0.1949
r ²	0.9725

Table 28. Growth parameters of longfin mojarra, *Pentaprion longimanus*. [Tabel 28. Parameter pertumbuhan ikan loncong, *Pentaprion longimanus*.]

Parameter	A	B	C	D	E	F
L_{∞} (cm)	13.4	13.5	13.7	14.2	15.6	15.6
K (year ⁻¹)	1.77	1.10	1.12	1.80	0.80	0.94

- A. Java Sea (Ref. 1452)
- B. Java Sea (Semarang) (Ref. 1314), L in TL
- C. Java Sea (Ref. 1452)
- D. Java Sea (Ref. 1452)
- E. Java Sea (Ref. 1452)
- F. Java Sea (southern) (Ref. 1381), L in TL




Fig. 45. (A) Extreme value plot for longfin mojarra, *Pentaptrion longimanus*, in Indonesia based on data from R/Vs Mutiara 4 and Dr. Fridtjof Nansen showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 15.5 \pm 0.54$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 39 length-frequency samples of longfin mojarra, *Pentaptrion longimanus*, from Western Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 27). Open dots represent outliers, not used for analysis.

[Gambar 45. (A) Gambaran nilai ekstrim dari ikan loncong, *Pentaptrion longimanus*, di Indonesia berdasarkan data kapal-kapal penelitian Mutiara 4 dan Dr. Fridtjof Nansen menunjukkan nilai maksimum dari 6 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 15.5 \pm 0.54$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 39 contoh frekuensi-panjang dari ikan loncong, *Pentaptrion longimanus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 27). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 46. Distribution of longfin mojarra, *Pentaptrion longimanus*, in Western Indonesia based on records of the surveys of R/Vs Dr. Fridtjof Nansen, Mutiara 4, Jurong and Bawal Putih 2.

[Gambar 46. Penyebaran ikan loncong, *Pentaptrion longimanus*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, Mutiara 4, Jurong dan Bawal Putih 2.]




Fig. 47. Depth distribution of longfin mojarra, *Pentaptrion longimanus*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong and (D) Bawal Putih 2.
 [Gambar 47. Penyebaran kedalaman ikan loncong, *Pentaptrion longimanus*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong dan (D) Bawal Putih 2.]

Diagramma pictum (Thunberg, 1792)

Painted sweetlips (English); Gadjih (Indonesian); Katji (Java); Katji-katji (Java); Domul (West Java, Jakarta); Gadji (Java, Jakarta); Gadji-gadji (Java, Jakarta); Ikan kadji (West Java, Jakarta); Kadji (West Java, Jakarta); Besiko (Riouw); Domul (Riouw); Radja bau (Ceram, Wahai, Ambon, Luhu, Saparua, Haria, Geser).

Body typically perciform. Flesh in maxilla thick. Usually cardiform jaw teeth. Vomer generally toothless. Usually with enlarged chin pore. Branchiostegal rays: 7. Dorsal spines: 9-10; soft rays: 21-26; anal spines: 3-3; soft rays: 7-8. $L_{max1} = 100$ cm FL; $L_{max2} = n.a.$; $L_{max3} = 84.2$ cm TL (Fig. 48A). See Fig. 48B and Table 29 for length-weight relationship.

Indo-West Pacific: East Africa and Red Sea, Indonesia to New Caledonia, north to Japan.

Occurs in shallow coastal areas and coral reefs. Depth range: 20-170 m. (Fig. 50). Juveniles usually occur in seaweed

beds, and large adults in small schools or solitary around coral. Carnivore; feeds on benthic invertebrates and fishes.

References: 280, 559, 1498, 1602, 1830, 2112, 2290, 2334, 2682, 2799, 2871, 2872, 3111, 3131, 3412, 3626, 3670, 3678, 4517, 5213, 5450, 5525, 5736, 5970, 5978, 6026, 6065, 6066, 6067, 6068, 6365, 6567, 6956, 9070, 9137

Table 29. Length-weight (g/[TL;cm]) relationship of painted sweetlips, *Diagramma pictum*, in Indonesia.
 [Tabel 29. Hubungan panjang-berat (g/[TL;cm]) ikan kaji, *Diagramma pictum*, di Indonesia.]

Parameter	Estimate
a	0.0077
s.e.(a)	0.0058
b	3.1314
s.e.(b)	0.1783
r ²	0.9867




Fig. 48. (A) Extreme value plot for painted sweetlips, *Diagramma pictum*, in Indonesia based on data from *R/Vs Mutiara 4* and *Jurong* showing maxima of 7 length-frequency samples, and estimate of $L_{max3} = 84.2 \pm 3.6$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 21 length-frequency samples of painted sweetlips, *Diagramma pictum*, from Western Indonesia based on data from *R/Vs Mutiara 4*, *Jurong* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 29). Open dots represent outliers, not used for analysis. [Gambar 48. (A) Penggambaran nilai ekstrim untuk ikan kaji, *Diagramma pictum*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* menunjukkan nilai maksimum dari 7 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 84.2 \pm 3.6$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 21 contoh frekuensi panjang ikan kaji, *Diagramma pictum*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Bawal Putih 2* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 29). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 49. Distribution of painted sweetlips, *Diagramma pictum*, based on records of the surveys of *R/Vs Dr. Fridtjof Nansen*, *Mutiara 4*, *Jurong* and *Bawal Putih 2*. [Gambar 49. Penyebaran ikan kaji, *Diagramma pictum*, berdasarkan laporan survei kapal-kapal penelitian Dr. Fridtjof Nansen, *Mutiara 4*, *Jurong* dan *Bawal Putih 2*.]




Fig. 50. Depth distribution of painted sweetlips, *Diagramma pictum*, based on surveys of R/Vs (A) *Jurong* and (B) *Bawal Putih 2*.
 [Gambar 50. Penyebaran kedalaman ikan kaji, *Diagramma pictum*, berdasarkan survei kapal-kapal penelitian (A) *Jurong* dan (B) *Bawal Putih 2*.]

***Pomadasys argenteus* (Forsskål, 1775)**

Silver grunt (English); Da-tanda (Indonesian); Krokot (Java); Gerot-gerot, Kerot-kerot, Krot, Krot-krot (West Java, Jakarta); Da-tanda, Mengantih, Towoito (Madura); Ronga (South Sulawesi, Makassar); Garut (West Borneo).

Body ovate; head profile almost straight. Mouth small; lips not thickened; two pores and a central groove under the chin. No antrorse spine before the dorsal fin origin; notch between the spinous and soft rayed portion of the dorsal fin shallow. Color is generally silver-mauve to fawn above, white below. Small specimens with numerous spots aligned horizontally or fused into horizontal lines; large specimens plain or with scattered charcoal scale spots on back and upper sides; the snout is dark brown; the upper operculum charcoal or purplish. Dorsal spines: 12-12; soft rays: 13-14; anal spines: 3-3; soft rays: 7-7. $L_{max1} = 66$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 60.4$ cm TL (Fig. 51A). See Fig. 51B and Table 30 for length-weight relationship.

From the Red Sea to Indonesia (Fig. 52) and the Philippines (but without record from the Persian Gulf) and southern to northern Australia. Also reported from New Caledonia.

Found in coastal waters. Depth range: 15-115 m (Fig. 53). Mainly carnivore, feeds on benthic animals. Table 31 presents a set of growth parameters from Indonesia.

References: 312, 1115, 1116, 1139, 1314, 3412, 3624, 3627, 3642, 3670, 3678, 4606, 4959, 5284, 5450, 5525, 5736, 5756, 6026, 6365

Table 30. Length-weight (g/[TL;cm]) relationship of silver grunt, *Pomadasys argenteus*, in Indonesia.
 [Tabel 30. Hubungan panjang-berat (g/[TL;cm]) ikan da-tanda, *Pomadasys argenteus*, di Indonesia.]

Parameter	Estimate
a	0.0267
s.e.(a)	0.0157
b	2.8551
s.e.(b)	0.1545
r ²	0.9758

Table 31. Growth parameter of silver grunt, *Pomadasys argenteus*.
 [Tabel 31. Parameter pertumbuhan ikan da-tanda, *Pomadasys argenteus*.]

Parameter	A
L _∞ (TL; cm)	54
K (year ⁻¹)	0.5

A. Java Sea (Tanjung Selatan, South Kalimantan) (Ref. 1314)




Fig. 51. (A) Extreme value plot for silver grunt, *Pomadasys argenteus*, in Indonesia based on data from *R/V Jurong* showing maxima of 8 length-frequency samples, and estimate of $L_{max3} = 60.4 \pm 2.2$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 42 length-frequency samples of silver grunt, *Pomadasys argenteus*, from Western Indonesia based on data from *R/Vs Mutiara 4, Jurong* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 30). Open dots represent outliers, not used for analysis.

[Gambar 51. (A) Gambaran nilai ekstrim ikan da-tanda, *Pomadasys argenteus*, di Indonesia berdasarkan data dari kapal penelitian *Jurong* menunjukkan nilai maksimum dari 8 contoh frekuensi-panjang, dan nilai perkiraan $L_{max3} = 60.4 \pm 2.2$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 42 contoh frekuensi-panjang ikan da tanda, *Pomadasys argenteus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4, Jurong* dan *Bawal Putih 2* sebagai output perangkat lunak ABee (lihat Boks 1), yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 30). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 52. Distribution of silver grunt, *Pomadasys argenteus*, in Western Indonesia based on records of the surveys of *R/Vs Mutiara 4, Jurong* and *Bawal Putih 2*.

[Gambar 52. Penyebaran ikan da-tanda, *Pomadasys argenteus*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4, Jurong* dan *Bawal Putih 2*.]




Fig. 53. Depth distribution of silver grunt, *Pomadasys argenteus*, in Western Indonesia based on surveys of R/Vs (A) *Jurong* and (B) *Bawal Putih 2*. [Gambar 53. Penyebaran kedalaman ikan da-tanda, *Pomadasys argenteus*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) *Jurong* dan (B) *Bawal Putih 2*.]

Pomadasys maculatus (Bloch, 1797)

Saddle grunt (English); Gerot-gerot (Indonesian); Gerot-gerot, Ikan krot, Kerot-kerot, Krot-krot (West Java, Jakarta).

Small-sized fish of moderately deep body. Isthmus narrow, forming a groove. Chin with two pairs of small pores. This species is characterized by several dark large elongate blotches on the upper back, one forming a saddle on the nape. Dorsal spines: 12-12; soft rays: 13-14; anal spines: 3-3; soft rays: 7-7. $L_{max1} = 59.3$ cm FL; $L_{max2} = n.a.$; $L_{max3} = 57.5$ cm FL (Fig. 54A). See Fig. 54B and Table 32 for length-weight relationship.

Indo-West Pacific: from the east coast of Africa and Madagascar, to Southeast Asia, via the Indonesian Archipelago (Fig. 55) and, thence northeast to China and southeast to Australia.

Found in coastal waters over sand near reefs; depth range: 20-110 m (Fig. 56). Feeds on crustaceans and fish.

References: 280, 393, 1498, 2112, 2135, 2857, 2871, 2872, 3225, 3626, 4749, 5213, 5450, 5736, 5756, 6567

Table 32. Length-weight (g/[FL;cm]) relationship of saddle grunt, *Pomadasys maculatus*, in Indonesia. [Tabel 32. Hubungan panjang-berat (g/[FL;cm]) dari ikan gerot-gerot, *Pomadasys maculatus*, di Indonesia.]

Parameter	Estimate
a	0.0799
s.e.(a)	0.0307
b	2.5693
s.e.(b)	0.1085
r ²	0.9856




Fig. 54. (A) Extreme value plot for saddle grunt, *Pomadasys maculatus*, in Indonesia based on data from R/Vs *Jurong* and *Bawal Putih 2* showing maxima of 9 length-frequency samples, and estimate of $L_{max3} = 57.5 \pm 1.5$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 13 length-frequency samples of silver grunt, *Pomadasys maculatus*, from Western Indonesia based on data from R/Vs *Mutiara 4*, Dr. Fridtjof Nansen, *Jurong* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 32). Open dots represent outliers, not used for analysis.

[Gambar 54. (A) Gambaran nilai ekstrim ikan gerot-gerot, *Pomadasys maculatus*, di Indonesia berdasarkan data kapal-kapal penelitian *Jurong* dan *Bawal Putih 2* menunjukkan nilai maksimum dari 9 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 57.5 \pm 1.5$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 13 contoh frekuensi-panjang ikan gerot-gerot, *Pomadasys maculatus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, Dr. Fridtjof Nansen, *Jurong* dan *Bawal Putih 2* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 32). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 55. Distribution of saddle grunt, *Pomadasys maculatus*, in Western Indonesia based on records of the surveys of R/Vs Mutiara 4, Dr. Fridtjof Nansen, Jurong and Bawal Putih 2.

[Gambar 55. Penyebaran ikan gerot-gerot, *Pomadasys maculatus*, di Indonesia bagian barat berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Dr. Fridtjof Nansen, Jurong dan Bawal Putih 2.]




Fig. 56. Depth distribution of saddle grunt, *Pomadasys maculatus*, in Western Indonesia based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Jurong and (C) Bawal Putih 2.

[Gambar 56. Penyebaran kedalaman ikan gerot-gerot, *Pomadasys maculatus*, di Indonesia bagian barat berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Jurong dan (C) Bawal Putih 2.]

Lactarius lactarius (Bloch & Schneider, 1801)

False trevally (English); Ikan susu (Indonesian); Lemahan, Limat, Tana (Java); Ikan lemah, Lelemah, Lemah, Susu (West Java, Jakarta); Klemes (Madura); Tambi-tambi (South Borneo); Bebete lubangang (South Sulawesi, Badjo).

Silvery gray with blue iridescence dorsally, silvery white ventrally; upper part of gill cover with a dusky black spot; fins pale yellow. Mouth large and oblique. Dorsal spines: 8-9; soft rays: 20-22; anal spines: 3-3; soft rays: 25-28. $L_{max1} = 40$ cm; $L_{max2} = n.a.$; $L_{max3} = 29.1$ cm TL (Fig. 57A). See Fig. 57B and Table 33 for length-weight relationship.

From the eastern Indian Ocean to Southeast Asia, extending northward to Japan, and southeastward through the

Indonesian Archipelago (Fig. 58) to Queensland, Australia.

Occurs in coastal waters; depth range: 15-90 m (Fig. 59)

Feeds on sand-dwelling and other benthic and zooplanktonic animals.

References: 312, 1012, 2857, 2872, 3404, 3423, 4789, 4931, 5193, 5213, 5736, 5756, 5978, 6313, 6567, 7300

Table 33. Length-weight (g/[TL;cm]) relationship of false trevally, *Lactarius lactarius*, in Indonesia.

Tabel 33. Hubungan panjang-berat (g/[TL;cm]) ikan susu, *Lactarius lactarius*, di Indonesia.

Parameter	Estimate
a	0.0098
s.e.(a)	0.0034
b	3.0469
s.e.(b)	0.1237
r ²	0.9942




Fig. 57. (A) Extreme value plot for false trevally, *Lactarius lactarius*, in Indonesia based on data from *R/Vs Mutiara 4, Jurong* and *Dr. Fridtjof Nansen* showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 29.1 \pm 4.97$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 8 length-frequency samples of false trevally, *Lactarius lactarius*, from Western Indonesia based on data from *R/Vs Mutiara 4, Jurong* and *Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 33).

[Gambar 57. (A) Gambaran nilai ekstrim untuk ikan susu, *Lactarius lactarius*, di Indonesia berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen menunjukkan nilai maksimum untuk 6 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 29.1 \pm 4.97$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 8 contoh frekuensi-panjang dari ikan susu, *Lactarius lactarius*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 33).

***Leiognathus splendens* (Cuvier, 1829)**

Splendid ponyfish (English); Bondol (Indonesian); Dodok, Gempar, Gemper (Java); Peperek Tjina (West Java, Jakarta); Bondol (West Java, Bandung).

Belly silvery; back grayish silvery with faint, gray wavy vertical lines above lateral lines in adults; lateral line scales and pectoral fin base yellow; pectoral axis black. Scales small and deciduous. Nuchal spine with a distinct median keel. Chest fully scaled. Mouth horizontal, pointing slightly downward when protracted; line of closed mouth passing below eye; a narrow brown band around end of snout. Lower edge of operculum and margin of supraorbital serrated. Third and fourth dorsal and third anal spines anteriorly serrated. Dorsal spines: 7-8; soft rays: 15-17; anal spines: 3-3; soft rays: 13-14. $L_{max1} = 17$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 21.1$ cm TL (Fig. 60A). See Fig. 60B and Table 34 for length-weight relationship.

Indian Ocean: Madagascar and Mauritius to the Red Sea, along the coasts of India and Sri Lanka; Indonesian Archipelago (Fig. 61) and throughout Western Central Pacific, reaching westward to Australia and Fiji.

This schooling species inhabits coastal waters. Depth range: 10-100 m (Fig. 62). Feeds on small fish, crustaceans, foraminiferans, and bivalves. Table 35 presents four sets of growth parameters from Indonesia.

References: 312, 393, 559, 560, 573, 1139, 1263, 1314, 1449, 1486, 1539, 1617, 1633, 1724, 1830, 1918, 2045, 2089, 2108,

2178, 2462, 2504, 2505, 2682, 3131, 3151, 3424, 3430, 3436, 3437, 3438, 3439, 3440, 3441, 3442, 3443, 3444, 3605, 3607, 3614, 3649, 3653, 3655, 3667, 4544, 4880, 4961, 4962, 4963, 5213, 5346, 5381, 5450, 5525, 5736, 5756, 5978, 6192, 6313, 6567, 6992, 7050, 7100

Table 34. Length-weight (g/[TL;cm]) relationship of splendid ponyfish, *Leiognathus splendens*, in Indonesia.

[Tabel 34. Hubungan panjang-berat (g/[TL;cm]) ikan bondol, *Leiognathus splendens*, di Indonesia.]

Parameter	Estimates	
	A	B
a	0.0112	0.0168
s.e.(a)	n.a.	0.0104
b	3.2170	3.0392
s.e.(b)	n.a.	0.2612
r ²	n.a.	0.9707

A. Northwestern coast of Java (Ref. 2089)
B. This study

Table 35. Growth parameters of splendid ponyfish, *Leiognathus splendens*. [Tabel 35. Parameter pertumbuhan ikan bondol, *Leiognathus splendens*.]

Parameter	Indonesia			
	A	B	C	D
L_{∞} (TL, cm)	14	14.5	16.7	16.9
K (year ⁻¹)	1.04	1.25	0.90	1.10

A. Southern Kalimantan (Ref. 1139)
B. Riau (Bintan) (Ref. 1314)
C. Java Sea (Central Java) (Ref. 1314)
D. Java Sea (South Kalimantan) (Ref. 1314)




Fig. 60. (A) Extreme value plot for splendid ponyfish, *Leiognathus splendens*, in Indonesia based on data from R/Vs Mutiara 4, Dr. Fridtjof Nansen and Jurong showing maxima of 7 length-frequency samples, and estimate of $L_{max3} = 21.1 \pm 3.35$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 19 length-frequency samples of splendid ponyfish, *Leiognathus splendens*, from Western Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 34). Open dots represent outliers, not used for analysis.

[Gambar 60. (A) Gambaran nilai ekstrim ikan bondol, *Leiognathus splendens*, di Indonesia berdasarkan data survei kapal-kapal penelitian Mutiara 4, Dr. Fridtjof Nansen dan Jurong yang menunjukkan 7 contoh frekuensi-panjang dan angka perkiraan $L_{max3} = 21.1 \pm 3.35$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 19 contoh frekuensi-panjang ikan bondol, *Leiognathus splendens*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 34). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 61. Distribution of splendid ponyfish, *Leiognathus splendidus*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.

[Gambar 61. Penyebaran ikan bondol, *Leiognathus splendidus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 62. Depth distribution of splendid ponyfish, *Leiognathus splendidus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong and (D) Bawal Putih 2.

[Gambar 62. Penyebaran kedalaman ikan bondol, *Leiognathus splendidus*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong dan (D) Bawal Putih 2.]

Leiognathus bindus (Valenciennes, 1835)

Orangefin ponyfish (English); Tjaria (Indonesian); Petah (Java); Peperek (West Java, Jakarta); Tjaria (South Sulawesi, Bugis).

Silvery body; snout with a dark band; dorsal and anal fins with orange tips. Head naked; with nuchal spine. Mouth pointing forward when protracted. Breast with small scales. Dorsal spines: 8-8; soft rays: 16-16; anal spines: 3-3; soft rays: 14-14. $L_{max1} = 14$ cm; $L_{max2} = n.a.$; $L_{max3} = 14.8$ cm TL (Fig. 63A). See Fig. 63B and Table 36 for length-weight relationship.

Indian Ocean: Red Sea (Port Sudan), Persian Gulf, India, Sri Lanka, Bangladesh. Western Central Pacific, including Indonesia (Fig. 64) and Australia; also reported from New Caledonia.

Found in shallow waters. Depth range: 10-100 m (Fig. 65). Forms schools. Table 37 presents a set of growth parameters from Indonesia.

References: 312, 393, 1015, 1016, 1263, 1314, 1372, 1403, 1449, 1486, 2044, 2088, 2108, 2857, 3424, 3605, 4789, 5346, 5381, 5525, 5756, 6365, 6567

Table 36. Length-weight (g/[TL;cm]) relationship of orangefin ponyfish, *Leiognathus bindus*, in Indonesia.

[Tabel 36. Hubungan panjang-berat (g/[TL;cm]) ikan caria, *Leiognathus bindus*, di Indonesia.]

Parameter	Estimate
a	0.0182
s.e.(a)	0.0044
b	2.9191
s.e.(b)	0.1210
r ²	0.9902

Table 37. Growth parameters of orangefin ponyfish, *Leiognathus bindus*.

[Tabel 37. Parameter pertumbuhan ikan caria, *Leiognathus bindus*.]

Parameter	A
L_{∞} (TL, cm)	12.5
K (year ⁻¹)	1.38

A. Java Sea (Central Java) (Ref. 1314)




Fig. 63. (A) Extreme value plot for orangefin ponyfish, *Leiognathus bindus*, in Indonesia based on data from R/V *Jurong* showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 14.8 \pm 1.15$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 17 length-frequency samples of orangefin ponyfish, *Leiognathus bindus*, from Western Indonesia based on data from R/Vs *Mutiara 4*, *Jurong* and *Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 35). Open dots represent outliers, not used for analysis. [Gambar 63. (A) Gambaran nilai ekstrim untuk ikan caria, *Leiognathus bindus*, di Indonesia berdasarkan data dari kapal penelitian *Jurong* menunjukkan 5 contoh frekuensi-panjang dan angka perkiraan $L_{max3} = 14.8 \pm 1.15$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 17 contoh frekuensi-panjang ikan caria, *Leiognathus bindus*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Dr. Fridtjof Nansen* sebagai output perangkat lunak ABee (lihat Boks 1), yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 35). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 64. Distribution of orangefin ponyfish, *Leiognathus bindus*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.

[Gambar 64. Penyebaran ikan caria, *Leiognathus bindus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 65. Depth distribution of orangefin ponyfish, *Leiognathus bindus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong and (D) Bawal Putih 2.

[Gambar 65. Penyebaran kedalaman ikan caria, *Leiognathus bindus*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong dan (D) Bawal Putih 2.]

Leiognathus equulus (Forsskål, 1775)

Common ponyfish (English); Peperek topang (Indonesian); Dodok (Java); Perek topang, Peperek topang, Peperek Tjina (West Java, Jakarta); Lokmolok (Madura); Molok-molok (Madura); Petek kuning (South Borneo); Bebeta (South Sulawesi, Badjo).

Body silvery; caudal peduncle with a small brown saddle; anal fins yellowish; dorsal fin transparent. Strongly arched back. Naked head, with nuchal spine. Protracted mouth pointing downward. Dorsal spines: 8-8; soft rays: 16-16; anal spines: 3-3; soft rays: 14-14. $L_{max1} = 25$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 28.4$ cm TL (Fig. 66A). See Fig. 66B and Table 38 for length-weight relationship.

Indo-West Pacific: from East London, South Africa including Réunion, Comores, Seychelles, Madagascar and Mauritius, Zanzibar, the Red Sea, Persian Gulf, India and Sri Lanka and thence to Southeast Asia and the islands of Indonesia (Fig. 67). Northeast to Okinawa, Ryukyu Islands; south to Australia and Fiji.

Occurs in river mouths and muddy inshore areas. Depth range: 10-110 m (Fig. 68). Feeds on polychaetes, small crustaceans, and small fish. Table 39 presents a set of growth parameters from Indonesia.

References: 186, 312, 393, 986, 1263, 1314, 1449, 1486, 1602, 2029, 2108, 2857, 3424, 3605, 3670, 3678, 4789, 4867, 4959, 5213, 5301, 5339, 5346, 5381, 5525, 5736, 5756, 6026, 6313, 6567

Table 38. Length-weight (g/[TL;cm]) relationship of common ponyfish, *Leiognathus equulus*, in Indonesia. [Tabel 38. Hubungan panjang-berat (g/[TL;cm]) ikan peperek topang, *Leiognathus equulus*, di Indonesia.]

Parameter	Estimate
a	0.0023
s.e.(a)	0.0031
b	3.6738
s.e.(b)	0.4120
r^2	0.9398

Table 39. Growth parameters of common ponyfish, *Leiognathus equulus*.

[Table 39. Parameter pertumbuhan ikan peperek topang, *Leiognathus equulus*.]

Parameter	A
L_{∞} (TL, cm)	21.5
K (year ⁻¹)	1.50

A. Java Sea (Central Java) (Ref. 1314)




Fig. 66. (A) Extreme value plot for common ponyfish, *Leiognathus equulus*, in Indonesia based on data from R/Vs Mutiara 4 and Jurong showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 28.4 \pm 2.65$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 14 length-frequency samples of common ponyfish, *Leiognathus equulus*, from Western Indonesia based on data from R/Vs Mutiara 4, Jurong and Dr. Fridtjof Nansen as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 37). Open dot(s) represent outliers, not used for analysis.

[Gambar 66. (A) Gambaran nilai ekstrim ikan peperek topang, *Leiognathus equulus*, di Indonesia berdasarkan data dari kapal-kapal penelitian Mutiara 4 dan Jurong yang menunjukkan nilai maksimum untuk 4 contoh frekuensi-panjang dan angka perkiraan $L_{max3} = 28.4 \pm 2.65$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 14 contoh frekuensi-panjang ikan peperek topang, *Leiognathus equulus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Mutiara 4, Jurong dan Dr. Fridtjof Nansen sebagai output perangkat lunak ABee (lihat Boks 1), dan memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 37). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 67. Distribution of common ponyfish, *Leiognathus equulus*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.
 [Gambar 67. Penyebaran ikan peperek topang, *Leiognathus equulus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 68. Depth distribution of common ponyfish, *Leiognathus equulus*, based on surveys of R/Vs (A) Mutiara 4, (B) Jurong and (C) Bawal Putih 2.
 [Gambar 68. Penyebaran kedalaman ikan peperek topang, *Leiognathus equulus*, berdasarkan survei kapal-kapal penelitian (A) Mutiara 4, (B) Jurong dan (C) Bawal Putih 2.]

Gymnocranius grandoculis (Valenciennes, 1830)

Rippled barenose.

Forehead profile moderately steep; large adults develop a bony ridge on the nape and a bony shelf over the front part of the eye. The inner surface of the pectoral fin axil is scaleless. Overall color is silvery with thin brown scale margins. The anterior half of the head is often brown with a series of narrow undulating, longitudinal lines on the cheek and side of the snout. Fins are yellow or orange; caudal fin is frequently dusky brown; a narrow brown bar across the base of pectoral fins. Juveniles under about 25 cm SL often with 5 or 6 dark bars on the side and a dark bar below the eye. Dorsal spines: 10-10; soft rays: 10-10; anal spines: 3-3; soft rays: 10-10. $L_{max1} = 80$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 74.3$ cm FL (Fig. 75A). See Fig. 75B and Table 43 for length-weight relationship.

Widely distributed from East Africa in the Indian Ocean via Southeast Asia to Japan in the north, and Indonesia (Fig. 76), Australia and Oceania.

Inhabits trawling grounds of the continental shelves and offshore rocky bottoms. Depth range: 20-170 m (Fig. 77). Feeds mostly on benthic invertebrates and small fishes.

References: 171, 1830, 2030, 2290, 2295, 4537, 4830, 5213, 5450, 5525, 5756, 6567

Table 43. Length-weight (g/[FL;cm]) relationship of rippled barenose, *Gymnocranius grandoculis*, in Indonesia. [Tabel 43. Hubungan panjang-berat (g/[FL;cm]) ikan *Gymnocranius grandoculis*, di Indonesia.]

Parameter	Estimate
a	0.2492
s.e.(a)	0.1445
b	2.3647
s.e.(b)	0.1538
r ²	0.8875




Fig. 75. (A) Extreme value plot for rippled barenose, *Gymnocranius grandoculis*, in Indonesia based on data from R/V *Jurong* showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 74.3 \pm 4.3$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 31 length-frequency samples of rippled barenose, *Gymnocranius grandoculis*, from Western Indonesia based on data from R/Vs *Mutiara 4*, *Jurong* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 43). Open dots represent outliers, not used for analysis. [Gambar 75. (A) Gambaran nilai ekstrim ikan *Gymnocranius grandoculis* di Indonesia berdasarkan data dari kapal penelitian *Jurong* menunjukkan nilai maksimum untuk 6 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 74.3 \pm 4.3$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 31 contoh frekuensi-panjang ikan *Gymnocranius grandoculis* dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Bawal Putih 2* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 43). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 76. Distribution of rippled barenose, *Gymnocranius grandoculis*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.
 [Gambar 76. Penyebaran ikan *Gymnocranius grandoculis* berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 77. Depth distribution of rippled barenose, *Gymnocranius grandoculis*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Jurong and (C) Bawal Putih 2.
 [Gambar 77. Penyebaran kedalaman ikan *Gymnocranius grandoculis* berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Jurong dan (C) Bawal Putih 2.]

***Aprion virescens* (Valenciennes, 1830)**

Green jobfish.

Preopercle edge smooth or sometimes denticulate in juveniles. There is a distinct horizontal groove in front of eye. Dorsal and anal fins scaleless. Scale rows on back parallel with lateral line. Color dark green to bluish or blue-gray. Dorsal spines: 10-10; soft rays: 11-11; anal spines: 3-3; soft rays: 8-8. $L_{max1} = 112$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 86.9$ cm FL (Fig. 78A). See Fig. 78B and Table 44 for length-weight relationship.

Widely distributed in the tropical Indo-Pacific Ocean from East Africa via Southeast Asia to southern Japan and Hawaii, and southward via Indonesia (Fig. 79) to Australia.

Inhabits inshore reef areas, usually solitary. Depth range: 20-100 m (Fig. 80). Feeds mainly on fishes, but also shrimps, crabs, cephalopods and planktonic organisms.

References: 55, 171, 245, 280, 583, 1602, 1830, 2290, 3084, 3090, 3111, 3670, 3678, 3804, 3807, 4517, 4690, 4699, 4795, 4821, 4868, 4887, 5213, 5358, 5450, 5525, 5579, 5736, 5756, 6089, 6273, 6306, 6365

Table 44. Length-weight (g/[FL;cm]) relationship of green jobfish, *Aprion virescens*, in Indonesia.

Tabel 44. Hubungan panjang-berat (g/[FL;cm]) ikan *Aprion virescens* di Indonesia.

Parameter	Estimate
a	0.0077
s.e.(a)	0.0039
b	3.1368
s.e.(b)	0.1181
r ²	0.9922




Fig. 78 (A) Extreme value plot for green jobfish, *Aprion virescens*, in Indonesia based on data from *R/V Jurong* showing maxima of 6 length-frequency samples, and estimate of $L_{max3} = 86.9 \pm 6.5$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 12 length-frequency samples of green jobfish, *Aprion virescens*, from Western Indonesia based on data from *R/V Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 44). Open dots represent outliers, not used for analysis.

[Gambar 78. (A) Gambaran nilai ekstrim ikan *Aprion virescens* di Indonesia berdasarkan data dari kapal penelitian *Jurong* yang menunjukkan nilai maksimum untuk 6 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 86.9 \pm 6.5$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 12 contoh frekuensi-panjang ikan *Aprion virescens* dari Indonesia bagian barat berdasarkan data dari kapal penelitian *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 44). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 79. Distribution of green jobfish, *Aprion virescens*, based on records of the surveys of R/Vs Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.

[Gambar 79. Penyebaran ikan *Aprion virescens* berdasarkan laporan survei kapal-kapal penelitian Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 80. Depth distribution of green jobfish, *Aprion virescens*, based on surveys of R/Vs (A) Jurong and (B) Bawal Putih 2.

[Gambar 80. Penyebaran kedalaman ikan *Aprion virescens* berdasarkan survei kapal-kapal penelitian (A) Jurong dan (B) Bawal Putih 2.]

Pristipomoides typus (Bleeker, 1852)

Sharptooth jobfish.

Interorbital space flat. Bases of dorsal and anal fins scaleless, their last soft rays extended into short filaments. Pectoral fins long, reaching level of anus. Scale rows on back parallel to lateral line. Overall color rosy red; the top of the head with longitudinal vermiculated lines and spots of brownish yellow; the dorsal fin with wavy yellow lines. Dorsal spines: 10-10; soft rays: 11-12; anal spines: 3-3; soft rays: 8-8. $L_{max1} = 70$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 68.3$ cm TL (Fig. 81A). See Fig. 81B and Table 45 for length-weight relationship.

Tropical western Pacific ranging in Indonesia from Sumatra to Irian Jaya (Fig 82) and northward to the Ryukyu Islands. Records from the western Indian Ocean need to be confirmed.

Occurs over rocky bottoms. Depth range: 40-120 m (Fig. 83). Feeds on benthic invertebrates and fishes.

References: 55, 171, 438, 1451, 2857, 3090, 4517, 4789, 5213, 5450, 5515, 5725, 5756, 6365, 6425, 6567

Table 45. Length-weight (g/[TL;cm]) relationship of sharptooth jobfish, *Pristipomoides typus*, in Indonesia.
 [Tabel 45. Hubungan panjang-berat [g/(TL;cm)] ikan *Pristipomoides typus* di Indonesia.]

Parameter	Estimate
a	0.0143
s.e.(a)	0.0175
b	2.9158
s.e.(b)	0.3156
r ²	0.9208




Fig. 81. (A) Extreme value plot for sharptooth jobfish, *Pristipomoides typus*, in Indonesia based on data from *R/Vs Mutiara 4* and *Dr. Fridtjof Nansen* showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 68.3 \pm 4.1$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 7 length-frequency samples of sharptooth jobfish, *Pristipomoides typus*, from Western Indonesia based on data from *R/Vs Mutiara 4*, *Bawal Putih 2* and *Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 45). Open dot represents outlier, not used for analysis.

[Gambar 81. (A) Gambaran nilai ekstrim ikan *Pristipomoides typus* di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Dr. Fridtjof Nansen* yang menunjukkan nilai maksimum untuk 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 68.3 \pm 4.1$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 7 contoh frekuensi-panjang *Pristipomoides typus* dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2* dan *Dr. Fridtjof Nansen* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 45). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]




Fig. 82. Distribution of sharptooth jobfish, *Pristipomoides typus*, based on records of the surveys of *R/Vs Mutiara 4*, *Bawal Putih 2*, *Jurong* and *Dr. Fridtjof Nansen*.
 [Gambar 82. Penyebaran ikan *Pristipomoides typus* berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2*, *Jurong* dan *Dr. Fridtjof Nansen*.]




Fig. 83. Depth distribution of sharptooth jobfish, *Pristipomoides typus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) Jurong. [Gambar 83. Penyebaran kedalaman ikan *Pristipomoides typus* berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) Jurong.]

Upeneus moluccensis (Bleeker, 1855)

Goldband goatfish (English); Bijnangka (Indonesian).

Body elongate, with relatively large ctenoid scales. Color is silvery white, with a bright yellow horizontal band running through the eye to the caudal fin. Dorsal fins with 3-4 orange or red bars; anal and pelvic fin pale. Upper lobe of the caudal fin with 5-6 orange-black bars, lower lobe plain yellow with dark margin. Dorsal spines: 13-13; soft rays: 9-9; anal spines: 0-0; soft rays: 7-7; $L_{max1} = 20$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 20.0$ cm FL (Fig. 84A). See Fig. 84B and Table 46 for length-weight relationship.

Occurs in the Indo-West Pacific from the east coast of Africa to Southeast Asia, the Indonesian Archipelago (Fig. 85) and the northern coasts of Australia; also reported from New Caledonia. Recently invaded the eastern Mediterranean from the Red Sea through the Suez Canal.

Found in coastal waters with a muddy substrate at depths ranging from 30 to 120 m (Fig. 86).

References: 393, 1263, 1449, 1486, 1975, 2029, 2178, 2795, 2857, 3397, 4789, 5213, 5381, 5385, 5450, 5525, 5756, 6306, 6328, 6567

Table 46. Length-weight (g/[FL;cm]) relationship of goldband goatfish, *Upeneus moluccensis*, in Indonesia.

[Tabel 46. Hubungan panjang-berat (g/[FL;cm]) ikan bijnangka, *Upeneus moluccensis*, di Indonesia.]

Parameter	Estimate
a	0.0451
s.e.(a)	0.0275
b	2.6364
s.e.(b)	0.2400
r^2	0.9631




Fig. 84. (A) Extreme value plot for goldband goatfish, *Upeneus moluccensis*, in Indonesia based on data from R/Vs Jurong and Dr. Fridtjof Nansen showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 20.0 \pm 1.1$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 15 length-frequency samples of goldband goatfish, *Upeneus moluccensis*, from Western Indonesia based on data from R/Vs Jurong, Dr. Fridtjof Nansen and Bawal Putih 2 as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 46).

[Gambar 84. (A) Gambaran nilai ekstrim ikan bijnangka, *Upeneus moluccensis*, di Indonesia berdasarkan data dari kapal-kapal penelitian Jurong dan Dr. Fridtjof Nansen yang menunjukkan nilai maksimum untuk 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 20.0 \pm 1.1$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 15 contoh frekuensi-panjang ikan bijnangka, *Upeneus moluccensis*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian Jurong, Dr. Fridtjof Nansen dan Bawal Putih 2 sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 46).




Fig. 85. Distribution of goldband goatfish, *Upeneus moluccensis*, based on records of the surveys of R/Vs *Jurong*, Dr. Fridtjof Nansen and *Bawal Putih 2*.

Gambar 85. Penyebaran ikan bijinangka, *Upeneus moluccensis*, berdasarkan laporan survei kapal-kapal penelitian *Jurong*, Dr. Fridtjof Nansen dan *Bawal Putih 2*.




Fig. 86. Depth distribution of goldband goatfish, *Upeneus moluccensis*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) *Jurong* and (C) *Bawal Putih 2*.

[Gambar 86. Penyebaran kedalaman ikan bijinangka, *Upeneus moluccensis*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) *Jurong* dan (C) *Bawal Putih 2*.]

Upeneus sulphureus (Cuvier, 1829)

Sulphur goatfish (English); Kunir (Indonesian); Kakunir, Kunir, Kuniran (Java); Bidji nangka (West Java, Jakarta).

Medium-sized fish of moderately elongate bodies. Head small; mouth small and slightly oblique; a pair of barbels under the chin. Dorsal fins with 2 to 3 olive bars, and black or dark brown tips; anal, pelvic and pectoral fins pale; caudal fin plain dull yellow, its hind margin dusky, its lower lobe tipped white. Two orange-yellow bands extend from the head to the caudal peduncle. Dorsal spines: 8-8; soft rays: 8-8; anal spines: 1-1; soft rays: 7-7. $L_{max1} = 23$ cm; $L_{max2} = n.a.$; $L_{max3} = 23$ cm TL (Fig. 87A). See Fig. 87B and Table 47 for length-weight relationship.

From East Africa to Southeast Asia; through Indonesia (Fig. 88); northward to the coast of China and southward to the northern coasts of Australia; also reported from New Caledonia.

Forms schools in coastal waters. Depth range: 10-90 m (Fig. 89). Table 48 presents four sets of growth parameters from Indonesia.

References: 393, 1263, 1314, 1379, 1392, 1435, 1449, 1474, 1486, 1966, 2029, 2110, 2178, 2857, 2871, 2926, 3470, 4749,

4789, 5213, 5381, 5405, 5450, 5525, 5736, 5756, 6292, 6365, 6567

Table 47. Length-weight (g/[TL;cm]) relationship of sulphur goatfish, *Upeneus sulphureus*, in Indonesia. [Tabel 47. Hubungan panjang-berat (g/[TL;cm]) ikan kunir, *Upeneus sulphureus*, di Indonesia.]

Parameter	Estimate	
	A	B
a	0.009	0.0081
s.e.(a)	n.a.	0.0027
b	3.193	3.2134
s.e.(b)	n.a.	0.1272
r ²	n.a.	0.9782

A. Java (north coast) (Ref. 1379)
B. This study

Table 48. Growth parameters of sulphur goatfish, *Upeneus sulphureus*.

[Tabel 48. Parameter pertumbuhan ikan kunir, *Upeneus sulphureus*.]

Parameter	A	B	C	D
L_{∞} (TL, cm)	15.8	16.5	17.5	19.9
K (year ⁻¹)	1.74	0.78	0.90	0.875

A. North Java Coast (Ref. 1435)
B. Java Sea (Central Java, 1978-79) (Ref. 1314)
C. Java Sea (Central Java, 1977-78) (Ref. 1314)
D. Java Sea (Ref. 1379)




Fig. 87. (A) Extreme value plot for sulphur goatfish, *Upeneus sulphureus*, in Indonesia based on data from *R/Vs Mutiara 4* and *Bawal Putih 2* showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 23 \pm 3.8$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 44 length-frequency samples of sulphur goatfish, *Upeneus sulphureus*, from Western Indonesia based on data from *R/Vs Mutiara 4*, *Jurong*, *Dr. Fridtjof Nansen* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 47). Open dots represent outliers, not used for analysis.

[Gambar 87. (A) Gambaran nilai ekstrim ikan kunir, *Upeneus sulphureus*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Bawal Putih 2* yang menunjukkan nilai maksimum untuk 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 23 \pm 3.8$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 44 contoh frekuensi-panjang ikan kunir, *Upeneus sulphureus*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian *Mutiara 4*, *Jurong*, *Dr. Fridtjof Nansen* dan *Bawal Putih 2* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 47). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 88. Distribution of sulphur goatfish, *Upeneus sulphureus*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.
 [Gambar 88. Penyebaran ikan kunir, *Upeneus sulphureus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 89. Depth distribution of sulphur goatfish, *Upeneus sulphureus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong and (D) Bawal Putih 2.
 [Gambar 89. Penyebaran kedalaman ikan kunir, *Upeneus sulphureus*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Mutiara 4, (C) Jurong dan (D) Bawal Putih 2.]

Nemipterus thosaporni (Russell, 1991)

Palefin threadfin bream (English); Kurisi (Indonesian).

Lower edge of eye touching or just above a line from tip of snout to upper pectoral-fin base; lower edge of suborbital slightly emarginate. Dorsal fin origin about 3-7 scale rows from imaginary line projected upward from posterior edge of suborbital to dorsal profile. Pectoral and pelvic fins long, reaching to or just short of level of anal-fin origin. Closely resembles *N. bathybius*, but has no yellow stripe on either side of the ventral midline and the upper tip of the caudal fin not drawn into a distinct filament. Axillary scale present. Color: Upper part pinkish, silvery below. Dorsal spines: 10-10; soft rays: 9-9; anal spines: 3-3; soft rays: 7-7. $L_{\max 1} = 21.5$ cm SL; $L_{\max 2} = 23$ cm TL; $L_{\max 3} = \text{n.a.}$ See Table 49 for length-weight relationship.

Widely distributed throughout the Western Pacific, notably in the Strait of Malacca, the Gulf of Thailand, the Sunda Islands, Indonesia (Fig. 90), and to southern Japan. This species has been previously misidentified as *N. marginatus* by most authors. Fig. 90 shows its distribution based on records of the *R/Vs Mutiara 4*, *Jurong* and *Dr. Fridtjof Nansen* surveys; Fig. 91 provides details on the distribution of *N. thosaporni* in the southern part of the South China Sea.

Found on sand or mud bottoms. Depth range: 10-80 m (Fig. 92). During that part of the *R/V Mutiara 4* survey which covered Area 5 in Pauly et al. (this vol.), i.e., the southern part of the South China Sea, D. Pauly and P. Martosubroto (Ref. 1158) measured a large number of nemipterids belonging to this species, which they thought was *Nemipterus marginatus*. This does not invalidate the results obtained by these two authors, and their main findings which are recalled here (see Box 4). Table 50 presents a set of growth parameters from Indonesia.

References: 171, 1066, 1139, 1158, 3207, 3810

Table 49. Length-weight (g/[TL;cm]) relationship of palefin threadfin bream, *Nemipterus thosaporni*, in Indonesia. [Tabel 49. Hubungan panjang-berat (g/[TL;cm]) ikan kurisi, *Nemipterus thosaporni*, di Indonesia.]

Parameter	Estimate*
a	0.0135
b	3.02
r	0.999

*West Kalimantan (Ref. 1158)

Table 50. Growth parameters of palefin threadfin bream, *Nemipterus thosaporni*. [Tabel 50. Parameter pertumbuhan ikan kurisi, *Nemipterus thosaporni*.]

Parameter	A	B
L_{∞} (TL, cm)	24.5	28.4
K (year ⁻¹)	0.420	0.363

A. Western Kalimantan (Ref. 1158)

B. Sarawak and Sabah (Northern Kalimantan) (Ref. 1139)




Fig. 90. Distribution of palefin threadfin bream, *Nemipterus thosaporni*, based on records of the surveys of *R/Vs Mutiara 4*, *Jurong* and *Dr. Fridtjof Nansen*.

[Gambar 90. Penyebaran ikan kurisi, *Nemipterus thosaporni*, berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Dr. Fridtjof Nansen*.]




Fig. 91. Distribution and relative abundance of palefin threadfin bream, *Nemipterus thosaporni*, in the southern part of the South China Sea from 9 August to 29 September 1975. See Pauly et al. (this vol.) for details on this survey.
 [Gambar 91. Penyebaran dan kelimpahan relatif ikan kurisi, *Nemipterus thosaporni*, di bagian selatan Laut Cina Selatan dari 9 Agustus hingga 29 September 1975. Lihat Pauly et al. (dalam buku ini) untuk rincian survei ini.]




Fig. 92. Depth distribution of palefin threadfin bream, *Nemipterus thosaporni*, based on surveys of R/Vs (A) *Dr. Fridtjof Nansen*, (B) *Mutiara 4* and (C) *Jurong*.
 [Gambar 92. Penyebaran kedalaman ikan kurisi, *Nemipterus thosaporni*, berdasarkan survei kapal-kapal penelitian (A) *Dr. Fridtjof Nansen*, (B) *Mutiara 4* dan (C) *Jurong*.]

Box 4. A case study of *Nemipterus thosaporni* a.k.a *N. marginatus*.
 [Boks 4. Suatu studi kasus dari *Nemipterus thosaporni*, yang dikenal juga dengan nama *N. marginatus*.]

Purwito Martosubroto and I measured, from 6 August to 29 September 1975 (Ref. 1158) the 3,283 specimens of *N. thosaporni* (which we called *N. marginatus*) in the table below during a survey off Western Kalimantan (see Fig. 91).

Summary of length-frequency data on *Nemipterus thosaporni* from Western Kalimantan (= 3,283).

Lower limit of class (TL; cm)	N*	Lower limit of class (TL; cm)	N*
7.5	1	15.5	172
8.0	3	16.0	140
8.5	9	16.5	133
9.0	16	17.0	102
9.5	40	17.5	79
10.0	55	18.0	82
10.5	100	18.5	78
11.0	122	19.0	59
11.5	157	19.5	39
12.0	200	20.0	24
12.5	218	20.5	12
13.0	211	21.5	13
13.5	287	21.5	8
14.0	356	22.0	2
14.5	334	22.5	3
15.0	226	23.0	2

* Sum of 44 samples: Stations 59-147: see Fig. 4 in Pauly et al. (this vol.).

In the absence of computers, we used the then popular, graphical "Cassie method" (Ref. 9564) to split our cumulative samples into three normally distributed components, to which we assigned relative ages, which were then used to estimate von Bertalanffy growth parameters that compared well with previous estimate from Northern Kalimantan (Table 50). These growth parameters, complemented with a length-weight relationship (Table 49), and an estimate of M - estimated from the size distribution in the then unexploited stock - allowed computation of yield-per-recruit curves.

This entire procedure - although involving no development of new methodology - was exemplary in that it illustrated how a wide range of analytical techniques could be applied to data obtained during a fairly standard trawl survey, and a more or less complete "assessment" thus being performed using data then generally not perceived as being sufficient for such purpose.

Although it has been cited perhaps 20 times to date, this work is now rather well known among fisheries scientists in the tropics because it formed the base of a "case study", taught in the 1980s by Dr. Erik Ursin, of the roving FAO/DANIDA Training Course in Tropical Fish Stock Assessment, and consisting of the following elements:

- i) evaluation of the work's methodology, based on copies of all paper cited in its "Methods" section;
- ii) evaluation of the "Results" section, based on recomputation of all estimates, and re-evaluation of all assumptions (explicit and implicit); and
- iii) evaluation of the "Discussion" section, through comparison with similar results in contemporary contributions (e.g., Ref. 1066), and later advances.

The paper survived this rather stringent test of its replicability, and the fish was thus allowed to migrate, via my textbook of 1984 (Ref. 4715) into the text that emerged from the above-mentioned training course (Ref. 9566).

I wish we had written more such papers.

Daniel Pauly
 ICLARM
 and
 Fisheries Centre, UBC

Priacanthus macracanthus (Cuvier 1829)

Red bigeye (English); Swanggi (Indonesian); Swanggi (Javanese).

Medium-sized fish of moderately deep body. The eyes large; the mouth oblique, with the lower jaw projecting upwards. The body tapers very slightly to beneath the middle of the soft portion of the dorsal fin, then abruptly to the peduncle. This species is distinguished from *Priacanthus fitchi* by the presence of numerous rusty brown to yellowish spots in the membranes of the dorsal and anal fins, and its less tapered body. Dorsal spines: 10-10; soft rays: 12-14; anal spines: 3-3; soft rays: 13-14. $L_{max1} = 29$ cm SL; $L_{max2} = n.a.$; $L_{max3} = 25.2$ cm TL (Fig. 93A). See Fig. 93B and Table 51 for length-weight relationship.

East Indo-West Pacific: from southern Japan in the north to Western Indonesia (Fig. 94) and Australia in the south.

Occurs in inshore and offshore reefs, apparently forms aggregations in open bottom areas. Depth range: 20-350 m (Fig. 95). Table 52 presents 2 sets of growth parameters from Indonesia.

References: 559, 1263, 1314, 1449, 2857, 3132, 3414, 4539, 4885, 5381, 5736, 5756

Table 51. Length-weight (g/[TL;cm]) relationship of red bigeye, *Priacanthus macracanthus*, in Indonesia.
 Tabel 51. Hubungan panjang-berat (g/[TL;cm]) ikan swanggi, *Priacanthus macracanthus*, di Indonesia.

Parameter	Estimate
a	0.0163
s.e.(a)	0.0072
b	2.9914
s.e.(b)	0.1648
r ²	0.9543

Table 52. Growth parameters of red bigeye, *Priacanthus macracanthus*.

Tabel 52. Parameter pertumbuhan ikan swanggi, *Priacanthus macracanthus*.

Parameter	A	B
L_{∞} (TL, cm)	23	23.8
K (year ⁻¹)	1.15	1.30

A. Java Sea (Central Java, 1978-79) (Ref. 1314)

B. Java Sea (Central Java, 1977-78) (Ref. 1314)




Fig. 95. Depth distribution of red bigeye, *Priacanthus macracanthus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) *Jurong*.
[Gambar 95. Penyebaran kedalaman ikan swanggi, *Priacanthus macracanthus*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) *Jurong*.]

Rastrelliger kanagurta (Cuvier, 1816)

Indian mackerel (English); Kembung lelaki (Indonesian); Banjar, Kembung lelaki (West Java, Jakarta); Gombong (Central Java); Bulus lake, Saängsa (Madura); Banjara (South Sulawesi, Makassar); Botto-botto (South Sulawesi, Bugis); Banjar (South Sulawesi, Badjo).

Head longer than body depth. Maxilla partly concealed, covered by lacrimal bone but extending to about hind margin of eye. Bristles on longest gillraker 105 on one side in specimens of 12.7 cm, 140 in 16 cm, and 160 in 19 cm fork length specimens. A black spot on body near lower margin of pectoral fin. Interpelvic process small and single. Swimbladder present. Anal spine rudimentary. Dorsal spines: 8-11; soft rays: 12-12; anal spines: 0-0; soft rays: 12-12. $L_{max1} = 36$ cm TL; $L_{max2} = 26$ cm; $L_{max3} = 26.8$ cm TL (Fig. 96A). See Fig. 96B and Table 53 for length-weight relationship.

Indo-West Pacific: from South Africa and the Seychelles in the east to the Red Sea, and Southeast Asia; Indonesia (Fig. 97); north to the Ryukyu Islands, China. Southeast to Northern Australia, Melanesia, Micronesia, Samoa. Entered the eastern Mediterranean Sea through the Suez Canal.

Form schools in coastal waters, bays and deep lagoons, usually in plankton-rich waters. Depth range: 20-90 m (Fig. 98). Feeds on phytoplankton (diatoms) and small zooplankton (cladocerans, ostracods, larval polychaetes, etc.). Adult individuals feed on macroplankton (larval shrimps and fish). Table 54 presents six sets of growth parameters from Indonesia.

References: 168, 171, 312, 762, 786, 821, 1139, 1195, 1196, 1197, 1198, 1263, 1314, 1389, 1392, 1447, 1449, 1462, 1463, 1464, 1465, 1466, 1467, 1485, 1488, 1531, 1602, 1687, 1751, 1836, 2178, 3557, 3626, 3621, 3579, 3669, 3670, 3678, 4546, 4547, 4593, 4749, 4789, 4838, 5213, 5284, 5385, 5450, 5756

Table 53. Length-weight (g/[TL;cm]) relationship of Indian mackerel, *Rastrelliger kanagurta*, in Indonesia.
[Tabel 53. Hubungan panjang-berat (g/[TL;cm]) ikan kembung lelaki, *Rastrelliger kanagurta*, di Indonesia.]

Parameter	Estimates				
	A	B	C	D	E
a	0.0039	0.0061	0.0022	0.0014	0.0061
s.e. (a)	n.a.	n.a.	n.a.	n.a.	0.0027
b	3.1900	3.1910	3.3300	3.3770	3.1743
s.e. (b)	n.a.	n.a.	n.a.	n.a.	0.1437
r ²	n.a.	n.a.	n.a.	n.a.	0.9909

A. Indonesia, Java Sea (Ref. 1463)
B. Indonesia, Java Sea (Ref. 1196)
C. Indonesia, Andaman Islands (Ref. 1463)
D. Indonesia, Malacca Strait (Ref. 1389)
E. This study

Table 54. Growth parameters of Indian mackerel, *Rastrelliger kanagurta*.
 Tabel 54. Parameter pertumbuhan ikan kembung lelaki, *Rastrelliger kanagurta*.

Parameter	A	B	C	D	E	F
L_{∞} (TL, cm)	23.9	25.7	25.8	26.5	28.5	28.7
K (year ⁻¹)	2.76	1.625	1.63	0.80	0.90	0.78

- A. Indonesia, Java Sea (Ref. 1196)
- B. Indonesia, Java Sea (Ref. 1447)
- C. Indonesia, Java Sea (Pekalongan, 1982-83) (Ref. 1314)
- D. Indonesia, Asahan, Sumatra (Ref. 1467)
- E. Indonesia, Banda Aceh (Ref. 4547)
- F. Indonesia, Strait of Malacca (1984-86) (Ref. 1389)




Fig. 96. (A) Extreme value plot for Indian mackerel, *Rastrelliger kanagurta*, in Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 26.8 \pm 0.85$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 16 length-frequency samples of Indian mackerel, *Rastrelliger kanagurta*, from Western Indonesia based on data from R/Vs *Mutiara 4*, *Bawal Putih 2*, *Jurong* and *Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 53). Open dots represent outliers, not used for analysis.

[Gambar 96. (A) Gambaran nilai ekstrim ikan kembung lelaki, *Rastrelliger kanagurta*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* yang menunjukkan nilai maksimum 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 26.8 \pm 0.85$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 16 contoh frekuensi-panjang ikan kembung lelaki, *Rastrelliger kanagurta*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2*, *Jurong* dan *Dr. Fridtjof Nansen* sebagai luaran perangkat lunak ABee (lihat Box 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 53). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 97. Distribution of Indian mackerel, *Rastrelliger kanagurta*, based on records of the surveys of R/Vs *Mutiara 4*, *Bawal Putih 2*, *Jurong* and *Dr. Fridtjof Nansen*.
 [Gambar 97. Penyebaran ikan kembung lelaki, *Rastrelliger kanagurta*, berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2*, *Jurong* dan *Dr. Fridtjof Nansen*.]




Fig. 98. Depth distribution of Indian mackerel, *Rastrelliger kanagurta*, based on surveys of R/Vs (A) *Dr. Fridtjof Nansen* and (B) *Bawal Putih 2*. [Gambar 98. Penyebaran kedalaman ikan kembung lelaki, *Rastrelliger kanagurta*, berdasarkan survei kapal-kapal penelitian (A) *Dr. Fridtjof Nansen* dan (B) *Bawal Putih 2*.]

***Scomberomorus commerson* (Lacepède, 1800)**

Narrow-barred Spanish mackerel (English); Tjalong (Indonesian); Tengiri (West Java, Jakarta); Langung, Tengere, Tjalong, Tjangetjang (Madura).

Interpelvic process small and bifid. Swimbladder absent. Lateral line abruptly bent downward below end of second dorsal

fin. Intestine with 2 folds and 3 limbs. Vertical bars on trunk sometimes break up into spots ventrally which number 40-50 in adults, and less than 20 in juveniles (which have jet black anterior first dorsal fin). Dorsal spines: 15-18; soft rays: 15-20; anal spines: 0-0; soft rays: 16-21. $L_{max1} = 220$ cm FL; $L_{max2} = n.a.$; $L_{max3} = 96.9$ cm FL (Fig. 99A). See Fig. 99B and Table 55 for length-weight relationship.

Indo-West Pacific: from South Africa and the Red Sea




Fig. 99. (A) Extreme value plot for narrow-barred Spanish mackerel, *Scomberomorus commerson*, in Indonesia based on data from R/V *Jurong* showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 96.9 \pm 26.2$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 32 length-frequency samples of narrow-barred Spanish mackerel, *Scomberomorus commerson*, from Western Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 55). Open dots represent outliers, not used for analysis.

[Gambar 99. (A) Gambaran nilai ekstrim ikan tenggiri papan, *Scomberomorus commerson*, di Indonesia berdasarkan data dari kapal penelitian *Jurong* menunjukkan nilai maksimum dari 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 96.9 \pm 26.2$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 32 contoh frekuensi panjang ikan tenggiri papan, *Scomberomorus commerson*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* sebagai luaran perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 55). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]

through Southeast Asia, north to China and Japan and southward to Indonesia (Fig. 100) and Southeast Australia. A recent immigrant to the eastern Mediterranean Sea by way of the Suez Canal. Atlantic Ocean: reported only from St. Helena.

Found in small schools and known to undertake lengthy longshore migrations, but permanently resident populations also seem to exist. Depth range: 10-70 m (Fig. 101). Feeds primarily on small fish such as anchovies, clupeids, carangids, squids and penaeid shrimps.

References: 168, 171, 1139, 1263, 1375, 1391, 1415, 1416, 1470, 1498, 1602, 2325, 2682, 2857, 3383, 3557, 3626, 3678, 4332, 4588, 4699, 4883, 4905, 5213, 5284, 5288, 5385, 5450, 5515, 5736, 5756, 5765, 5766, 5970, 6026, 6323, 6365, 6783

Table 55. Length-weight (g/[FL;cm]) relationship of narrow-barred Spanish mackerel, *Scomberomorus commerson*, in Indonesia.

[Tabel 55. Hubungan panjang-berat (g/[FL;cm]) ikan tenggiri papan, *Scomberomorus commerson*, di Indonesia.]

Parameter	Estimate
a	0.0057
s.e. (a)	0.0046
b	3.1247
s.e. (b)	0.2094
r ²	0.9271




Fig. 100. Distribution of narrow-barred Spanish mackerel, *Scomberomorus commerson*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.

[Gambar 100. Penyebaran ikan tenggiri papan, *Scomberomorus commerson*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 101. Depth distribution of narrow-barred Spanish mackerel, *Scomberomorus commerson*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) Bawal Putih 2.

[Gambar 101. Penyebaran kedalaman ikan tenggiri papan, *Scomberomorus commerson*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) Bawal Putih 2.]

Scomberomorus guttatus (Bloch & Schneider, 1801)

Indo-Pacific king mackerel (English); Tenggiri (Indonesian); Ajong-ajong, Usek-usek (Java); Tengiri (West Java, Jakarta); Langung, Tengere, Tjalong, Tjangetjang (Madura); Tengiri (South Borneo).

Interpelvic process small and bifid. Swimbladder absent. Body entirely covered with small scales. Lateral line with many auxiliary branches extending dorsally and ventrally in anterior third, curving down toward caudal peduncle. Intestine with 2 folds and 3 limbs. Sides silvery white with several rows of round dark brownish spots scattered in about three irregular rows along the lateral line. First dorsal fin membrane black. Dorsal spines: 15-18; soft rays: 18-24; anal spines: 0-0; soft rays: 19-23. $L_{max1} = 76$ cm FL; $L_{max2} = n.a.$; $L_{max3} = 64.4$ cm FL (Fig. 102A). See Fig. 102B and Table 56 for length-weight relationship.

Indo-West Pacific from the Persian Gulf, India and Sri Lanka to Southeast Asia, Indonesia (Fig. 103); north to Hong Kong and Wakasa Bay, Sea of Japan.

Depth range: 20-90 m (Fig. 104). A pelagic migratory fish inhabiting coastal waters; sometimes entering turbid estuarine waters, usually found in small schools. Feeds mainly on small schooling fishes (especially sardines and anchovies), squids and crustaceans.

References: 168, 171, 280, 298, 2682, 3383, 4515, 4588, 4883, 5515, 5285, 5736, 5756, 6313, 6365, 6567

Table 56. Length-weight (g/[FL;cm]) relationship of Indo-Pacific king mackerel, *Scomberomorus guttatus*, in Indonesia. [Tabel 56. Hubungan panjang-berat (g/[FL;cm]) ikan tenggiri, *Scomberomorus guttatus*, di Indonesia.]

Parameter	Estimate
a	0.0096
s.e.	0.0053
b	3.0020
s.e.	0.1515
r ²	0.9777




Fig. 102. (A) Extreme value plot for Indo-Pacific king mackerel, *Scomberomorus guttatus*, in Indonesia based on data from *R/Vs Mutiara 4, Bawal Putih 2* and *Dr. Fridtjof Nansen* showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 64.4 \pm 4.35$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 16 length-frequency samples of Indo-Pacific king mackerel, *Scomberomorus guttatus*, from Western Indonesia based on data from *R/Vs Mutiara 4, Bawal Putih 2, Dr. Fridtjof Nansen* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 56). Open dots represent outliers, not used for analysis.

[Gambar 102. (A) Gambaran nilai ekstrim ikan tenggiri, *Scomberomorus guttatus*, di Indonesia berdasarkan data kapal-kapal penelitian *Mutiara 4, Bawal Putih 2* dan *Dr. Fridtjof Nansen* menunjukkan nilai maksimum dari 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 64.4 \pm 4.35$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 16 contoh frekuensi-panjang ikan tenggiri, *Scomberomorus guttatus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4, Bawal Putih 2, Dr. Fridtjof Nansen* dan *Jurong* sebagai luaran perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 56). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 103. Distribution of Indo-Pacific king mackerel, *Scomberomorus guttatus*, based on records of the surveys of R/Vs *Mutiara 4*, *Bawal Putih 2*, *Jurong* and *Dr. Fridtjof Nansen*.

[Gambar 103. Penyebaran ikan tenggiri, *Scomberomorus guttatus*, berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2*, *Jurong* dan *Dr. Fridtjof Nansen*.]




Fig. 104. Depth distribution of Indo-Pacific king mackerel, *Scomberomorus guttatus*, based on surveys of R/Vs (A) *Dr. Fridtjof Nansen* and (B) *Jurong*.

[Gambar 104. Penyebaran kedalaman ikan tenggiri, *Scomberomorus guttatus*, berdasarkan survei kapal-kapal penelitian (A) *Dr. Fridtjof Nansen* dan (B) *Jurong*.]

Sphyaena obtusata (Cuvier, 1829)

Obtuse barracuda (English); Tantjak (Indonesia); Alu-alu, Kutjul, Langsar (Java); Alu-alu (West Java, Jakarta); Kotjol, Tantjak (Madura).

Body elongate and subcylindrical with small cycloid scales; head long and pointed. Mouth large and horizontal, the tip of the lower jaw protruding; intermaxilla non-protractile. Preoperculum rectangular, with wide naked skin flap. First dorsal fin origin slightly before the pectoral fin tip, the first spine equal to the second. Pelvic fins well before the tip of the pectoral, closer to the anal than the tip of the lower jaw. Color is generally green above and silvery below. Dorsal spines: 6-6; soft rays: 9-9; anal spines: 2-2; soft rays: 9-9. $L_{max1} = 55$ cm; $L_{max2} =$

n.a.; $L_{max3} = 47.3$ cm FL (Fig. 105A). See Fig. 105B and Table 57 for length-weight relationship.

Indo-Pacific Ocean: East Africa and Red Sea to Philippines and Indonesia (Fig. 106); from Samoa north to Ryukyus, south to Lord Howe Islands; Kapingamarangi and Marianas in Micronesia. Migrated to eastern Mediterranean from the Red Sea via the Suez Canal.

Inhabits bays and estuaries. Found in schools in seagrass beds and rocky reefs. Depth range: 20-120 (Fig. 107). Feeds mainly on fishes.

References: 560, 1365, 1602, 2857, 4752, 5213, 5381, 5385, 5450, 5525, 5579, 5736, 5756, 6328, 6365, 6567

Table 57. Length-weight (g/[FL;cm]) relationship of obtuse barracuda, *Sphyraena obtusata*, in Indonesia.
 [Tabel 55. Hubungan panjang-berat (g/[FL;cm]) ikan alu-
 alu, *Sphyraena obtusata*, di Indonesia.]

Parameter	Estimate
a	0.0095
s.e.(a)	0.0031
b	2.8678
s.e.(b)	0.0977
r ²	0.9961




Fig. 105. (A) Extreme value plot for obtuse barracuda, *Sphyraena obtusata*, in Indonesia based on data from R/V *Jurong* showing maxima of 4 length-frequency samples, and estimate of $L_{max3} = 47.3 \pm 7.0$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 19 length-frequency samples of obtuse barracuda, *Sphyraena obtusata*, from Western Indonesia based on data from R/Vs *Mutiara 4*, *Jurong* and *Dr. Fridtjof Nansen* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 57). Open dots represent outliers, not used for analysis. [Gambar 105. (A) Gambaran nilai ekstrim ikan alu-*alu*, *Sphyraena obtusata*; di Indonesia berdasarkan data kapal penelitian *Jurong* menunjukkan nilai maksimum dari 4 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 47.3 \pm 7.0$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 19 contoh frekuensi-panjang ikan alu-*alu*, *Sphyraena obtusata*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Dr. Fridtjof Nansen*, sebagai luaran perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 57). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 106. Distribution of obtuse barracuda, *Sphyraena obtusata* based on records of the surveys of R/Vs *Mutiara 4*, *Bawal Putih 2*, *Jurong* and *Dr. Fridtjof Nansen*. [Gambar 106. Penyebaran ikan alu-*alu*, *Sphyraena obtusata*, berdasarkan laporan survei kapal-kapal penelitian *Mutiara 4*, *Bawal Putih 2*, *Jurong* dan *Dr. Fridtjof Nansen*.]




Fig. 107. Depth distribution of obtuse barracuda, *Sphyraena obtusata* based on surveys of R/Vs (A) Dr. Fridtjof Nansen, (B) Jurong and (C) Bawal Putih 2. [Gambar 107. Penyebaran kedalaman ikan alu-alu, *Sphyraena obtusata*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen, (B) Jurong dan (C) Bawal Putih 2.]

Pampus argenteus (Euphrasen, 1788)

Silver pomfret (English); Bawal putih (Indonesian); Dawah, Dawahan, Lawang, Lowang (Java); Bawal, Bawal putih (West Java, Jakarta); Njiuran, Njor njoran, Potean, Potian, Tangkolok, Tjeplak (Madura); Manriwasakebo (South Sulawesi, Makassar); Peda-peda puti (South Sulawesi, Bugis).

Body very deep, and compressed. Operculum absent; gill opening reduced to a vertical slit on the side of the body; gill membrane broadly united to isthmus. Dorsal and anal fins preceded by a series of 5 to 10 blade-like spines with anterior and posterior points. Pelvic fins absent. Caudal fin deeply forked, the lower lobe longer than the upper. Color is gray above grading to silvery white towards the belly, with small black dots all over the body. Fins are faintly yellow; vertical fins with dark edges. $L_{max1} = 60$ cm; $L_{max2} = n.a.$; $L_{max3} = 30.7$ cm FL (Fig. 108A). See Fig. 108B and Table 58 for length-weight relationship.

Indo-West Pacific: from the Persian Gulf east to Southeast Asia, Indonesia (Fig. 109) and north to southern Japan.

Found in coastal waters over muddy bottoms, associated with prawns and *Nemipterus* and *Leiognathus* species. Forms schools which can be large and abundant. Depth range: 10-110 m (Fig. 110). Feeds on ctenophores, salps, medusae and

other zooplankton groups. Table 59 presents a set of growth parameters from Indonesia.

References: 559, 1314, 2047, 3517, 4606, 4789, 5204, 5736, 5756, 6365

Table 58. Length-weight (g/[FL;cm]) relationship of silver pomfret, *Pampus argenteus*, in Indonesia. [Tabel 58. Hubungan panjang-berat (g/[FL; cm]) ikan bawal putih, *Pampus argenteus*, di Indonesia.]

Parameter	Estimate
a	0.1660
s.e.(a)	0.0496
b	2.5033
s.e.(b)	0.1043
r^2	0.9715

Table 59. Growth parameters of silver pomfret, *Pampus argenteus*. [Tabel 59. Parameter pertumbuhan ikan bawal putih, *Pampus argenteus*.]

Parameter	A
L_{∞} (TL, cm)	31.5
K (year ⁻¹)	0.95

A. Java Sea (Central Java) (Ref. 1314)




Fig. 108. (A) Extreme value plot for silver pomfret, *Pampus argenteus*, in Indonesia based on data from *R/Vs Mutiara 4* and *Jurong* showing maxima of 4 length-frequency samples, and estimate of $L_{\max 3} = 30.7 \pm 1.15$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 13 length-frequency samples of silver pomfret, *Pampus argenteus*, from Western Indonesia based on data from *R/Vs Mutiara 4* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 58). Open dots represent outliers, not used for analysis.

[Gambar 108. (A) Gambaran nilai ekstrim ikan bawal putih, *Pampus argenteus*, di Indonesia berdasarkan data dari kapal-kapal penelitian Mutiara 4 dan *Jurong* menunjukkan nilai maksimum 4 contoh frekuensi-panjang, dan angka perkiraan $L_{\max 3} = 30.7 \pm 1.15$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 13 contoh frekuensi-panjang ikan bawal putih, *Pampus argenteus*, dari Indonesia bagian barat berdasarkan data kapal-kapal penelitian Mutiara 4 dan *Jurong* sebagai luaran perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 58). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 109. Distribution of silver pomfret, *Pampus argenteus* based on records of the surveys of *R/Vs Mutiara 4*, *Bawal Putih 2*, *Jurong*, *Lemuru* and *Dr. Fridtjof Nansen*.

[Gambar 109. Penyebaran ikan bawal putih, *Pampus argenteus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, *Bawal Putih 2*, *Jurong*, *Lemuru* dan *Dr. Fridtjof Nansen*.]




Fig. 110. Depth distribution of silver pomfret, *Pampus argenteus*, based on surveys of R/Vs (A) *Mutiara 4* and (B) *Jurong*.
[Gambar 110. Penyebaran kedalaman ikan bawal putih, *Pampus argenteus*, berdasarkan survei kapal-kapal penelitian (A) *Mutiara 4* dan (B) *Jurong*.]

Terapon jarbua (Forsskål, 1775)

Jarbua terapon (English); Kerong-kerong tambu (Indonesian); Djambung, Djandjan, Djangdjan, Kerong-kerong (Java); Djambun, Erong-erong, Kerong-kerong tambu (West Java, Jakarta); Kerongan (Central Java); Djambun, Longkerong (Madura); Keretang (East Sumatra); Kerung-kerung, Mangahua (South Sulawesi, Makassar); Karong-karong (South Sulawesi, Bugis).

Lower opercular spine extending well beyond the opercular flap. Post temporal bone exposed posteriorly and serrate. Body color is fawn above, cream below, nape dark;

head, body and fins with an iridescent sheen. Three or four curved dark brown bands run from the nape to the hind part of the body, the lowermost continuing across the middle of the caudal fin. Dorsal spines: 12-12; soft rays: 10-10; anal spines: 3-3; soft rays: 8-8. $L_{max1} = 33$ cm TL; $L_{max2} = n.a.$; $L_{max3} = 19.7$ cm FL (Fig. 111A). See Fig. 111B and Table 60 for length-weight relationship.

From the Red Sea in the Indian Ocean to Southeast Asia, Indonesia (Fig. 112): north to southern Japan, south to Samoa, Belau in Micronesia and Lord Howe Islands.

Occurs over shallow sandy bottoms, in the vicinity of river mouths. Depth range: 20-290 m (Fig. 113). Feeds on sand-dwelling invertebrates.




Fig. 111. (A) Extreme value plot for Jarbua terapon, *Terapon jarbua*, in Indonesia based on data from R/V *Jurong* showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 19.7 \pm 0.65$ cm FL. (B) Predicted vs. observed weights (in g wet weight) of 8 length-frequency samples of Jarbua terapon, *Terapon jarbua*, from Western Indonesia based on data from R/V *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 60). Open dot represents outlier, not used for analysis.

[Gambar 111. (A) Gambaran nilai ekstrim ikan kerong-kerong tambu, *Terapon jarbua*, di Indonesia berdasarkan data dari kapal penelitian *Jurong* menunjukkan nilai maksimum 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 19.7 \pm 0.65$ cm FL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 8 contoh ikan kerong-kerong tambu, *Terapon jarbua*, dari Indonesia bagian barat berdasarkan data dari kapal penelitian *Jurong* sebagai output perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 60). Bulatan kosong mewakili suatu pengamatan yang tidak dipakai dalam analisis.]

References: 1602, 2857, 3539, 4327, 4515, 4959, 4967, 5213, 5255, 5450, 5525, 5736, 5756, 5970, 6026, 6365

Table 60. Length-weight (g/[FL;cm]) relationship of Jarbua terapon, *Terapon jarbua*, in Indonesia.
 [Tabel 60. Hubungan panjang-berat (g/[FL; cm]) ikan kerong-kerong tambu, Terapon jarbua, di Indonesia.]

Parameter	Estimate
a	0.0748
s.e. (a)	0.0896
b	2.5241
s.e. (b)	0.4443
r ²	0.9824




Fig. 112. Distribution of Jarbua terapon, *Terapon jarbua*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.

[Gambar 112. Penyebaran ikan kerong-kerong tambu, Terapon jarbua, berdasarkan laporan survei dari kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 113. Depth distribution of Jarbua terapon, *Terapon jarbua*, based on surveys of R/Vs (A) Jurong and (B) Bawal Putih 2.

[Gambar 113. Penyebaran kedalaman ikan kerong-kerong tambu, Terapon jarbua, berdasarkan survei kapal-kapal penelitian (A) Jurong dan (B) Bawal Putih 2.]

Trichiurus lepturus (Linnaeus, 1758)

Largehead hairtail (English); Lajur (Indonesian); Djogor (Java); Lajur (West Java, Jakarta); Ladjur (Madura); Ladjuru (South Sulawesi, Makassar).

Body extremely elongate, compressed and tapering to a point. Mouth large with a dermal process at the tip of each jaw. Dorsal fin relatively high; anal fin reduced to minute spinules usually embedded in the skin or slightly breaking through; anterior margin of pectoral fin spine not serrated. Pelvic and caudal fins absent. Lateral line beginning at the upper margin of the gill cover, running oblique to behind the tip of the pectoral fins, then straight close to the ventral contour. Fresh specimens steely blue with silvery reflections, becoming uniformly silvery gray sometime after death. Dorsal spines: 3-3; soft rays: 130-135; anal spines: -; soft rays: 100-105. $L_{\max 1} = 213$ cm TL; $L_{\max 2} = n.a.$; $L_{\max 3} = 125.8$ cm TL (Fig. 114A) and Table 61 for length-weight relationship.

Throughout tropical waters such as Indonesia (Fig. 115) and temperate waters of the world.

Occurs on continental shelf, occasionally in shallow waters and at surface at night. Depth range: 55-385 m (Fig. 116). Immature fish feed mostly on euphausiids, small pelagic

planktonic crustaceans and small fishes while adults feed on anchovies, sardines, myctophiids etc. and occasionally on squid and crustaceans. Adults and juveniles have opposing complementary vertical diurnal feeding migrations.

References: 171, 181, 245, 276, 312, 559, 591, 637, 1263, 1348, 1349, 1350, 1351, 1652, 1751, 1809, 2221, 2302, 2308, 2311, 2682, 2857, 3136, 3383, 3397, 3669, 3670, 3678, 4604, 4733, 4743, 4789, 4830, 4868, 4883, 4931, 5204, 5213, 5217, 5219, 5252, 5287, 5516, 5525, 5541, 5756, 6181, 6365, 6490

Table 61. Length-weight (g/[TL;cm]) relationship of largehead hairtail, *Trichiurus lepturus*, in Indonesia. [Tabel 61. Hubungan panjang-berat (g/[TL;cm]) ikan layur, *Trichiurus lepturus*, di Indonesia.]

Parameter	Estimate
a	0.0009
s.e. (a)	0.0014
b	2.9686
s.e. (b)	0.2967
r^2	0.9019




Fig. 114. (A) Extreme value plot for largehead hairtail, *Trichiurus lepturus*, in Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* showing maxima of 4 length-frequency samples, and estimate of $L_{\max 3} = 125.8 \pm 6.0$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 17 length-frequency samples of largehead hairtail, *Trichiurus lepturus*, from Western Indonesia based on data from R/Vs *Mutiara 4* and *Jurong* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 61). Open dots represent outliers, not used for analysis.

[Gambar 114. (A) Gambaran nilai ekstrim ikan layur, *Trichiurus lepturus*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* menunjukkan nilai maksimum 4 contoh frekuensi-panjang, dan angka perkiraan $L_{\max 3} = 125.8 \pm 6.0$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) 17 contoh frekuensi-panjang ikan layur, *Trichiurus lepturus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4* dan *Jurong* sebagai luaran perangkat lunak ABee (lihat Boks 1), dan yang memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 61). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 115. Distribution of largehead hairtail, *Trichiurus lepturus*, based on records of the surveys of R/Vs Mutiara 4, Bawal Putih 2, Jurong and Dr. Fridtjof Nansen.
 [Gambar 115. Penyebaran ikan layur, *Trichiurus lepturus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan Dr. Fridtjof Nansen.]




Fig. 116. Depth distribution of largehead hairtail, *Trichiurus lepturus*, based on surveys of R/Vs (A) Dr. Fridtjof Nansen and (B) Jurong.
 [Gambar 116. Penyebaran kedalaman ikan layur, *Trichiurus lepturus*, berdasarkan survei kapal-kapal penelitian (A) Dr. Fridtjof Nansen dan (B) Jurong.]

Abalistes stellatus (Lacepède, 1798)

Starry triggerfish (English); Kambing-kambing (Indonesian).

Scales enlarged above the pectoral fin base and just behind the gill slit to form a flexible tympanum; scales of posterior body with prominent keels, forming longitudinal ridges. A prominent groove in the skin extending anteriorly from front of eye for a distance of about 1 eye diameter. Caudal peduncle depressed. Caudal fin rays of adults prolonged above and below. Dorsal spines: 3-3; soft rays: 25-27; anal spines: 0-0; soft rays: 24-26. $L_{max1} = 60$ cm; $L_{max2} = n.a.$; $L_{max3} = 51.5$ cm TL (Fig. 117A). See Fig. 117B and Table 62 for length-weight relationship.

Indo-West Pacific, from East Africa and the Red Sea, Southeast Asia, Indonesia (Fig.118) and thence to Northern Australia and Japan; also reported from the eastern tropical Atlantic.

Inhabits coastal areas, usually found over muddy and sandy bottoms, also around reefs, together with the sponges and algae. Depth range: 20-170 m (Fig. 119). Feeds on benthic animals.

References: 28, 182, 2683, 2857, 3109, 3128, 3804, 3807, 4789, 5193, 5213, 5255, 5450, 5736, 5756, 6026, 6365, 6567

Table 62. Length-weight (g/[TL;cm]) relationship of starry triggerfish, *Abalistes stellatus*, in Indonesia. *Tabel 62. Hubungan panjang-berat (g/[TL;cm]) ikan kambing-kambing, Abalistes stellatus, di Indonesia.*

Parameter	Estimate
a	0.0281
s.e. (a)	0.0085
b	2.8746
s.e. (b)	0.0845
r ²	0.9877




Fig. 117. (A) Extreme value plot for starry triggerfish, *Abalistes stellatus*, in Indonesia based on data from R/Vs *Bawal Putih 2* and *Jurong* showing maxima of 5 length-frequency samples, and estimate of $L_{max3} = 51.5 \pm 1.25$ cm TL. (B) Predicted vs. observed weights (in g wet weight) of 31 length-frequency samples of starry triggerfish, *Abalistes stellatus*, from Western Indonesia based on data from R/Vs *Mutiara 4*, *Jurong* and *Bawal Putih 2* as output by the ABee software (see Box 1), and allowing estimation of a length-weight relationship (see Table 62). Open dot(s) represent outlier(s), not used for analysis. [Gambar 117. (A) Gambaran nilai ekstrim ikan kambing-kambing, *Abalistes stellatus*, di Indonesia berdasarkan data dari kapal-kapal penelitian *Bawal Putih 2* dan *Jurong* menunjukkan nilai maksimum 5 contoh frekuensi-panjang, dan angka perkiraan $L_{max3} = 51.5 \pm 1.25$ cm TL. (B) Berat prediksi terhadap berat observasi (dalam g berat basah) dari 31 contoh frekuensi-panjang ikan kambing-kambing, *Abalistes stellatus*, dari Indonesia bagian barat berdasarkan data dari kapal-kapal penelitian *Mutiara 4*, *Jurong* dan *Bawal Putih 2* sebagai luaran perangkat lunak ABee (lihat Box 1), dan memungkinkan estimasi suatu hubungan panjang-berat (lihat Tabel 62). Bulatan-bulatan kosong mewakili pengamatan-pengamatan yang tidak dipakai dalam analisis.]




Fig. 118. Distribution of starry triggerfish, *Abalistes stellatus*, based on records of the surveys of *R/Vs* Mutiara 4, Bawal Putih 2, Jurong and *Dr. Fridtjof Nansen*.

[Gambar 118. Penyebaran ikan kambing-kambing, *Abalistes stellatus*, berdasarkan laporan survei kapal-kapal penelitian Mutiara 4, Bawal Putih 2, Jurong dan *Dr. Fridtjof Nansen*.]




Fig. 119. Depth distribution of starry triggerfish, *Abalistes stellatus*, based on surveys of *R/Vs* (A) *Dr. Fridtjof Nansen*, (B) *Mutiara 4*, (C) *Jurong* and (D) *Bawal Putih 2*.

[Gambar 119. Penyebaran kedalaman ikan kambing-kambing, *Abalistes stellatus*, berdasarkan survei kapal-kapal penelitian (A) *Dr. Fridtjof Nansen*, (B) *Mutiara 4*, (C) *Jurong* dan (D) *Bawal Putih 2*.]

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

Elasmobranchii (sharks and rays)

Heterodontiformes (bullhead and horn sharks)

Heterodontidae

Heterodontus zebra
 (Gray 1831)

Bullhead, horn, or Port Jackson sharks

Zebra bullhead shark, (M, Dan), Ref. 247. Max. 125 cm TL. From Sulawesi and Ambon. Also Ref.: 559, 5978.

Orectolobiformes (carpet sharks)

Ginglymostomatidae

Nebrius ferrugineus
 (Lesson 1830)

Nurse sharks

Tawny nurse shark, (M, Fi, Sport, Dan), Ref. 247. Max. 320 cm TL. Also Ref.: 5978.

Hemiscylliidae

Chiloscyllium griseum
 Müller & Henle 1838

Chiloscyllium indicum
 (Gmelin 1789)

Chiloscyllium plagiosum
 (Bennett 1830)

Chiloscyllium punctatum
 Müller & Henle 1838

Hemiscyllium freycineti
 (Quoy & Gaimard 1824)

Hemiscyllium hallstromi
 Whitley 1967

Hemiscyllium ocellatum
 (Bonnaterre 1788)

Hemiscyllium strahani
 1967

Hemiscyllium trispeculare
 Richardson 1843

Bamboo sharks

Grey bambooshark, (M, Br, Fi), Ref. 247. Max. 74 cm TL. Also Ref.: 5978.

Slender bambooshark, (M, Fi), Ref. 247. Max. 65 cm TL. Also Ref.: 5978.

Whitespotted bambooshark, (M, Fi), Ref. 247. Max. 69 cm TL. Also Ref.: 5978.

Brownbanded bambooshark, (M, Fi), Ref. 247. Max. 104 cm TL. Also Ref.: 5978.

Indonesia speckled carpetshark, (M), Ref. 247. Max. 46 cm TL. Museum. MNHN A.7792 (Syntypes, Waigiu (Waigeo), Indonesia). Also Ref.: 5978.

Papuan epaulette shark, (M), Ref. 247. Max. 75 cm TL. Also Ref.: 5978.

Epaulette shark, (M), Ref. 247. Max. 107 cm TL. Also Ref.: 6445.

Hooded carpetshark, (M), Ref. 247. Max. 75 cm TL. Also Ref.: 5978.

Speckled carpetshark, (M), Ref. 5978. Max. 64 cm TL.

Orectolobidae

Eucrossorhinus dasypogon
 (Bleeker 1867)

Orectolobus ornatus
 (De Vis 1883)

Rhincodontidae

Rhincodon typus
 Smith 1828

Stegostomatidae

Stegostoma fasciatum
 (Hermann 1783)

Carchariniformes (ground sharks)

Carcharhinidae

Carcharhinus albimarginatus
 (Rüppell 1837)

Carcharhinus amblyrhynchoides
 (Whitley 1934)

Carcharhinus amblyrhynchos
 (Bleeker 1856)

Carcharhinus amboinensis
 (Müller & Henle 1839)

Carcharhinus brevipinna
 (Müller & Henle 1839)

Carpet or nurse sharks

Tasselled wobbegong, (M, Fi, Dan), Ref. 247. Max. 366 cm TL. Museum. BMNH 1867.11.28.209 (Syntypes, Waigiu and Aru, Indonesia).

Ornate wobbegong, (M, Fi, Dan), Ref. 247. Max. 288 cm TL.

Whale shark

Whale shark, (M, Fi, Thr), Ref. 247. Max. 1370 cm TL. Also Ref.: 5978.

Zebra sharks

Zebra shark, (M, Fi), Ref. 247. Max. 354 cm TL.

Requiem sharks

Silvertip shark, (M, Fi, Sport, Dan), Ref. 244. Max. 300 cm TL.

Graceful shark, (M, Fi, Dan), Ref. 244. Max. 140 cm TL. Also Ref.: 5978.

Grey reef shark, (M, Fi, Dan), Ref. 244. Max. 255 cm TL. Also Ref.: 5978.

Pigeye shark, (M, Fi, Dan), Ref. 244. Max. 280 cm TL. Also Ref.: 5978.

Spinner shark, (M, Fi, Sport), Ref. 244. Also Ref.: 5978.

Carcharhinus dussumieri
 (Valenciennes 1839)

Carcharhinus falciformis
 (Bibron 1839)

Carcharhinus hemiodon
 (Valenciennes 1839)

Carcharhinus leucas
 (Valenciennes 1839)

Carcharhinus limbatus
 (Valenciennes 1839)

Carcharhinus longimanus
 (Poey 1861)

Carcharhinus macloti
 (Müller & Henle 1839)

Carcharhinus melanopterus
 (Quoy & Gaimard 1824)

Carcharhinus plumbeus
 (Nardo 1827)

Carcharhinus sealei
 (Pietschmann 1913)

Carcharhinus sorrah
 (Valenciennes 1839)

Galeocerdo cuvier
 (Péron & Lesueur 1822)

Glyphis glyphis
 (Müller & Henle 1839)

Lamiopsis temmincki
 (Müller & Henle 1839)

Loxodon macrorhinus
 Müller & Henle 1839

Negaprion acutidens
 (Rüppell 1837)

Prionace glauca
 (Linnaeus 1758)

Rhizoprionodon acutus
 (Rüppell 1837)

Rhizoprionodon oligolinx
 Springer 1964

Scoliodon laticaudus
 Müller & Henle 1838

Triaenodon obesus
 (Rüppell 1837)

Hemigaleidae

Chaenogaleus macrostoma
 (Bleeker 1852)

Hemigaleus microstoma
 Bleeker 1852

Proscylliidae

Proscyllium habereri
 Hilgendorf 1904

Scyliorhinidae

Apristurus sibogae
 (Weber 1913)

Table 2. Continuation.
 [Tabel 2. Sambungan.]

<i>Apristurus spongiceps</i> (Gilbert 1895)	Spongehead catshark, (M), Ref. 244. Max. 50 cm TL.	<i>Pristis pectinata</i> Latham 1794	Sma
<i>Apristurus verweyi</i> (Fowler 1934)	Borneo catshark, (M, En), Ref. 244. Max. 30 cm TL.	<i>Pristis zijsron</i> Bleeker 1851	Long M.
<i>Atelomycterus marmoratus</i> (Bennett 1830)	Coral catshark, (M, Fi), Ref. 244. Max. 70 cm TL. Also Ref.: 5978.		fre Al
<i>Halaaelurus boesemani</i> Springer & D'Aubrey 1972	Speckled catshark, (M), Ref. 244. Max. 48 cm TL. Ambon Island.	Torpediniformes (electric rays)	
<i>Scyliorhinus garmani</i> (Fowler 1934)	Brownspotted catshark, (M), Ref. 244. Max. 36 cm TL.	Narkidae	
Sphyrnidae	Hammerhead, bonnethead, scoophead shark	<i>Narcine indica</i> Henle 1834	(M, E Re
<i>Eusphyra blochii</i> (Cuvier 1816)	Winghead shark, (M, Fi), Ref. 244 Max. 152 cm TL. From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Narcine timlei</i> (Bloch & Schneider 1801)	Spot Kr
<i>Sphyrna lewini</i> (Griffith & Smith 1834)	Scalloped hammerhead, (M, Br, Fi, Sport, Dan), Ref. 244. Max. 420 cm TL. From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Narke dipterygia</i> (Bloch & Schneider 1801)	Num Kr
<i>Sphyrna mokarran</i> (Rüppell 1837)	Great hammerhead, (M, Dan), Ref. 5978. Max. 610 cm TL.	Rajiformes (skates and rays)	
Lamniformes (mackerel sharks)		Anacanthobatidae	Sm
Alopiidae	Thresher sharks	<i>Anacanthobatis borneensis</i> Chan 1965	(M),
<i>Alopias vulpinus</i> (Bonnaterre 1788)	Thresher shark, (M, Fi, Dan), Ref. 247. Max. 609 cm TL. Also Ref.: 5978.	Rajidae	Skat
Lamnidae	Mackerel sharks, white sharks	<i>Bathyraja andriashevi</i> Dolganev 1985	(M),
<i>Isurus oxyrinchus</i> Rafinesque 1810	Shortfin mako, (M, Fi, Sport, Dan), Ref. 247. Max. 400 cm TL. Also Ref.: 5978.	<i>Bathyraja tzinovskii</i> Dolganev 1985	(M),
Odontaspidae	Sand tigers	<i>Gurgesiella sibogae</i> (Weber 1913)	(M),
<i>Carcharias taurus</i> Rafinesque 1810	Sandtiger shark, (M, Dan), Ref. 5978. Max. 320 cm.	<i>Raja annandalei</i> Weber 1913	(M),
<i>Pseudocarcharias kamoharai</i> (Matsubara 1936)	Crocodile shark, (M), Ref. 5978. Max. 110 cm TL.		
Hexanchiformes (frill and cow sharks)		Rhinobatidae	Guitt
Hexanchidae	Cow sharks	<i>Rhina ancylostoma</i> Bloch & Schneider 1801	Bow Ma (T Re
<i>Hepttranchias perlo</i> (Bonnaterre 1788)	Sharpnose sevengill shark, (M, Fi, Dan), Ref. 247. Max. 137 cm TL. Also Ref.: 5978.	<i>Rhinobatos granulatus</i> (Cuvier 1829)	Gran Po
<i>Hexanchus griseus</i> (Bonnaterre 1788)	Bluntnose sixgill shark, (M, Fi, Sport, Dan), Ref. 247. Max. 500 cm. Also Ref.: 5978.	<i>Rhinobatos halavi</i> (Forsskål 1775)	Hal co
Squaliformes (bramble, sleeper and dogfish sharks)		<i>Rhinobatos obtusus</i> (Müller & Henle 1841)	(M),
Squalidae	Dogfish sharks	<i>Rhinobatos schlegelii</i> (Müller & Henle 1841)	Brow
<i>Centrophorus moluccensis</i> Bleeker 1860	Smallfin gulper shark, (M, Fi), Ref. 247. Max. 100 cm. Museum. Rijksmuseum van Natuurlijke Histoire RMNH 7415 (Holotype, Ambon).	<i>Rhinobatos thouin</i> Anon. 1798	Thou
<i>Etmopterus lucifer</i> Jordan & Snyder 1902	Blackbelly lanternshark, (M), Ref. 247. Max. 42 cm TL. Also Ref.: 5978.	<i>Rhinobatos typus</i> (Bennett 1830)	Gian Ma Su
Pristiformes (sawfishes)		<i>Rhynchobatus australiae</i> Whitley 1939	(M), In
Pristidae	Sawfishes	<i>Rhynchobatus djiddensis</i> (Forsskål 1775)	Whit TU C (R
<i>Anoxypristis cuspidata</i> (Latham 1794)	Pointed sawfish, (M, Br, Fi, Sport, Dan), Ref. 8630. Max. 470 cm TL. Also Ref.: 5978.		
<i>Pristis microdon</i> Latham 1794	Large-tooth sawfish, (M, Br, Fr, Fi, Dan), Ref. 7050. Max. 600 cm TL. Confirmed records from several major river basins (Ref. 6871). Known from Java, Borneo, and Sumatra (Ref. 9859). In range Ref.: 4429.		

Table 2. Marine and brackishwater fishes of Indonesia .

[Tabel 2. Ikan-ikan laut dan pavau Indonesia.]

Myliobatiformes (eagle rays, stingrays and mantas)	Sting rays	<i>Mobula japonica</i>
Dasyatidae	Estuary stingray, (M, Br), Ref. 8630. Max. 130 cm TL.	(Müller & Henle 1841)
<i>Dasyatis fluviatorum</i>	Blue-spotted stingray, (M, Dan), Ref. 3263. Max. 80 cm TL. Museum. ISH 64/82. From southwest Sumatra to Timor Sea (Ref. 5978).	Myliobatidae
Ogilby 1908	Painted maskray, (M), Ref. 9840. Max. 53 cm TL.	<i>Aetobatus flagellum</i>
<i>Dasyatis kuhlii</i> (Müller & Henle 1841)	Pale-edged stingray, (M, Br, Fi), Ref. 8630. Max. 29 cm WD.	(Bloch & Schneider 1801)
<i>Dasyatis leylandi</i> Last 1987	(M), Ref. 8630	<i>Aetobatus narinari</i>
<i>Dasyatis zugei</i> (Müller & Henle 1841)	Bleeker's whipray, (M, Br, Fi, Dan), Ref. 9840. Max. 105 cm WD.	(Euphrasen 1790)
<i>Himantura alcocki</i> (Annandale 1909)	Sharpnose stingray, (M), Ref. 9840. Max. 200 cm TL.	<i>Aetobatus ocellatus</i>
<i>Himantura bleekeri</i> (Blyth 1860)	Mangrove whipray, (M, Br, Dan), Ref. 8630. Max. 97 cm WD. Known from southern New Guinea and northern Java.	(Kuhl & van Hasselt 1823)
<i>Himantura gerrardi</i> (Gray 1851)	Jenkins whipray, (M, Fi), Ref. 8630. Max. 200 cm TL.	<i>Aetomylaeus maculatus</i>
<i>Himantura granulata</i> (Macleay 1883)	Black-spotted whipray, (M), Ref. 9840. Max. 179 cm TL. Known from southern New Guinea.	(Gray 1834)
<i>Himantura jenkinsii</i> (Annandale 1909)	Honeycomb stingray, (M, Br, Sport, Dan), Ref. 6871. Max. 200 cm WD. Museum: CSIRO CA1245, from Bali Strait to Timor (Ref. 5978).	<i>Aetomylaeus milvus</i>
<i>Himantura toshi</i> Whitley 1939	Leopard whipray, (M, Dan), Ref. 8630. Max. 410 cm TL.	(Valenciennes 1838-1841)
<i>Himantura uarnak</i> (Forsskål 1775)	Dwarf whipray, (M), Ref. 9840. Max. 40 cm TL.	<i>Aetomylaeus nicholfii</i>
<i>Himantura undulata</i> (Bleeker 1852)	Cowtail stingray, (M, Br, Fr, Fi, Dan), Ref. 7050. Max. 180 cm WD. Museum: CSIRO CA1247 (PW101), Max. 40 cm TL. from southwest Sumatra to Bali Strait (Ref. 5978).	(Bloch & Schneider 1801)
<i>Himantura walga</i> (Müller & Henle 1841)	Ribbon-tail stingray, (M, Dan), Ref. 6871. Max. 70 cm TL. Museum: ISH 66/82, CSIRO CA1246 (TGT1552), from Bali Strait to Timor (Ref. 5978).	<i>Aetomylaeus vespertilio</i>
<i>Hypolophus sephen</i> (Forsskål 1775)	Porcupine ray, (M, Br, Dan), Ref. 8630. Max. 300 cm TL.	(Bleeker 1852)
<i>Taeniura lymma</i> (Forsskål 1775)	Butterfly rays	<i>Myliobatis tobijei</i>
<i>Urogymnus asperrimus</i> (Bloch & Schneider 1801)	Australian butterfly ray, (M), Ref. 9918. Max. 56 cm TL. Known from the southern coast of New Guinea. Possibly occurs in the northern coast as well.	(Bleeker 1854)
Gymnuridae	Long-tailed butterfly ray, (M), Ref. 8630. Known from Sumatra, Java, and Borneo (Ref. 9918).	<i>Rhinoptera javanica</i>
<i>Gymnura australis</i> (Ramsay & Ogilby 1886)	(M), Ref. 8630. Known from Java (Ref. 9918).	Müller & Henle 1841
<i>Gymnura poecilura</i> (Shaw 1804)	Manta rays and devil rays	Urolophidae
<i>Gymnura zonura</i> (Bleeker 1852)	Giant manta, (M, Br, Dan), Ref. 5978. Max. 700 cm WD.	<i>Trygonoptera javanica</i>
Mobulidae	Devil ray, (M), Ref. 5978. Max. 178 cm WD.	Martens 1864
<i>Manta birostris</i> (Walbaum 1792)		<i>Trygonoptera kaiana</i>
<i>Mobula diabolus</i> (Shaw 1804)		(Günther 1880)
		Actinopterygii (ray-finned fishes)
		Elopiiformes (tarpons and tenpounders)
		Elopidae
		<i>Elops hawaiiensis</i>
		Regan 1909
		<i>Elops machnata</i>
		(Forsskål 1775)
		Megalopidae
		<i>Megalops cyprinoides</i>
		(Broussonet 1782)
		Albuliformes (bonefishes)
		Albulidae
		<i>Albula glossodonta</i>
		(Forsskål 1775)
		<i>Albula neoguinaica</i>
		Valenciennes 1847

Table 2. Continuation.
[Tabel 2. Sambungan.]

<p><i>Anguilla celebensis</i> Kaup 1856</p>	<p>Celebes longfin eel, (M, Br, Fr, Fi), Ref. 7050. Max. 150 cm TL. Sulawesi, Bali, Moluccas, and Irian Jaya.</p>	<p><i>Muraenesox cinereus</i> (Forsskal 1775)</p>
<p><i>Anguilla marmorata</i> Quoy & Gaimard 1824</p>	<p>Giant mottled eel, (M, Br, Fr, Fi, Aq, Sport), Ref. 7050. Max. 200 cm TL.</p>	<p>Muraenidae</p>
<p><i>Anguilla nebulosa</i> McClelland 1844</p>	<p>Mottled eel, (M, Br, Fr), Ref. 7050. Max. 121 cm TL. Known from Sumatra.</p>	<p><i>Echidna nebulosa</i> (Ahl 1789)</p>
<p>Congridae</p>	<p>Conger eels</p>	<p><i>Echidna rhodochilus</i> Bleeker 1863</p>
<p><i>Ariosoma anago</i> (Temminck & Schlegel 1846)</p>	<p>(M), Ref. 5978. Max. 60 cm TL. Museum: BMNH 1984.1.1.3. From southwest Sumatra to Timor Sea.</p>	<p><i>Gymnothorax buroensis</i> (Bleeker 1857)</p>
<p><i>Ariosoma anagoides</i> (Bleeker 1854)</p>	<p>(M), Ref. 559. Max. 51 cm TL.</p>	<p><i>Gymnothorax chilospilus</i> Bleeker 1865</p>
<p><i>Ariosoma scheelei</i> (Stromman 1896)</p>	<p>Tropical conger, (M), Ref. 5323. Max. 20 cm TL.</p>	<p><i>Gymnothorax enigmaticus</i> McCosker & Randall 1982</p>
<p><i>Bathycongrus guttulatus</i> (Günther 1887)</p>	<p>(M), Ref. 5978. Museum: BMNH 1984.1.1.5. From southwest Sumatra to Bali Strait.</p>	<p><i>Gymnothorax fimbriatus</i> (Bennett 1832)</p>
<p><i>Bathymyrus smithi</i> Castle 1968</p>	<p>Maputo conger, (M, Br, Fr), Ref. 5978. Max. 58 cm TL. Museum: BMNH 1984.1.1.4 (PJPW135). From southwest Sumatra to Bali Strait.</p>	<p><i>Gymnothorax javanicus</i> (Bleeker 1859)</p>
<p><i>Bathyroconger vicinus</i> (Vaillant 1888)</p>	<p>(M), Ref. 4453. Max. 88 cm TL.</p>	<p><i>Gymnothorax polyuranodon</i> (Bleeker 1853)</p>
<p><i>Conger cinereus</i> Rüppell 1828</p>	<p>Longfin African conger, (M), Ref. 583. Max. 130 cm.</p>	<p><i>Gymnothorax pseudothyrsoides</i> (Bleeker 1852)</p>
<p><i>Heteroconger hassi</i> (Klausewitz & Eibl-Eibesfeldt 1959)</p>	<p>Spotted garden-eel, Garden eel, (M), Ref. 8631. Max. 35 cm TL. Known from Bali.</p>	<p><i>Gymnothorax reticularis</i> Bloch 1795</p>
<p><i>Heteroconger perissodon</i> Böhle & Randall 1981</p>	<p>(M), Ref. 8912. Museum: Molucca Is., Ambon, Poka, ANSP 142731 BPBM 18543, CAS 45889, MNHN 1980-1190, USNM 221380, WAM P26788-001.</p>	<p><i>Gymnothorax richardsoni</i> (Bleeker 1852)</p>
<p><i>Macrocephenchelys brachialis</i> Fowler 1934</p>	<p>(M, Thr), Ref. 245.</p>	<p><i>Gymnothorax tile</i> (Hamilton 1822)</p>
<p><i>Parabathymyrus macrophthalmus</i> Kamohara 1938</p>	<p>(M), Ref. 5978. Max. 47 cm TL. From Bali Strait to Timor Sea.</p>	<p><i>Gymnothorax undulatus</i> (Lacepede 1803)</p>
<p><i>Rhynchoconger brevirostris</i> Chen & Weng 1967</p>	<p>(M), Ref. 5978. Museum: NMNZ 15178 (TGT1894). From Bali Strait to Timor Sea.</p>	<p><i>Gymnothorax zonipectis</i> Seale 1906</p>
<p><i>Uroconger lepturus</i> (Richardson 1845)</p>	<p>Slender conger, (M), Ref. 5978. Max. 52 cm TL. Museum: NTM S.10751-001 (TGT1677). From southwest Sumatra to Timor Sea.</p>	<p><i>Pseudechidna brummeri</i> (Bleeker 1858-59)</p>
<p>Moringuidae</p>	<p>Worm or spaghetti eels</p>	<p><i>Rhinomuraena quaesita</i> Garman 1888</p>
<p><i>Moringua javanica</i> (Kaup 1856)</p>	<p>Java spaghetti eel, (M, Br), Ref. 5501. Max. 90 cm TL. Recorded from Java and Sulawesi. Also Ref.: 7050.</p>	<p><i>Siderea picta</i> (Ahl 1789)</p>
<p><i>Moringua microchir</i> Bleeker 1853</p>	<p>(M), Ref. 7050. Max. 30 cm TL.</p>	<p><i>Strophidon sathete</i> (Hamilton 1822)</p>
<p>Muraenesocidae</p>	<p>Pike congers</p>	<p>Nemichthyidae</p>
<p><i>Congresox talabon</i> (Cuvier 1829)</p>	<p>Yellow pike conger, (M, Br, Fi), Ref. 7238. Max. 80 cm TL. Recorded from Sulawesi and Lesser Sundas. Also Ref.: 7050.</p>	<p><i>Avocettina infans</i> (Günther 1878)</p>
<p><i>Congresox talabonoides</i> (Bleeker 1853)</p>	<p>Indian pike conger, (M, Br, Fi), Ref. 7050. Max. 250 cm TL. Recorded from Sulawesi.</p>	<p><i>Nemichthys scolopaceus</i> Richardson 1848</p>
<p><i>Gavialiceps taeniola</i> Alcock 1889</p>	<p>(M), Ref. 5978. Max. 84 cm TL. Museum: BMNH 1984.1.1.6, 1984.1.1.7 (PJPW136). From southwest Sumatra to Bali Strait.</p>	<p>Nettastomatidae</p>
<p><i>Muraenesox bagio</i> (Hamilton 1822)</p>	<p>Common pike conger, (M, Br, Fi, Dan), Ref. 5978. Max. 200 cm TL. Museum: CSIRO CA1088. From southwest Sumatra to Bali Strait. Also Ref.: 7050.</p>	<p><i>Nettenchelys gephyra</i> Castle & Smith 1981</p>

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

226	<p>Ophichthidae <i>Apterichtus klazingai</i> (Weber 1913) <i>Callechelys catostomus</i> (Schneider & Forster 1801) <i>Lamnostoma mindora</i> (Jordan & Richardson 1908) <i>Lamnostoma orientalis</i> (McClelland 1844) <i>Muraenichthys gymnopterus</i> (Bleeker 1853) <i>Muraenichthys macropterus</i> Bleeker 1857 <i>Myrichthys bleekeri</i> Gosline 1951 <i>Ophichthus bonaparti</i> (Kaup 1856) <i>Ophichthus macrochir</i> (Bleeker 1853) <i>Ophichthus urolophus</i> (Temminck & Schlegel 1846) <i>Pisodonophis boro</i> (Hamilton 1822) <i>Pisodonophis cancrivorus</i> (Richardson 1848) <i>Pisodonophis hypselopterus</i> (Bleeker 1851) Synaphobranchidae <i>Dysomma anguillare</i> Barnard 1923 Clupeiformes (herrings) Chirocentridae <i>Chirocentrus dorab</i> (Forsskål 1775) <i>Chirocentrus nudus</i> (Swainson 1839) Clupeidae <i>Amblygaster clupeioides</i> Bleeker 1849 <i>Amblygaster leiogaster</i> (Valenciennes 1847) <i>Amblygaster sirm</i> (Walbaum 1792)</p>	<p>Snake eels Sharpnose snake eel, (M), Ref. 3972. Max. 40 cm TL. (M), Ref. 2334. Max. 85 cm TL. (M, Br), Ref. 7050. Max. 40 cm TL. Expected to be present in the area (Ref 7050). Oriental worm-eel, (M, Br, Bait), Ref. 7050. Max. 30 cm. Reported from Irian Jaya. (M, Br), Ref. 7050. Max. 26.6 cm TL. Slender snake-eel, (M), Ref. 637. Max. 25 cm TL. (M), Ref. 1602. Max. 39.5 cm TL. (M), Ref. 3972. Max. 75 cm TL. (M, Br), Ref. 7050. Max. 51 cm TL. Known from Sumatra and Java. (M), Ref. 5978. Max. 60 cm TL. Museum: NTM S.10750-001 (TGT1444). From Bali Strait to Timor Sea. Rice-paddy eel, (M, Br, Fr, Fi), Ref. 7050. Max. 100 cm TL. Known from Sumatra, Java, Sulawesi and Moluccas. Longfin snake-eel, (M, Br, Fr, Fi), Ref. 5978. Max. 100 cm TL. Museum: BMNH 1984.1.1.8 (PJPW63). From southwest Sumatra to Bali Strait Bangka, Sulawesi, and Moluccas (Ref. 7050). (M, Br, Fr), Ref. 7050. Max. 75 cm TL. Recorded from Borneo. Cutthroat eels Shortbelly eel, (M), Ref. 5978. Max. 52 cm TL. Museum: NTM S.10745-001 (TGT1895). From Bali Strait to Timor Sea. Wolf herring Dorab wolf-herring, (M, Fi), Ref. 188. Max. 100 cm SL. Museum: LPPL JIF4 (TGT2222). Found from southwest Sumatra to Timor Sea (Ref. 5978). Whitefin wolf-herring, (M, Fi), Ref. 188. Max. 100 cm SL. Herrings, shads, sardines, menhadens Bleeker smoothbelly sardinella, (M, Fi, Bait), Ref. 188. Max. 17 cm SL. Smoothbelly sardinella, (M, Fi), Ref. 188. Max. 23 cm SL. Found from southwest Sumatra to Bali Strait (Ref. 5978). Spotted sardinella, Sardin, (M, Fi, Bait), Ref. 188. Max. 23 cm SL. Found from southwest Sumatra to Timor Sea. Museum: NTM S.11000-005 (TGT2382) (Ref. 5978).</p>	<p><i>Anodontostoma chacunda</i> (Hamilton 1822) <i>Anodontostoma selangkat</i> (Bleeker 1852) <i>Anodontostoma thailandiae</i> Wongratana 1983 <i>Dussumieria acuta</i> Valenciennes 1847 <i>Dussumieria elopsoidea</i> Bleeker 1849 <i>Escualosa thoracata</i> (Valenciennes 1847) <i>Herklotsichthys dispilonotus</i> (Bleeker 1852) <i>Herklotsichthys gotoi</i> Wongratana 1983 <i>Herklotsichthys quadrimaculatus</i> (Rüppell 1837) <i>Hilsa kelee</i> (Cuvier 1829) <i>Ilisha elongata</i> (Bennett 1830) <i>Ilisha filigera</i> (Valenciennes 1847) <i>Ilisha kampeni</i> (Weber & de Beaufort 1913) <i>Ilisha macrogaster</i> Bleeker 1866 <i>Ilisha megaloptera</i> (Swainson 1839) <i>Ilisha melastoma</i> (Schneider 1801) <i>Ilisha pristigastroides</i> (Bleeker 1852) <i>Ilisha sirishai</i> Seshagiri Rao 1975 <i>Nematalosa come</i> (Richardson 1846) <i>Opisthopterus tardoore</i> (Cuvier 1829)</p>
-----	--	---	---

Table 2. Continuation.
[Tabel 2. Sambungan.]

	<i>Opisthopterus valenciennesi</i> Bleeker 1872	Slender tardoor, (M, Br, Fi), Ref. 188. Max. 20 cm SL. Known from Sumatra, Kalimantan, and Java (Ref. 7050).	Bleeker 1849
	<i>Pellona ditchela</i> Valenciennes 1847	Indian pellona, (M, Br, Fr, Fi, Bait), Ref. 188. Max. 16 cm SL. Also Ref.: 6567, 7050. Museum: NTM S.10734-020 (TGT1226). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Encrasicholina devisi</i> (Whitley 1940)
	<i>Raconda russeliana</i> Gray 1831	Raconda, (M, Br, Fi), Ref. 188. Max. 19 cm SL. Sumatra, Kalimantan, and Java (Ref. 7050).	<i>Encrasicholina heteroloba</i> (Rüppell 1837)
	<i>Sardinella albella</i> (Valenciennes 1847)	White sardinella, (M, Fi), Ref. 188. Max. 14 cm SL.	<i>Encrasicholina punctifer</i> Fowler 1938
	<i>Sardinella atricauda</i> (Günther 1868)	Bleeker's blacktip sardinella, (M, Fi), Ref. 188. Max. 12.6 cm SL.	<i>Engraulis japonicus</i> Temminck & Schlegel 1846
	<i>Sardinella brachysoma</i> Bleeker 1852	Deepbody sardinella, (M, Fi), Ref. 188. Max. 13 cm SL. Museum: NTM S.10733-023 (TGT1228). Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Papuengraulis micropinna</i> Munro 1964
	<i>Sardinella gibbosa</i> (Bleeker 1849)	Goldstripe sardinella, Tembang, (M, Fi), Ref. 188. Max. 17 cm SL. Museum: NTM S.10733-024 (TGT2110). Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Setipinna breviceps</i> (Cantor 1850)
	<i>Sardinella lemuru</i> Bleeker 1853	Bali sardinella, Lemuru, (M, Fi), Ref. 188. Max. 23 cm SL. Strait of Bali, south of Ternate and Djakarta Bay and off Central Java. Fished during rainy season with ca. 15-20 days/mo. active fishing with 'payang besar' or 'djala oras' seine nets. Museum: NTM S.10733-032 (TGT1255) (Ref. 5978).	<i>Setipinna melanochir</i> (Bleeker 1849)
227	<i>Sardinella melanura</i> (Cuvier 1829)	Blacktip sardinella, (M, Fi, Bait), Ref. 188. Max. 12.2 cm SL.	<i>Setipinna taty</i> (Valenciennes 1848)
	<i>Spratelloides delicatulus</i> (Bennett 1831)	Delicate round herring, (M, Fi, Bait), Ref. 188. Max. 7 cm SL.	<i>Setipinna tenuifilis</i> (Valenciennes 1848)
	<i>Spratelloides gracilis</i> (Temminck & Schlegel 1846)	Silverstriped round herring, (M, Fi, Bait), Ref. 188. Max. 9.5 cm SL.	<i>Stolephorus andhraensis</i> Babu Rao 1966
	<i>Spratelloides lewisi</i> Wongratana 1983	Lewis' round herring, (M, Fi, Bait), Ref. 188. Max. 6 cm SL.	<i>Stolephorus baganensis</i> Hardenberg 1933
	<i>Tenualosa macrura</i> (Bleeker 1852)	Longtail shad, (M, Br, Fr, Fi), Ref. 188. Max. 52 cm SL. Also Ref.: 7050.	<i>Stolephorus carpentariae</i> (De Vis 1882)
	<i>Tenualosa toli</i> (Valenciennes 1847)	Toli shad, (M, Br, Fr, Fi), Ref. 188. Max. 50 cm SL. Also Ref.: 7050.	<i>Stolephorus commersonii</i> Lacepède 1803
	Engraulidae	Anchovies	<i>Stolephorus dubiosus</i> Wongratana 1983
	<i>Coilia borneensis</i> Bleeker 1852	Bornean grenadier anchovy, (M, Br, Fr), Ref. 189. Max. 12.4 cm SL. Reported from Sumatra and Kalimantan (Ref. 7050).	<i>Stolephorus indicus</i> (van Hasselt 1823)
	<i>Coilia coomansi</i> Hardenberg 1934	Cooman's grenadier anchovy, (M, Br, Fi), Ref. 189. Max. 12.3 cm SL. Reported from Sumatra and Kalimantan (Ref. 7050).	<i>Stolephorus insularis</i> Hardenberg 1933
	<i>Coilia dussumieri</i> Valenciennes 1848	Goldspotted grenadier anchovy, (M, Br, Fi), Ref. 189. Max. 20 cm SL. Known from Sumatra and Java (Ref. 7050).	<i>Stolephorus tri</i> (Bleeker 1852)
	<i>Coilia macrognathos</i> Bleeker 1852	Longjaw grenadier anchovy, (M, Br, Fi), Ref. 189. Max. 26 cm SL. Known from Kalimantan (Ref. 7050).	<i>Stolephorus waitei</i> Jordan & Seale 1926
	<i>Coilia neglecta</i> Whitehead 1968	Neglected grenadier anchovy, (M, Br, Fi), Ref. 189. Max. 17 cm SL. Known from Kalimantan (Ref. 7050).	<i>Thryssa baelama</i> (Forsskål 1775)
	<i>Coilia rebentischii</i>	Many-fingered grenadier anchovy, (M, Br, Fi), Ref. 189.	<i>Thryssa dussumieri</i> (Valenciennes 1848)
			<i>Thryssa encrasicholoides</i> (Bleeker 1852)
			<i>Thryssa hamiltonii</i> (Gray 1830)

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	<i>Thryssa kammalensis</i> (Bleeker 1849)	Kammal thryssa, (M, Br, Fi), Ref. 189. Max. 8.3 cm SL.	Valenciennes 1840	Max
	<i>Thryssa mystax</i> (Schneider 1801)	Moustached thryssa, (M, Br, Fi), Ref. 189. Max. 15.5 cm SL. Museum: NTM S.10733-038. Found from southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 7050.	<i>Batrachocephalus mino</i> (Hamilton 1822)	Beard
	<i>Thryssa setirostris</i> (Broussonet 1782)	Longjaw thryssa, (M, Br, Fi, Bait), Ref. 189. Max. 18 cm SL. Museum: LPPL JIF3 (TGT2171). Found from southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 7050.	<i>Cephalocassis melanochir</i> (Bleeker 1852)	(M, Br)
			<i>Hemipimelodus borneensis</i> (Bleeker 1851)	(M, Br)
			<i>Ketengus typus</i> Bleeker 1857	(M, Br)
			<i>Netuma thalassina</i> (Rüppell 1837)	Ma
			<i>Osteogeneiosus militaris</i> (Linnaeus 1758)	Jav
				Giant
				Als
				Frc
				Soldie
				TL.
				Eelta
				Long-
				Ma
				White
				Ma
				Gray
				Ha
				Ja
				Stripe
				Me
				Fo
				Alc
				Salmoniformes (salmons, pikes and smelts)
				Aleocephalidae
				Slick
				(M), I
				(M), I
				(P)
				Blunt
				Me
				(P)
				Deep
				(M),
				1E
				Sl
				Microstomatidae
				(M),
				Stomiiformes (lightfishes and dragonfishes)
				Astronesthidae
				Sna
				(M),
				Fr
				(M),
				M
				pe
				Fi
				(M),
				B

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Astronesthes martensii</i> Klunzinger 1871	From southwest Sumatra to Bali Strait. (M), Ref. 5978. Max. 14 cm TL. Museum: BMNH 1984.1.1.22 (TGT (PJPW) 774). From southwest Sumatra to Timor Sea.	<i>Stomias nebulosus</i> Alcock 1889	(M), 19 to
<i>Astronesthes splendidus</i> Brauer 1902	(M), Ref. 5978. From southwest Sumatra to Bali Strait.	Aulopiformes (grinners)	Lanc
Chauliodontidae	Viperfishes	Alepisauridae	Long
<i>Chauliodus sloani</i> Schneider 1801	Sloane's viperfish, (M), Ref. 5978. Max. 35 cm SL. Museum: AMS I.24338-001 (TGT1672). From southwest Sumatra to Bali Strait.	<i>Alepisaurus ferox</i> Lowe 1833	Green
Gonostomatidae	Bristlemouths or lightfishes	Chlorophthalmidae	Shor
<i>Diplophos greyae</i> Johnson 1970	(M), Ref. 5978. Museum: BMNH 1984.1.1.17 (TGT (PJPW) 839). From southwest Sumatra to Bali Strait.	<i>Chlorophthalmus agassizi</i> Bonaparte 1840	M Fc 59
<i>Gonostoma elongatum</i> Günther 1878	Elongated bristlemouth fish, (M), Ref. 5978. Max. 27.5 cm TL. Museum: BMNH 1984.1.1.23 (PJPW116), from southwest Sumatra to Bali Strait.	<i>Chlorophthalmus albatrossis</i> Jordan & Starks 1904	(M), 01 St
Malacosteidae	Loosejaws	<i>Chlorophthalmus bicornis</i> Norman 1939	Spin Mu
<i>Malacosteus niger</i> Ayres 1848	(M), Ref. 4469. Max. 21.6 cm SL.	<i>Chlorophthalmus nigromarginatus</i> Kamohara 1953	(M), 01 Se
Melanostomiidae	Scaleless black dragonfishes	<i>Chlorophthalmus oblongus</i> Kamohara 1953	(M), 01
<i>Echiostoma barbatum</i> Lowe 1843	(M), Ref. 5978. Max. 36.8 cm SL. From southwest Sumatra to Bali Strait.	<i>Ipnops agassizii</i> Garman 1899	(M), Mu
<i>Eustomias bifilis</i> Gibbs 1960	(M), Ref. 5978. From southwest Sumatra to Bali Strait.	Paralepididae	Barra
<i>Melanostomias macrophotus</i> Regan & Trewavas 1930	(M), Ref. 5978 Max. 22.9 cm SL. Museum: BMNH 1984.1.1.25 (PJPW 42 in part). From southwest Sumatra to Bali Strait.	<i>Lestidiops mirabilis</i> (Ege 1933)	Stran Su
<i>Melanostomias valdiviae</i> Brauer 1902	Valdivia black dragon fish, (M), Ref. 4468. Max. 23.2 cm SL. West coast of Sumatra.	<i>Lestidium atlanticum</i> Borodin 1928	Atlan Mu
<i>Photonectes albipennis</i> (Döderlein 1882)	(M), Ref. 5978. Museum: BMNH 1984.1.1.27 (PJPW) 42 in part). From southwest Sumatra to Bali Strait.	<i>Lestrolepis intermedia</i> (Poey 1868)	(M), BM Su
Phosichthyidae	Lightfishes	<i>Stemonosudis elegans</i> (Ege 1933)	(M), 19 Su
<i>Pollichthys maui</i>	(M), Ref. 5978. Max. 6 cm SL. Museum: BMNH 1984.1.1.13 (Poll 1953) (PJPW) 2031). From southwest Sumatra to Bali Strait.	<i>Stemonosudis rothschildi</i> Richards 1967	(M), 19 Ba
<i>Polymetme corythaeola</i> (Alcock 1898)	(M), Ref. 5978. Max. 26 cm SL. Museum: BMNH 1984.1.1.14 (PJPW205). From southwest Sumatra to Bali Strait.	Scopelarchidae	Pear
Sternoptychidae	Brock's bristle-mouth fish, (M), Ref. 5978. Museum: AMS I.24338-002 (TGT1676). From Bali Strait to Timor Sea.	<i>Melamphaes danae</i> Ebeling 1962	(M), to
<i>Argyripnus brocki</i> Struhsaker 1973	Pearlsides, (M, Fi), Ref. 5978. Max. 6.5 cm SL. Museum: BMNH 1984.1.1.19. From southwest Sumatra to Bali Strait.	Synodontidae	Lizan
<i>Maurolicus muelleri</i> (Gmelin 1789)	(M), Ref. 5282.	<i>Harpadon microchir</i> Günther 1878	(M), 19 to
<i>Polyipnus spinosus</i> (Günther 1887)	(M), Ref. 5978. Museum: AMS I.24318-002 (TGT2533). From southwest Sumatra to Bali Strait.	<i>Harpadon nehereus</i> (Hamilton 1822)	Bom Mu
<i>Polyipnus tridentifer</i> McCulloch 1914	(M), Ref. 5978. Museum: AMS I.24338-001 (TGT1672 in part). From Bali Strait to Timor Sea.	<i>Saurida gracilis</i> (Quoy & Gaimard 1824)	Fo Gr
<i>Polyipnus triphanos</i> Schultz 1938	Scaly dragonfishes		
Stomiidae			

Table 2. Marine and brackishwater fishes of Indonesia

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Saurida longimanus</i> Norman 1939	Longfin lizardfish, (M), Ref. 5978. Max. 25 cm. Museum: NTM S.10761-005 (TGT3164). From Bali Strait to Timor Sea.	Cressey 1981 <i>Synodus variegatus</i> (Lacepède 1803)	Mu: F Var
<i>Saurida micropectoralis</i> Shindo & Yamada 1972	Shortfin lizardfish, (M), Ref. 2117. Max. 38 cm. Museum: BMNH 1984.1.1.33 (TGT1325). Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Trachinocephalus myops</i> (Forster 1801)	(S Sne
<i>Saurida nebulosa</i> Valenciennes 1849	Clouded lizardfish, (M, Br, Fi), Ref. 5978. Max. 18.5 cm FL. Museum: NTM S.10749-001 (TGT1805). From Bali Strait to Timor Sea.	Myctophiformes (lanternfishes)	
<i>Saurida tumbil</i> (Bloch 1795)	Greater lizardfish, (M, Fi), Ref. 6567. Max. 60 cm FL. Museum: LPPL JIF6 (TGT2384). From southwest Sumatra to Timor Sea.	Myctophidae	Lar
<i>Saurida undosquamis</i> (Richardson 1848)	Brushtooth lizardfish, (M, Fi), Ref. 6567. Max. 50 cm SL. Museum: LPPL JIF7 (TGT2248). Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Benthoosema fibulatum</i> (Gilbert & Cramer 1897)	Spil M (
<i>Saurida wanieso</i> Shindo & Yamada 1972	Wanieso lizardfish, (M), Ref. 5978. Max. 65 cm SL. Museum: NTM S.11029-002 (TGT2253). From southwest Sumatra to Bali Strait. In range Ref.: 559.	<i>Benthoosema pterotum</i> (Alcock 1890)	Skir M F
<i>Synodus dermatogenys</i> Fowler 1912	Sand lizardfish, (M), Ref. 8631. Max. 18 cm SL. Known from Flores.	<i>Benthoosema suborbitale</i> (Gilbert 1913)	Sm M
<i>Synodus englemani</i> Schultz 1953	(M), Ref. 5978. Museum: NTM S.10758-001 (TGT1877). From Bali Strait to Timor.	<i>Diaphus chrysorhynchus</i> Gilbert & Cramer 1896	Gol M fi
<i>Synodus hoshinonis</i> Tanaka 1917	Blackear lizardfish, (M, Fi), Ref. 5978. Max. 20 cm. Museum: USNM 264322 (TGT2535). From southwest Sumatra to Bali Strait.	<i>Diaphus coeruleus</i> (Klunzinger 1871)	(M) 2
<i>Synodus indicus</i> (Day 1873)	Indian lizardfish, (M, Fi), Ref. 5978. Max. 20 cm. Museum: USNM 264332 (TGT2602). From southwest Sumatra to Timor Sea.	<i>Diaphus effulgens</i> (Goode & Bean 1896)	(M) I. T
<i>Synodus jaculum</i> Russell & Cressey 1979	Lighthouse lizardfish, (M, Fi), Ref. 5978. Max. 20 cm TL. Also Ref.: 8631. Museum: NTM S.10744-003. From Bali Strait to Timor Sea.	<i>Diaphus fragilis</i> Tåning 1928	Fra- M S
<i>Synodus kaianus</i> (Günther 1880)	Gunther's lizard fish, (M), Ref. 5978. Max. 30 cm SL. Museum: NTM S.10760-001 (TGT1724). From Bali Strait to Timor Sea.	<i>Diaphus garmani</i> Gilbert 1906	(M) 1 te
<i>Synodus macrocephalus</i> Cressey 1981	(M), Ref. 5978. Museum: USNM 235455 (TGT3092). From southwest Sumatra to Timor Sea.	<i>Diaphus lucidus</i> (Goode & Bean 1896)	(M) 1 S
<i>Synodus macrops</i> Tanaka 1917	Triplecross lizardfish, (M, Fi), Ref. 5978. Max. 20 cm. Museum: USNM 235456 (TGT3215). From Bali Strait to Timor Sea.	<i>Diaphus signatus</i> Gilbert 1908	(M) s 4
<i>Synodus oculateus</i> Cressey 1981	(M), Ref. 5978. Museum: NTM S.10743-002 (TGT3162). From Bali Strait to Timor Sea.	<i>Diaphus splendidus</i> (Brauer 1904)	(M) C
<i>Synodus rubromarmoratus</i> Russell & Cressey 1979	Redmarbled lizardfish, (M), Ref. 5978. Max. 12 cm TL. Museum: USNM 264327 (TGT2281B). From southwest Sumatra to Timor Sea.	<i>Diaphus suborbitalis</i> Weber 1913	(M) 1
<i>Synodus sageneus</i> Waite 1905	Speartoothed grinner, (M, Fi), Ref. 3520. Max. 26 cm TL.	<i>Diaphus thiollieri</i> Fowler 1934	(M) F
<i>Synodus tectus</i> Cressey 1981	(M), Ref. 5978. Museum: NTM S.10740-002 (TGT1632). From Bali Strait to Timor Sea.	<i>Diaphus watasei</i> Jordan & Starks 1904007.	(M) F
<i>Synodus ulae</i> Schultz 1953	Red lizard fish, (M), Ref. 8631. Max. 30 cm SL. Known from Flores.	<i>Lampadena luminosa</i> (Garman 1899)	(M) 1 S
<i>Synodus usitatus</i> (M), Ref. 5978.		<i>Lampanyctus lineatus</i> Tåning 1928	(M) 1 S
		<i>Myctophum asperum</i>	Pric

Table 2. Continuation.
[Tabel 2. Sambungan.]

Richardson 1845	Museum: BMNH 1984.1.1.43TGT(PJPW)436). From southwest Sumatra to Bali Strait.	<i>Ophidion muraenolepis</i> (Günther 1880)	(M)
<i>Myctophum brachygnathum</i> (Bleeker 1856)	Short-jawed lanternfish, (M), Ref. 5978. Museum: BMNH 1984.1.1.44 (PJPW 120 in part). From southwest Sumatra to Timor Sea. RMNH 6932 (syntypes, 2), Makassar.	<i>Sirembo jerdoni</i> (Day 1888)	Bro
<i>Notoscopelus resplendens</i> (Richardson 1845)	Patchwork lampfish, (M), Ref. 5978. Max. 9.5 cm SL. Museum: BMNH 1984.1.1.46 (PJPW 120 in part). From southwest Sumatra to Bali Strait.		
<i>Scopelopsis multipunctatus</i>	(M), Ref. 5978 Brauer 1906. Max. 8.1 cm SL. Museum: BMNH 1984.1.1.47 (PJPW 120 in part). From southwest Sumatra to Bali Strait.		
Lampriformes (velifers, tube-eyes and ribbonfishes)			
Megalomycteridae		Largenose fishes	
<i>Ataxolepis apus</i> Myers & Friehofer 1966	(M), Ref. 9844.		
Trachipteridae		Ribbonfishes	
<i>Desmodema polystictum</i> (Ogilby 1897)	Polka-dot ribbonfish, (M), Ref. 5978. Max. 110 cm TL. Museum: NTM S.11036-001 (TGT1560). From Bali Strait to Timor Sea.		
<i>Trachipterus trachipterus</i> (Gmelin 1789)	Ribbon fish, (M), Ref. 5978. Max. 300 cm TL. Museum: BMNH 1984.1.1.62 (TGT (PJPW) 834). From southwest Sumatra to Bali Strait.		
Veliferidae		Velifers	
<i>Velifer hypselopterus</i> Bleeker 1879	Sailfin velifer, (M), Ref. 5978. Max. 40 cm SL. Museum: WAM P.26194-010. From Bali Strait to Timor Sea, though the lateral line count (59-63) of this material is consistently below that stated for the species (70-72).		
Ophidiiformes (cusk eels)			
Carapidae		Pearlfishes	
<i>Encheliophis gracilis</i> (Bleeker 1856)	Graceful pearlfish, (M), Ref. 4104. Max. 30 cm SL. Banda Islands, type locality.		
<i>Pyramodon ventralis</i> Smith & Radcliffe 1913	(M), Ref. 559		
Ophidiidae		Cusk-eels	
<i>Glyptophidium macropus</i> Alcock 1894	(M), Ref. 5978. Museum: ZMUC P77737 (TGT1448). From Bali Strait to Timor Sea.		
<i>Glyptophidium oceanicum</i> Smith & Radcliffe 1913	(M), Ref. 5978. Max. 22 cm SL. Museum: ZMUC P77751 (TGT2518). From southwest Sumatra to Bali Strait.		
<i>Homostolus acer</i> Smith & Radcliffe 1913	(M), Ref. 6225. Max. 18.4 cm SL. Museum: BSKU 16704, 16678; MNHN 1984-640.		
<i>Hypopleuron caninum</i> Smith & Radcliffe 1913	(M), Ref. 5978. Museum: ZMUC P77752 (TGT2519). From southwest Sumatra to Bali Strait.		
<i>Lamprogrammus brunswigi</i> (Brauer 1906)	(M), Ref. 3686. Max. 90 cm. Museum: USNM 74146 (Holotype, Buton Strait, Indonesia).		
<i>Neobythites longipes</i> Smith & Radcliffe 1913	(M), Ref. 5978. Museum: ZMUC P77740 (TGT1718). From Bali Strait to Timor Sea.		
<i>Neobythites macrops</i> Günther 1887	(M), Ref. 5978. Museum: BMNH 1983.1.1.57 (TGT (PJPW) 453). From southwest Sumatra to Bali Strait.		
<i>Neobythites malayanus</i> Weber 1913	(M), Ref. 5978. Museum: ZMUC P77742 (TGT1897). From Bali		
		Gadiformes (cods)	
		Bregmacerotidae	Co
		<i>Bregmaceros atlanticus</i> Goode & Bean 1886	An
		<i>Bregmaceros lanceolatus</i> Shen 1960	(M)
		<i>Bregmaceros maclellandi</i> Thompson 1840	Sp
		<i>Bregmaceros nectabanus</i> Whitley 1941	(M)
		Macrouridae	Gr
		<i>Caelorinchus argentatus</i> Smith & Radcliffe 1912	Sil
		<i>Caelorinchus argus</i> Weber 1913	Ey
		<i>Cetonurus globiceps</i> (Vaillant 1888)	(M)
		<i>Malacocephalus laevis</i> (Lowe 1843)	So
		<i>Ventrifossa divergens</i> Gilbert & Hubbs 1920	Pla
		<i>Ventrifossa nigrodorsalis</i> Gilbert & Hubbs 1920	Sp
		<i>Ventrifossa petersoni</i> (Alcock 1891)	Pe
		Moridae	Moric
		<i>Physiculus nigrescens</i> Smith & Radcliffe 1912	(M)
		<i>Tripterophycis gilchristi</i> Boulenger 1902	Gr
		Batrachoidiformes (toadfishes)	
		Batrachoididae	To
		<i>Batrachomoeus trispinosus</i> (Günther 1861)	Three
		<i>Halophryne diemensis</i> (LeSueur 1824)	Kn Band
		Lophiiformes (anglerfishes)	
		Antennariidae	Frog
		<i>Antennarius biocellatus</i> (Cuvier 1817)	Brack
		<i>Antennarius coccineus</i>	Ma Scarl

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

(Lesson 1830)			<i>Tetrabrachium ocellatum</i>	(M), F
<i>Antennarius commersoni</i>	Commerson's frogfish, (M), Ref. 5978. Max. 29.1 cm SL.		Günther 1880	
(Latreille 1804)	Museum: LPPL JIF10 (TGT1770). From Bali Strait to Timor Sea.		Atheriniformes (silversides)	
<i>Antennarius dorehensis</i>	New Guinean frogfish, (M), Ref. 6773. Max. 5.1 cm SL.		Atherinidae	Silve
Bleeker 1859			<i>Atherinomorus cylindricus</i>	Waige
<i>Antennarius hispidus</i>	Shaggy angler, (M), Ref. 5978.		(Valenciennes 1835)	For
(Bloch & Schneider 1801)	Max. 20 cm TL. Museum: CSIRO CA1689.		<i>Atherinomorus duodecimalis</i>	Tropic
	From Bali Strait to Timor Sea.		(Valenciennes 1835)	Kn
<i>Antennarius maculatus</i>	Warty frogfish, (M), Ref. 6773. Max. 9 cm.		<i>Atherinomorus lineatus</i>	Line s
(Desjardins 1840)			(Günther 1872)	Re
<i>Antennarius nummifer</i>	Spotfin frogfish, (M), Ref. 6773. Max. 10 cm SL.		<i>Atherion elymus</i>	Beard
(Cuvier 1817)			Jordan & Starks 1901	
<i>Antennarius pictus</i>	Painted angler, (M), Ref. 5978. Max. 30 cm TL.		<i>Dentatherina mercei</i>	Merce
(Shaw & Nodder 1794)	From Bali Strait to Timor Sea. In range Ref.: 4113.		Patten & Ivantsoff 1983	SL
<i>Antennarius randalli</i>	Randall's frogfish, (M), Ref. 1602. Max. 2.1 cm SL.		<i>Hypoatherina barnesi</i>	Barne
Allen 1970			Schultz 1953	
<i>Antennarius striatus</i>	Striated frogfish, (M, Br), Ref. 6773. Max. 22 cm.		<i>Hypoatherina ovalaua</i>	Fijian
(Shaw & Nodder 1794)			(Herre 1935)	
<i>Antennatus tuberosus</i>	Tuberculated frogfish, (M, Dan), Ref. 6773. Max. 7 cm SL.		<i>Hypoatherina valenciennesi</i>	Suma
(Cuvier 1817)			(Bleeker 1853)	
<i>Histrio histrio</i>	Sargassumfish, (M), Ref. 6773. Max. 20 cm TL.		<i>Kalyptatherina helodes</i>	Marin
(Linnaeus 1758)			(Ivantsoff & Allen 1984)	Ma
Chaunacidae	Sea toads		<i>Pseudomugil inconspicuus</i>	Incon
<i>Chaunax fimbriatus</i>	(M), Ref. 5978. Max. 14 cm SL. Museum:		Roberts 1978	Ma
Hilgendorf 1879	NTM S.10998-010 (TGT2531). From southwest Sumatra to Timor Sea.		<i>Stenatherina panatela</i>	Pana
			(Jordan & Richardson 1908)	TL
Gigantactinidae			Beloniformes (needle fishes)	
<i>Gigantactis perlatus</i>	(M), Ref. 559		Belonidae	Need
Beebe & Crane 1947			<i>Strongylura urvillii</i>	(M, B
Lophichthyidae	Lophichthyid frogfishes		(Valenciennes 1846)	
<i>Lophichthys boschmai</i>	(M, En), Ref. 245.		<i>Tylosurus crocodilus crocodilus</i>	Houn
Boeseman 1964			(Peron & Lesueur 1821)	Ma
Lophiidae	Goosefishes		<i>Tylosurus punctulatus</i>	(M), F
<i>Lophiodes gracilimanus</i>	(M), Ref. 5978. Museum: BMNH 1984.1.1.53.		(Günther 1872)	
(Alcock 1899)	From southwest Sumatra to Bali Strait.		Exocoetidae	Flyin
<i>Lophiodes mutilus</i>	Smooth angler, (M), Ref. 5978. Max. 45 cm. Museum:		<i>Cheilopogon arcticeps</i>	(M), F
(Alcock 1894)	BMNH 1984.1.1.54 (PJPW2035). From southwest Sumatra to Bali Strait. In range Ref.: 3461.		(Günther 1866)	
<i>Lophiomus setigerus</i>	Blackmouth angler, (M), Ref. 5978. Max. 40 cm.		<i>Cheilopogon katoptron</i>	(M), F
(Vahl 1797)	From southwest Sumatra to Timor Sea.		(Bleeker 1866)	
Ogcocephalidae	Batfishes		<i>Cypselurus hexazona</i>	(M), F
<i>Halieutaea coccinea</i>	(M), Ref. 5978. Museum: NTM S.11117-002. From southwest Sumatra to Bali Strait.		(Bleeker 1853)	
Alcock 1889			<i>Cypselurus oligolepis</i>	(M, F
<i>Halieutaea fumosa</i>	(M), Ref. 5978. Max. 14 cm SL. Museum: NTM S.		(Bleeker 1866)	
Alcock 1894	10761-010 (TGT3152). From Bali Strait to Timor Sea.		<i>Cypselurus opisthopus</i>	Black
<i>Halieutaea indica</i>	Indian handfish, (M), Ref. 5978.		(Bleeker 1866)	SL
Annandale & Jenkins 1910	Museum: NTM S.11001-005 (TGT2556). From southwest Sumatra to Bali Strait.		<i>Cypselurus poecilopterus</i>	Yellow
			(Valenciennes 1847)	
<i>Halieutaea stellata</i>	Starry handfish, (M), Ref. 5978. Max. 30 cm TL.		<i>Hirundichthys oxycephalus</i>	Bony
(Vahl 1797)	Museum: NTM S.10995-005 (TGT2556).		(Bleeker 1852)	
<i>Malthopsis luteus</i>	From southwest Sumatra to Timor Sea.		Hemiramphidae	Halfb
Alcock 1891	(M), Ref. 5978. Max. 6 cm SL. Museum: NTM S.10760-013 (TGT1711). From Bali Strait to Timor Sea.		<i>Hemiramphus archipelagicus</i>	Jump
Tetrabrachiidae			Collette & Parin 1978	
			<i>Hemiramphus lutkei</i>	Lutke
			(Valenciennes 1847)	
			<i>Hyporhamphus balinensis</i>	(M), F

Table 2. Continuation.
[Tabel 2. Sambungan.]

	(Bleeker 1859)			<i>Myripristis pralinia</i>	Scarle
	<i>Hyporhamphus dussumieri</i>	Dussumier's halfbeak, (M), Ref. 9843. Max. 29.8 cm SL.		Cuvier 1829	Alsi
	(Valenciennes 1846)			<i>Myripristis trachyacron</i>	(M), R
	<i>Hyporhamphus melanopterus</i>	(M), Ref. 9843. Max. 17 cm SL.		Bleeker 1863	
	Collette & Parin 1978			<i>Myripristis violacea</i>	Lattice
	<i>Hyporhamphus neglectissimus</i>	(M), Ref. 9843. Max. 14.4 cm SL. Irian Jaya.		Bleeker 1851	from
	Parin, Collette & Schcherbachev 1980			<i>Myripristis vittatus</i>	Whitel
	<i>Hyporhamphus neglectus</i>	(M), Ref. 9843. Max. 16.5 cm SL.		Valenciennes 1831	Kn
	(Bleeker 1866)			<i>Neoniphon argenteus</i>	Clearf
	<i>Hyporhamphus quoyi</i>	Quoy's garfish, (M, Br, Fr), Ref. 9843. Max. 31.2 cm SL.		(Valenciennes 1831)	Kn
	(Valenciennes 1847)			<i>Neoniphon opercularis</i>	Black
	<i>Oxyporhamphus convexus convexus</i>	(M), Ref. 9843.		(Valenciennes 1831)	Mus
	(Weber & de Beaufort 1922)				Tim
	<i>Oxyporhamphus micropterus micropter</i>	Bigwing halfbeak, (M), Ref. 9843. Max. 18.5 cm SL.		<i>Neoniphon sammara</i>	Samm
	(Valenciennes 1847)			(Forsskål 1775)	Ma
	<i>Rhynchorhamphus georgii</i>	Long billed half beak, (M), Ref. 9843.		<i>Ostichthys acanthorinus</i>	(M), F
	(Valenciennes 1847)			Ahl 1923	26
	<i>Zenarchopterus buffonis</i>	(M, Br), Ref. 7050. Max. 12.5 cm SL.			Sea
	(Valenciennes 1847)			<i>Ostichthys japonicus</i>	(M), F
	<i>Zenarchopterus rasori</i>	(M), Ref. 10988. Muna Island, Celebes.		(Cuvier 1829)	(TG
	(Popta 1912)			<i>Ostichthys kaianus</i>	Deepv
				(Günther 1880)	
Beryciformes (sawbellies)				<i>Sargocentron caudimaculatum</i>	Silver
Anomalopidae		Lanterneye fishes		(Rüppell 1838)	Ma
<i>Anomalops katoptron</i>		Splitfin flashlightfish, (M), Ref. 5004. Max. 35 cm.		<i>Sargocentron comutum</i>	Three
(Bleeker 1856)				(Bleeker 1853)	Ma
<i>Photoblepharon palpebratus palpebra</i>		Eyelight fish, (M), Ref. 5004. Max. 12 cm. Also Ref.: 1602, 4537, 8631.		<i>Sargocentron diadema</i>	Crown
(Boddaert 1781)				(Lacepède 1801)	Alsi
					(TG
Anoplogastridae		Fangtooth			Str
<i>Anoplogaster cornuta</i>		Common fangtooth, (M), Ref. 4737. Max. 15.2 cm SL.		<i>Sargocentron melanospilos</i>	Black
(Valenciennes 1833)				(Bleeker 1858)	TL
				<i>Sargocentron punctatissimum</i>	Speck
Diretmidae		Spinyfins		(Cuvier 1829)	Mu
<i>Diretmoides veriginiae</i>		(M), Ref. 9927. Max. 23.3 cm SL.			Tim
Kotlyar 1987				<i>Sargocentron rubrum</i>	Redcc
				(Forsskål 1775)	Ret
Holocentridae		Squirrelfishes, soldierfishes			Bal
<i>Myripristis adusta</i>		Shadowfin soldierfish, (M), Ref. 5300. Max. 33 cm.		<i>Sargocentron spiniferum</i>	Sabre
Bleeker 1853				(Forsskål 1775)	Mu
<i>Myripristis amaenus</i>		Brick soldierfish, (M), Ref. 1602. Max. 26 cm TL.			Sur
(Castelnau 1873)				<i>Sargocentron violaceum</i>	Violet
<i>Myripristis berndti</i>		Blotcheye soldierfish, (M), Ref. 5378. Max. 30 cm TL.		(Bleeker 1853)	Kn
Jordan & Evermann 1903		Museum: BPBM 29389 (TGT2190). Found from southwest Sumatra to Timor Sea (Ref. 5978).			Bigsc
		Blacktip soldierfish, (M), Ref. 5300. Max. 30 cm.		Melamphaidae	(M), F
<i>Myripristis hexagona</i>				<i>Melamphaes eulepis</i>	
(Lacepède 1802)				Ebeling 1962	
<i>Myripristis kuntee</i>		Pearly soldierfish, (M), Ref. 5300. Max. 17 cm SL.		<i>Poromitra oscitans</i>	(M), F
Cuvier 1831		Also Ref.: 8631. Museum: BPBM 29388 (TGT2180). Found from Bali Strait to Timor Sea (Ref. 5978).		Ebeling 1975	
				<i>Scopelogadus mizolepis mizolepis</i>	(M), F
<i>Myripristis melanosticta</i>		Splendid squirrelfish, (M), Ref. 5378. Max. 30 cm TL.		(Günther 1878)	BV
Bleeker 1863		Museum: CSIRO CA1545. From Bali Strait to Timor Sea (Ref. 5978).			Frc
<i>Myripristis murdjan</i>		Pinecone soldierfish, (M, Dan), Ref. 5300. Max. 30 cm.			42-
(Forsskål 1775)		Also Ref.: 8631. Museum: LPPL JIF12 (PAN9). Found from Bali Strait to Timor Sea (Ref. 5978).		Monocentridae	Pinec
				<i>Monocentris japonica</i>	Pinec
<i>Myripristis parvidens</i>		Small-eyed squirrelfish, (M), Ref. 5300. Max. 17.3 cm SL.			
Cuvier 1829					

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

(Houttuyn 1782)	Museum: CSIRO CA1807. From southwest Sumatra to Timor Sea.	<i>Fistularia petimba</i> Lacepède 1803	Red co Mus Sum
Trachichthyidae	Slimeheads	Solenostomidae	Ghost
<i>Aulotrachichthys latus</i> (Fowler 1938)	(M), Ref. 5978. Museum: CSIRO CA1545. From Bali Strait to Timor Sea.	<i>Solenostomus cyanopterus</i> Bleeker 1854	Ghost Kno
<i>Hoplostethus melanopus</i> (Weber 1913)	(M), Ref. 4181. Max. 25 cm TL.	<i>Solenostomus paradoxus</i> (Pallas 1770)	Harleq
<i>Hoplostethus rubellopterus</i> Kotlyar 1980	(M), Ref. 9872.	Syngnathidae	Pipefis
<i>Hoplostethus shubnikovii</i> Kotlyar 1980	(M), Ref. 9872.	<i>Apterygocampus epinnulatus</i> Weber 1913	(M), Re
Zeiformes (dories)		<i>Bhanotia fasciolata</i> (Duméril 1870)	(M), Re
Caproidae	Boarfishes	<i>Bulbonaricus brauni</i> (Dawson & Allen 1978)	Pughe
<i>Antigonia capros</i> Lowe 1843	Deepbody boarfish, (M), Ref. 5978. Max. 30 cm SL. Museum: NTM S.10760-015 (TGT1742). From southwest Sumatra to Timor Sea.	<i>Bulbonaricus davaoensis</i> (Herald 1953)	(M), Re
<i>Antigonia malayana</i> Weber 1913	(M), Ref. 5978. Museum: NTM S.11034-002 (TGT2330). From southwest Sumatra to Timor Sea.	<i>Choeroichthys brachysoma</i> (Bleeker 1855)	Short-b Fro
<i>Antigonia rubescens</i> (Günther 1860)	Indo-Pacific boarfish, (M), Ref. 5978. Max. 22 cm TL. Museum: NTM S.11116-004. From southwest Sumatra to Timor Sea.	<i>Choeroichthys cinctus</i> Dawson 1976	(M), Re 214
Macrurocyttidae		<i>Choeroichthys sculptus</i> (Günther 1870)	Sculpt
<i>Cyttula macropus</i> Weber 1913	(M), Ref. 6543. Max. 6.8 cm TL. Specimens collected from Flores Sea.	<i>Corythoichthys amplexus</i> Dawson & Randall 1975	Brown- Fro
Oreosomatidae	Oreos	<i>Corythoichthys flavofasciatus</i> (Rüppell 1838)	Netwo fro
<i>Alloctytus verrucosus</i> (Gilchrist 1906)	Warty oreo, (M), Ref. 6545. Max. 42 cm.	<i>Corythoichthys haematopterus</i> (Bleeker 1851)	(M), Re Flo
Zeidae Dories		<i>Corythoichthys intestinalis</i> (Ramsay 1881)	Bande
<i>Cyttopsis cypho</i> (Fowler 1934)	(M), Ref. 5978. Museum: NTM S.10998-011 (TGT2529). From southwest Sumatra to Timor Sea.	<i>Corythoichthys schultzi</i> Herald 1953	Schult.
<i>Zenopsis conchifer</i> (Lowe 1852)	Silvery John dory, (M), Ref. 5978. Max. 70 cm TL. Museum: LPPL JIF16 (TGT(PJPW)390). From southwest Sumatra to Bali Strait.	<i>Doryrhamphus dactylophorus</i> (Bleeker 1853)	Alsc Ringec
<i>Zenopsis nebulosus</i> (Temminck & Schlegel 1847)	Mirror dory, (M, Fi), Ref. 5978. Max. 70 cm. Museum: NTM S.10752-010 (TGT3229). From Bali Strait to Timor Sea.	<i>Doryrhamphus excisus excisus</i> Kaup 1856	Alsc Bluest Ref.
Gasterosteiformes (sticklebacks and seamoths)		<i>Doryrhamphus janssi</i> (Herald & Randall 1972)	Janss' Kno
Pegasidae	Seamoths	<i>Doryrhamphus multiannulatus</i> (Regan 1903)	Many-
<i>Eurypegasus draconis</i> (Linnaeus 1766)	Short dragonfish, (M, Br), Ref. 1418. Max. 10 cm TL. Known from Sumatra and Java.	<i>Doryrhamphus negrosensis negrosensis</i> (M), Herre 1934	(M),
<i>Pegasus volitans</i> Linnaeus 1758	Longtail seamouth, (M, Br), Ref. 1418. Max. 17.5 cm TL. Known from Sumatera and Java.	<i>Festucalex erythraeus</i> (Gilbert 1905)	Red pi
Syngnathiformes (pipefishes and seahorses)		<i>Festucalex prolixus</i> Dawson 1984	(M), R
Aulostomidae	Trumpetfishes	<i>Halicampus grayi</i> Kaup 1856	Gray's
<i>Aulostomus chinensis</i> (Linnaeus 1766)	Chinese trumpetfish, (M), Ref. 8631. Max. 80 cm TL. Known from Flores.	<i>Halicampus macrorhynchus</i> Bamber 1915	Ornate Fou rec
Centriscidae	Snipefishes and shrimpfishes	<i>Haliichthys taeniophorus</i> Gray 1859	Ribbo
<i>Aeoliscus strigatus</i> (Günther 1860)	Razorfish, (M), Ref. 8631. Max. 15 cm TL. Known from Flores.		
<i>Centriscus scutatus</i> Linnaeus 1758	Grooved razorfish, (M, Br), Ref. 561. Max. 15 cm TL. Museum: NTM S.10749-007 (TGT1812). Found from southwest Sumatra to Timor Sea (Ref. 5978).		
Fistulariidae	Cornetfishes		
<i>Fistularia commersonii</i> Rüppell 1838	Bluespotted cornetfish, (M), Ref. 1602. Max. 160 cm TL.		

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Hippichthys cyanospilos</i> (Bleeker 1854)	Blue-spotted pipefish, (M, Br, Fr), Ref. 5316. Max. 16 cm SL. Nias, Java, and Banda (Ref. 7050).	<i>Dactyloptena orientalis</i> (Cuvier 1829)	Orie Al Fr in
<i>Hippichthys penicillus</i> (Cantor 1849)	Beady pipefish, (M, Br, Fr), Ref. 5316. Max. 18 cm SL. Kalimantan and Java (Ref. 7050).	<i>Dactyloptena peterseni</i> (Nyström 1887)	Star M Se
<i>Hippichthys spicifer</i> (Rüppell 1838)	Bellybarred pipefish, (M, Br, Fr), Ref. 5316. Max. 17 cm SL. Sumatra, Java, and Sulawesi (Ref. 7050).	Hoplichthyidae	Gho
<i>Hippocampus kuda</i> Bleeker 1852	Spotted seahorse, (M, Br), Ref. 5978. Max. 30 cm TL. Also Ref.: 8631. Museum: NTM S.10749-006. Known from Bali Strait to Timor Sea, including Flores.	<i>Hoplichthys gilberti</i> Jordan & Richardson 1908	(M), 2E
<i>Ichthyocampus carce</i> (Hamilton 1822)	(M, Br, Fr), Ref. 5316. Max. 14 cm SL. Sumatra, Kalimantan, Sulawesi, and Java (Ref. 7050).	<i>Hoplichthys regani</i> Jordan & Richardson 1908	Ghos Mi
<i>Micrognathus brevirostris pygmaeus</i> Fritzsche 1981	(M), Ref. 1602. Known from Moluccas.	Platycephalidae	to
<i>Micrognathus micronotopterus</i> (Fowler 1938)	(M), Ref. 5316. Max. 5.7 cm SL.	<i>Cociella crocodila</i> (Tilesius 1812)	Flat Croc M
<i>Microphis argulus</i> (Peters 1855)	Flat-nosed pipefish, (M, Br, Fr), Ref. 5316. Max. 18 cm SL. Java and Flores (Ref. 7050).	<i>Cociella punctatus</i> (Cuvier 1829)	19 In Spot
<i>Minyichthys brachyrhinus</i> (Herald 1953)	(M), Ref. 5316. Max. 4.6 cm SL.	<i>Cymbacephalus beauforti</i> (Knapp 1973)	Croc Mc
<i>Minyichthys myersi</i> (Herald & Randall 1972)	Myers' pipefish, (M), Ref. 1602. Max. 5.8 cm SL.	<i>Cymbacephalus nematophthalmus</i> (Günther 1860)	Fring
<i>Phoxocampus belcheri</i> (Kaup 1856)	Rock pipefish, (M), Ref. 5316. Max. 7.2 cm SL.	<i>Elates ransonnetii</i> (Steindachner 1876)	Dwar US to
<i>Phoxocampus tetrophthalmus</i> (Bleeker 1858)	(M), Ref. 5316. Max. 7.5 cm SL.	<i>Grammolites scaber</i> (Linnaeus 1758)	Roug M Str Ka
<i>Siokunichthys breviceps</i> Smith 1963	(M), Ref. 5173. Max. 8 cm SL.	<i>Inegocia japonica</i> (Tilesius 1812)	Japar M Se
<i>Siokunichthys herrei</i> Herald 1953	(M), Ref. 5173. Max. 7.6 cm SL.	<i>Levanaora bosschei</i> (Bleeker 1860)	Small
<i>Siokunichthys nigrolineatus</i> Dawson 1983	(M), Ref. 5173. Max. 7.4 cm SL. Also Ref.: 8631. Known from Flores.	<i>Onigocia macrolepis</i> (Bleeker 1854)	Notch M Str
<i>Syngnathoides biaculeatus</i> (Bloch 1785)	Alligator pipefish, (M), Ref. 8631. Max. 28.3 cm SL. Known from Bali.	<i>Onigocia pedimacula</i> (Regan 1908)	(M), F
<i>Trachyrhamphus bicoarctatus</i> (Bleeker 1857)	Double-ended pipefish, (M), Ref. 1602. Max. 39 cm SL.	<i>Onigocia spinosa</i> (Temminck & Schlegel 1844)	(M), F
Scorpaeniformes (scorpionfishes and flatheads)		<i>Platycephalus arenarius</i> Ramsay & Ogilby 1886	North TL Bal diff eas
Aploactinidae	Velvetfishes	<i>Platycephalus indicus</i> (Linnaeus 1758)	Indiar In n
<i>Erisphex philippinus</i> (Fowler 1938)	(M), Ref. 5978. Museum: CAS 54604. From southwest Sumatra to Bali Strait.	<i>Rogadius asper</i> (Cuvier 1829)	Thorn Mu Stra
<i>Kanekonia aniara</i> Thompson 1968	Darkfined velvetfish, (M), Ref. 5978. Museum: CAS 54596. From Bali Strait to Timor Sea.	<i>Rogadius pristiger</i> (Cuvier 1829)	(M), F
<i>Kanekonia pelta</i> Poss 1982	(M, En), Ref. 8937. Max. 2.85 cm SL. Halmahera Is. off Teluk Kau.		
Bembridae	Deepwater flatheads		
<i>Bembradium roseum</i> Gilbert 1905	(M), Ref. 5978. Max. 11 cm SL. Museum: NTM S.10760- 004 (TGT1710). From Bali Strait to Timor Sea.		
<i>Bembras japonicus</i> Cuvier 1829	(M), Ref. 5978. Max. 30 cm SL. Museum: USNM 264796 (TGT1726). From Bali Strait to Timor Sea.		
Caracanthidae	Orbicular velvetfishes		
<i>Caracanthus maculatus</i> (Gray 1831)	Spotted coral croucher, (M), Ref. 1602. Max. 5 cm.		
Dactylopteridae	Flying gurnards		
<i>Dactyloptena macracanthus</i> (Bleeker 1854)	Spotwing flying gurnard, (M), Ref. 3392. Max. 16.5 cm TL. Museum: CAS 53136 (TGT2464). From southwest Sumatra to Timor Sea (Ref. 5978).		

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	<i>Rogadius serratus</i> (Cuvier 1829)	Serrated flathead, (M), Ref. 9790. Max. 24 cm TL.	<i>Lioscorpius longiceps</i> Günther 1880	(M, Dan)
	<i>Rogadius welanderi</i> (Schultz 1966)	Welander's flathead, (M), Ref. 9790. Max. 13 cm TL. Museum: USNM 264803 (TGT3030). From Bali Strait to Timor Sea (Ref. 5978).	<i>Minous monodactylus</i> (Bloch & Schneider 1801)	Grey st
	<i>Sorsogona tuberculata</i> (Cuvier 1829)	Tuberculated flathead, (M), Ref. 5978. Max. 14 cm TL. Museum: NTM S.11013-002 (TGT1799). From Bali Strait to Timor Sea. In range Ref.: 5999.	<i>Minous pictus</i> Günther 1880	Painted 002 (
	<i>Suggrundus macracanthus</i> (Bleeker 1869)	Large-spined flathead, (M), Ref. 5978. Max. 26 cm TL. Museum: CSIRO B.2126. From southwest Sumatra to Timor Sea.	<i>Neocentropogon aeglefinis</i> (Weber 1913)	(M), Re From
	<i>Suggrundus rodericiensis</i> (Cuvier 1829)	Spiny flathead, (M), Ref. 9790. Max. 25 cm.	<i>Neomerinthe amplisquamiceps</i> (Fowler 1938)	(M), Re
	<i>Thysanophrys arenicola</i> Schultz 1966	Broadhead flathead, (M), Ref. 9790. Max. 37 cm TL.	<i>Neomerinthe megalepis</i> (Fowler 1938)	(M), Re From
	<i>Thysanophrys carbunculus</i> (Valenciennes 1833)	Papillose flathead, (M), Ref. 9790. Max. 18 cm TL.	<i>Neomerinthe procurva</i> Chen 1981	(M), Re From
	<i>Thysanophrys celebica</i> (Bleeker 1854)	Celebes flathead, (M), Ref. 9790. Max. 15 cm TL. Celebes and Irian Barat.	<i>Neomerinthe rotunda</i> Chen 1981	(M), Re From
	<i>Thysanophrys chiltonae</i> Schultz 1966	Longsnout flathead, (M), Ref. 2334. Max. 23 cm TL.	<i>Paracentropogon longispinus</i> (Cuvier 1829)	Wispy 1531
	<i>Thysanophrys malayanus</i> (Bleeker 1853)	(M), Ref. 5978. Museum: NTM S.10734-015 (TGT1221). From southwest Sumatra to Timor Sea.	<i>Pontinus macrocephalus</i> (Sauvage 1882)	(M), Re From
	<i>Thysanophrys otaitensis</i> (Parkinson 1829)	Fringelip flathead, (M), Ref. 9790. Max. 25 cm TL.	<i>Pteroidichthys amboinensis</i> Bleeker 1856	(M), Re and
	Scorpaenidae	Scorpionfishes or rockfishes	<i>Pterois antennata</i> (Bloch 1787)	Broadb TL. A (TG
236	<i>Ablabys taenianotus</i> (Cuvier 1829)	Cockatoo waspfish, (M, Dan), Ref. 3132. Max. 10 cm TL.	<i>Pterois mombasae</i> (Smith 1957)	Frillfin Mus Sea
	<i>Apistus carinatus</i> (Bloch & Schneider 1801)	Ocellated waspfish, (M, Fi, Dan), Ref. 5978. Max. 18 cm. Museum: LPPL JIF19 (TGT1198). From southwest Sumatra to Darwin.	<i>Pterois radiata</i> Cuvier 1829	Radial Mus Time
	<i>Cottapistus cottoides</i> (Linnaeus 1764)	Marbled stingfish, (M), Ref. 5978. Museum: UMMZ 212292 (TGT3245). From Bali Strait to Timor Sea.	<i>Pterois russelli</i> Bennett 1831	Plainta
	<i>Dendrochirus biocellatus</i> (Fowler 1938)	Twospot turkeyfish, (M, Dan), Ref. 8631. Max. 10 cm. Known from Flores.	<i>Pterois volitans</i> (Linnaeus 1758)	Lionfis Mus Time
	<i>Dendrochirus brachypterus</i> (Cuvier 1829)	Shortfin turkeyfish, (M, Dan), Ref. 5978. Max. 17 cm. Also Ref.: 8631. Known from Bali Strait to Timor Sea, including Flores.	<i>Rhinopias frondosa</i> (Günther 1891)	Weedy Mus to TI
	<i>Dendrochirus zebra</i> (Cuvier 1829)	Zebra turkeyfish, (M, Dan), Ref. 5978. Max. 25 cm SL. Also Ref.: 8631. Museum: CSIRO CA1303. From Bali Strait to Timor Sea.	<i>Scorpaena picta</i> (Cuvier 1829)	Northe Mus
	<i>Ebosia bleekeri</i> (Döderlein 1884)	(M), Ref. 5978. Museum: ANSP 152033 (TGT3189). From Bali Strait to Timor Sea.	<i>Scorpaenodes albaiensis</i> (Evermann & Seale 1907)	Longfir Max
	<i>Ectreposebastes imus</i> Garman 1899	(M, Fi), Ref. 5978. Max. 18 cm SL. Museum: BMNH 1984. 1.1.63 (TGT (PJPW) 824). From southwest Sumatra to Bali Strait.	<i>Scorpaenodes guamensis</i> (Quoy & Gaimard 1824)	Guam Kno
	<i>Inimicus cuvieri</i> (Gray 1835)	(M), Ref. 5978. Museum: LPPL JIF20 (TGT1394). From Bali Strait to Timor Sea.	<i>Scorpaenopsis cirrhosa</i> (Thunberg 1793)	Weedy, Kno
	<i>Inimicus didactylus</i> (Pallas 1769)	Bearded ghou, (M, Br, Dan), Ref. 5978. Max. 21.5 cm TL. Museum: LPPL JIF (TGT2366). From southwest Sumatra to Bali Strait.	<i>Scorpaenopsis diabolus</i> Cuvier 1829	False :
	<i>Inimicus sinensis</i> (Valenciennes 1833)	Spotted ghou, (M, Dan), Ref. 5978. Max. 26 cm. Museum: BMNH 1984.1.1.64 (TGT (PJPW) 711). From southwest Sumatra to Bali Strait.	<i>Scorpaenopsis macrochir</i> Ogilby 1910	Flashe
			<i>Scorpaenopsis neglecta</i> Heckel 1837	(M, Da
			<i>Scorpaenopsis oxycephalus</i>	Tassle

Table 2. Continuation.
[Tabel 2. Sambungan.]

(Bleeker 1849)	Museum: NTM S.10734-014 (TGT1227). From southwest Sumatra to Timor Sea.	<i>Satyrichthys moluccense</i> (Bleeker 1850)	re Blac M sc
<i>Scorpaenopsis venosa</i> (Cuvier 1829)	Raggy scorpionfish, (M, Dan), Ref. 2334. Max. 18 cm TL.	<i>Satyrichthys rieffeli</i> (Kaup 1859)	Spot M sc
<i>Sebastapistes cyanostigma</i> (Bleeker 1856)	Yellowspotted scorpionfish, (M, Dan), Ref. 8631. Max. 8 cm. Known from Bali.		
<i>Sebastiscus tertius</i> (Basukov & Chen 1978)	(M), Ref. 5978. Max. 37 cm SL. Museum: SDSU 83-14 (TGT1751). From Bali Strait to Timor Sea.	Perciformes (perch-likes)	
<i>Setarches guentheri</i> Johnson 1862	Channeled rockfish, (M, Fi, Dan), Ref. 3503. Max. 24 cm.	Acanthoclinidae	Spin
<i>Setarches longimanus</i> (Alcock 1894)	(M, Dan), Ref. 5978. Max. 18 cm SL. Museum: CSIRO CA1584. From southwest Sumatra to Bali Strait. In range Ref.: 559.	<i>Acanthoplesiops hiatti</i> Schultz 1953	(M),
<i>Snyderina yamanokami</i> Jordan & Starks 1901	(M), Ref. 5978. Max. 24 cm TL. Museum: UMMZ 212291 (TGT3192). From Bali Strait to Timor Sea.	Acanthuridae	Surg
<i>Synanceia verrucosa</i> Bloch & Schneider 1801	Stonefish, (M, Dan), Ref. 5978. Max. 40 cm SL. Museum: LPPL JIF18 (TGT1020). From Bali Strait to Timor Sea.	<i>Acanthurus auranticavus</i> Randall 1956	Oran cr
<i>Tetraroge barbata</i> (Cuvier 1829)	Bearded roguefish, (M, Br, Fr, Dan), Ref. 7050. Max. 10 cm TL. Known in Sumatra, Java, Sulawesi, Seram, and Irian Jaya.	<i>Acanthurus bariene</i> (Lesson 1830)	Black Al Str
<i>Tetraroge niger</i> (Cuvier 1829)	(M, Br, Fr, Dan), Ref. 7050. Max. 13.5 cm TL. Known in Sumatra, Bali, Sulawesi and Moluccas.	<i>Acanthurus blochii</i> Valenciennes 1835	Ringt
<i>Vesplicula depressifrons</i> (Richardson 1848)	(M, Br, Fr, Dan), Ref. 7050. Max. 10 cm TL. Sumatra, Bali, Sulawesi, Moluccas, and Irian Jaya.	<i>Acanthurus dussumieri</i> Valenciennes 1835	Eyest Frc
Triglidae	Searobins	<i>Acanthurus fowleri</i> De Beaufort 1951	Fowl Flc
<i>Gargariscus prionocephalus</i> (Duméril 1869)	(M), Ref. 559. Max. 30 cm TL.	<i>Acanthurus grammoptilus</i> Richardson 1843	Fineli
<i>Lepidotrigla japonica</i> (Bleeker 1857)	(M), Ref. 5978. Max. 20 cm TL. Museum: NTM S.11013-004 (TGT1797). From Bali Strait to Timor Sea. Also Ref.: 559.	<i>Acanthurus japonicus</i> Schmidt 1930	(M), F
<i>Lepidotrigla punctipectoralis</i> Fowler 1938	(M), Ref. 5978. Max. 20 cm TL. Museum: NTM S.10752-015 (TGT3198). From southwest Sumatra to Timor Sea.	<i>Acanthurus leucocheilus</i> Herre 1927	Paleli Knc
<i>Lepidotrigla spiloptera</i> Günther 1880	Spotwing gurnard, (M), Ref. 5978. Max. 10 cm. Museum: NTM S.10999-004 (TGT2460). From southwest Sumatra to Timor Sea. In range Ref.: 3542.	<i>Acanthurus leucosternon</i> Bennett 1832	Powd cm
<i>Paraheminodus murrayi</i> (Günther 1880)	(M), Ref. 9771	<i>Acanthurus lineatus</i> (Linnaeus 1758)	Lined TL (PA)
<i>Peristedion liorhynchus</i> Günther 1872	Armoured gurnard, (M), Ref. 5978. Max. 40 cm. Museum: AMS I.22807-028. From Bali Strait to Timor Sea.	<i>Acanthurus maculiceps</i> (Ahl 1923)	White Ma
<i>Pterygotrigla hemisticta</i> (Temminck & Schlegel 1842)	Blackspotted gurnard, (M), Ref. 5978. Max. 25 cm. Museum: NTM S.10760-016 (TGT1737). From southwest Sumatra to Timor Sea.	<i>Acanthurus mata</i> Cuvier 1829	Blue-li cm Str
<i>Pterygotrigla leptacanthus</i> (Günther 1880)	Black-finned gurnard, (M), Ref. 3132. Max. 15 cm TL.	<i>Acanthurus nigricans</i> (Linnaeus 1758)	White TL
<i>Pterygotrigla multiocellata</i> (Matsubara 1937)	(M), Ref. 9771.	<i>Acanthurus nigricauda</i> Duncker & Mohr 1929	Epaul cm (Re)
<i>Pterygotrigla ryukyuensis</i> Matsubara & Hiyama 1932	(M), Ref. 5978. Max. 30 cm TL. Museum: NTM S.10998-007 (TGT2525). From southwest Sumatra to Timor Sea. Also Ref.: 559.	<i>Acanthurus nigrofuscus</i> (Forsskäl 1775)	Brown Knc 192
<i>Satyrichthys adeni</i> (Lloyd 1907)	(M), Ref. 5978. Max. 70 cm TL. Museum: NTM S.10760-017 (TGT1740). From Bali Strait to Timor Sea. A new	<i>Acanthurus nubilus</i> (Fowler & Bean 1929)	Bluelir Ref.
		<i>Acanthurus olivaceus</i> Bloch & Schneider 1801	Orang Max Fron
		<i>Acanthurus pyroferus</i>	Choco

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

Kittlitz 1834	TL. Known from Bali and Flores.	Acropomatidae	Lanter
<i>Acanthurus thompsoni</i> (Fowler 1923)	Thompson's surgeonfish, (M), Ref. 8631. Max. 27 cm TL. Known from Flores. Also from Java and Sumatra (Ref. 1920).	<i>Acropoma japonicum</i> Günther 1859	Glow-b JIF4 Sea.
<i>Acanthurus triostegus</i> (Linnaeus 1758)	Convict surgeonfish, (M, Fi, Dan), Ref. 8631. Max. 27 cm TL. Known from Bali. Also from Java and Sumatra (Ref. 1920).	<i>Doederleinia berycoides</i> (Hilgendorf 1879)	(M), Re (con Timor
<i>Acanthurus tristis</i> Randall 1993	Indian Ocean mimic surgeonfish, (M), Ref. 8940. Max. 16.5 cm SL. Also Ref.: 8631.	<i>Malakichthys elegans</i> Matsubara & Yamaguti 1943	(M), Re 019 Strai
<i>Acanthurus xanthopterus</i> Valenciennes 1835	Yellowfin surgeonfish, (M, Fi, Dan), Ref. 5978. Max. 70 cm. Also Ref.: 8631. Museum: LPPL JIF210 (TGT1034). From southwest Sumatra to Timor Sea.	<i>Synagrops japonicus</i> (Döderlein 1883)	Japane Mus Sum
<i>Ctenochaetus binotatus</i> Randall 1955	Twospot surgeonfish, (M, Dan), Ref. 1602. Max. 22 cm TL.	<i>Synagrops philippinensis</i> (Günther 1880)	(M), Re Front
<i>Ctenochaetus striatus</i> (Quoy & Gaimard 1825)	Striated surgeonfish, (M, Dan), Ref. 8631. Max. 26 cm TL. Known from Bali.	Ambassidae	Glass
<i>Ctenochaetus strigosus</i> (Bennett 1828)	Spotted surgeonfish, (M), Ref. 1602. Max. 13.9 cm SL.	<i>Ambassis gymnocephalus</i> (Lacepède 1802)	Bald gl TL.
<i>Ctenochaetus tominiensis</i> Randall 1955	Tomini surgeonfish, (M), Ref. 1602. Max. 10 cm SL. Also Ref.: 8631. Known from Bali and Sulawesi (Celebes).	<i>Ambassis urotaenia</i> Bleeker 1852	(M, Br,
<i>Naso annulatus</i> (Quoy & Gaimard 1825)	White margin unicornfish, (M), Ref. 1602. Max. 100 cm TL.	Ammodytidae	Sand I
<i>Naso brachycentron</i> (Valenciennes 1835)	Humpback unicornfish, (M), Ref. 5978. Max. 90 cm FL. Museum: LPPL JIF130 (TGT3175). From southwest Sumatra to Timor Sea.	<i>Bleekeria mitsukurii</i> Jordan & Evermann 1902	(M), Re 007
<i>Naso brevirostris</i> (Valenciennes 1835)	Spotted unicornfish, (M), Ref. 1602. Max. 60 cm FL.	<i>Bleekeria viridianguilla</i> (Fowler 1931)	(M), Re Front
<i>Naso fageni</i> Morrow 1954	Horseface unicornfish, (M), Ref. 8631. Max. 80 cm. Known from Flores.	Apogonidae	Cardin
<i>Naso hexacanthus</i> (Bleeker 1855)	Sleek unicornfish, (M, Fi, Sport), Ref. 8631. Max. 75 cm FL. Known from Flores.	<i>Apogon angustatus</i> (Smith & Radcliffe 1911)	Striped Kno
<i>Naso lituratus</i> (Bloch & Schneider 1801)	Orangespine unicornfish, (M, Dan), Ref. 5978. Max. 45 cm TL. From southwest Sumatra to Timor Sea. Also Ref.: 8631.	<i>Apogon apogonides</i> (Bleeker 1856)	Shortc Ref. P.28 Sur
<i>Nasolopezi</i> Herre 1927	Elongate unicornfish, (M), Ref. 1602. Max. 54 cm FL. Also Ref.: 8631. From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Apogon aureus</i> (Lacepède 1802)	Ringtai Also (TG 5978
<i>Naso minor</i> (Smith 1966)	Slender unicorn, (M), Ref. 4974. Max. 20 cm TL.	<i>Apogon bandanensis</i> Bleeker 1854	Bigeye
<i>Naso thynnoides</i> (Valenciennes 1835)	Oneknife unicornfish, (M), Ref. 5978. Max. 40 cm. From southwest Sumatra to Timor Sea. Also Ref.: 8631.	<i>Apogon brevicaudata</i> Weber 1909	Manyb WAI
<i>Naso tuberosus</i> Lacepède 1801	Humpnose unicornfish, (M), Ref. 5978. Max. 60 cm FL. Museum: LPPI JIF133 (TGT2343). From southwest Sumatra to Timor Sea.	<i>Apogon ceramensis</i> Bleeker 1852	Ceram Mus Stra
<i>Naso unicornis</i> (Forsskål 1775)	Bluespine unicornfish, (M, Sport, Dan), Ref. 5978. Max. 70 cm TL. Also Ref.: 8631. From southwest Sumatra to Timor Sea.	<i>Apogon chrysoaenia</i> Bleeker 1851	Many-I Mar Flor
<i>Naso vlamingii</i> (Valenciennes 1835)	Bignose unicornfish, (M), Ref. 8631. Max. 55 cm TL. Known from Bali.	<i>Apogon coccineus</i> Rüppell 1838	Ruby c Kno Ma corr
<i>Zebrasoma scopas</i> (Cuvier 1829)	Twotone tang, (M), Ref. 1602. Max. 20 cm SL.	<i>Apogon compressus</i> (Smith & Radcliffe 1911)	Ochre- 863
<i>Zebrasoma veliferum</i> (Bloch 1795)	Sailfin tang, (M), Ref. 1602. Max. 40 cm TL.	<i>Apogon cookii</i> Macleay 1881	Cook's Kno
		<i>Apogon cyanosoma</i>	Yellow

Table 2. Continuation.
[Tabel 2. Sambungan.]

	Bleeker 1853	the Manado area, Sulawesi (Celebes).	<i>Apogon nigrocincta</i>	(M),
	<i>Apogon dispar</i>	Redspot cardinalfish, (M), Ref. 8631. Max. 4.8 cm SL.	(Smith & Radcliffe 1912)	Fr
	Fraser & Randall 1976	Known from Flores.	<i>Apogon nigrofasciatus</i>	Black
	<i>Apogon ellioti</i>	Flag-in cardinalfish, (M), Ref. 5978. Max. 16 cm TL.	Lachner 1953	Al
	Day 1875	Museum: NTM S.10996-003 (TGT2308).	<i>Apogon notatus</i>	Spot
		From southwest Sumatra to Timor Sea. In range Ref.: 1602.	(Houttuyn 1782)	Kr
	<i>Apogon evermanni</i>	Evermann's cardinalfish, (M), Ref. 8631. Max. 11 cm SL.	<i>Apogon novemfasciatus</i>	Seve
	Jordan & Snyder 1904	Known from Flores.	Cuvier 1828	SL
	<i>Apogon exostigma</i>	Narrowstripe cardinalfish, (M), Ref. 8631. Max. 9.4 cm		(T
	(Jordan & Starks 1906)	SL. Known from Flores.	<i>Apogon perlitus</i>	Pearl
	<i>Apogon fasciatus</i>	Broadbanded cardinalfish, Glaga, (M), Ref. 5978. Max.	Fraser & Lachner 1985	Kr
		13 cm TL. White 1790 Museum: LPPL JIF41		Ko
		(TGT2160). From southwest Sumatra to Timor Sea.	<i>Apogon poecilopterus</i>	Pearl
		In range Ref.: 4329.	Cuvier 1828	Ma
	<i>Apogon fleuriu</i>	Cardinalfish, (M), Ref. 2142. Max. 11 cm SL.		Tir
	(Lacepède 1802)	Museum: BMNH 1855.3.24.41, Ambon.	<i>Apogon sangiensis</i>	Sang
	<i>Apogon fraenatus</i>	Bridled cardinalfish, (M), Ref. 8631. Max. 8.5 cm SL.	Bleeker 1857	
	Valenciennes 1832	Known from Bali.	<i>Apogon sealei</i>	Seale
	<i>Apogon fragilis</i>	Fragile cardinalfish, (M), Ref. 1602. Max. 4.3 cm SL.	Fowler 1918	Kr
	Smith 1961	Known from Maumere Bay, Flores (Allen, pers. comm.).	<i>Apogon semilineatus</i>	Half-
	<i>Apogon fuscus</i>	Samoan cardinalfish, (M), Ref. 8631. Max. 7.7 cm SL.	Temminck & Schlegel 1842	TL
	Quoy & Gaimard 1825	Known from Flores.		Fr
	<i>Apogon gilberti</i>	Gilbert's cardinalfish, (M), Ref. 1602. Max. 4.2 cm SL.	<i>Apogon semiornatus</i>	Obliq
	(Jordan & Seale 1905)		Peters 1876	TL
	<i>Apogon guamensis</i>	Guam cardinalfish, (M), Ref. 8631. Max. 7.9 cm SL.		(A
	Valenciennes 1832	Known from Bali.	<i>Apogon septemstriatus</i>	(M),
	<i>Apogon hartzfeldii</i>	Hartzfeld's cardinalfish, (M), Ref. 8631. Max. 8.2 cm SL.	Günther 1880	Fr
	Bleeker 1852	Known from Bali, Flores, and the Moluccas. Also Ref.: 1602.	<i>Apogon taeniophorus</i>	Reef
	<i>Apogon hoevenii</i>	Frostfin cardinalfish, (M), Ref. 5978. Max. 5 cm TL.	Regan 1908	16
	Bleeker 1854	Also Ref.: 8631. Museum: WAM P.28046-001		Su
	<i>Apogon kallopterus</i>	(TGT1051). From Bali Strait to Timor Sea.	<i>Apogon thermalis</i>	Ko
	Bleeker 1856	Iridescent cardinalfish, (M), Ref. 8631. Max. 12.2 cm SL.	Cuvier 1829	Half-
	<i>Apogon kiensis</i>	Known from Bali.	<i>Apogon timorensis</i>	Ba
	Jordan & Snyder 1901	Rifle cardinal, (M, Br), Ref. 8631. Max. 8 cm TL. Known from Flores.	Bleeker 1854	Timor
	<i>Apogon lateralis</i>	Humpback cardinal, (M, Br, Fr), Ref. 7050. Max. 8 cm		cm
	Valenciennes 1832	SL. Also Ref.: 8631. Known from Flores.	<i>Apogon trimaculatus</i>	Ba
	<i>Apogon leptacanthus</i>	Threadfin cardinalfish, (M), Ref. 1602. Max. 4.5 cm SL.	Cuvier 1828	Thre
	Bleeker 1856	Known from Maumere Bay, Flores and Komodo I., Allen, pers. comm.	<i>Archamia biguttata</i>	SL
	<i>Apogon melas</i>	Black cardinalfish, (M), Ref. 1602. Max. 9.2 cm SL.	Lachner 1951	Twins
	Bleeker 1848		<i>Archamia dispilus</i>	Al
	<i>Apogon moluccensis</i>	Moluccan cardinalfish, (M), Ref. 5978. Max. 9 cm TL.	Lachner 1951	(M),
	Valenciennes 1832	Also Ref.: 8631. Museum: WAM P.28044-004	<i>Archamia fucata</i>	Oran
	<i>Apogon multilineatus</i>	(TGT1049). From Bali Strait to Timor Sea.	(Cantor 1849)	SL
	(Bleeker 1865)	Many-lined cardinalfish, Multi-striped cardinal, (M), Ref.		(T
	<i>Apogon nigripinnis</i>	8631. Max. 10 cm TL. From Flores. In range Ref.: 6192.	<i>Archamia lineolata</i>	inc
	Cuvier 1828	Bullseye, (M), Ref. 5978. Max. 10 cm TL. Museum: CSIRO CA1617. From Bali Strait to Timor Sea. In range Ref.: 4329.	(Ehrenberg 1828)	Shim
			<i>Archamia zosterophora</i>	Black
			(Bleeker 1858)	Al
			<i>Cheilodipterus alleni</i>	(M),
			Gon 1993	Fl
			<i>Cheilodipterus artus</i>	Wolf
			Smith 1961	Al
				Ce

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	Ambon, BPBM 31471; ZMH 14372. Bone Rate Is., BPBM 31512. Also known from Flores.	Lachner 1953	SL
<i>Cheilodipterus isostigmus</i> (Schultz 1940)	Dog-toothed cardinalfish, (M), Ref. 1602. Max. 9.3 cm SL. Museum: NTM S.10733-019 (TGT1064). From Bali Strait to Timor Sea (Ref. 5978).	<i>Siphamia majimae</i> Matsubara & Iwai 1958	(M), I co
<i>Cheilodipterus macrodon</i> (Lacepède 1802)	Large-toothed cardinalfish, (M), Ref. 5978. Max. 20 cm SL. Also Ref.: 8525. Museum: LPPL JIF42 (TGT1589). From Bali Strait to Timor Sea. Bay of Jakarta, ZMA 101.377. Lombok, BPBM 30048. Sulawesi, Ujung Pandang (Makassar), RMNH 74.	<i>Siphamia versicolor</i> (Smith & Radcliffe 1911)	(M), I M Str
<i>Cheilodipterus nigrotaeniatus</i> Smith & Radcliffe 1912	(M), Ref. 8525. Max. 6.4 cm SL. Museum: Molucca Is., Halmahera I., USNM 112305.	<i>Sphaeramia nematoptera</i> (Bleeker 1856)	Pajar Als
<i>Cheilodipterus quinquelineatus</i> Cuvier 1828	Five-lined cardinalfish, (M), Ref. 8631. Max. 13 cm TL. Known from Flores.	<i>Sphaeramia orbicularis</i> (Cuvier 1828)	Orbic Als Kn
<i>Cheilodipterus singapurensis</i> Bleeker 1859	Truncate cardinalfish, (M), Ref. 8525. Max. 17.5 cm SL. Museum: Java, USNM 261570. Krimundjawa I., USNM 261573. Celebes, (Makassar), USNM 149337; ZMA 101.379 (Holotype of <i>C. subulatus</i> Weber). Kabaena I., USNM 261567. Buru I., USNM 149335. Doworra I., USNM 149336. Misol I., BMNH 1870.8.31.15.	Ariommatidae <i>Ariomma brevimanum</i> (Klunzinger 1884)	Arior (M), F 10' Tin
<i>Foa brachygramma</i> (Jenkins 1903)	Weed cardinalfish, (M), Ref. 1602. Max. 4 cm SL. Known from Maumere Bay, Flores, Allen pers. comm.	<i>Ariomma indica</i> (Day 1870)	Indiar M sol
<i>Fowleria abocellata</i> Goren & Karplus 1980	(M), Ref. 2334. Max. 5 cm TL.	Banjosiidae <i>Banjos banjos</i> (Richardson 1846)	(M), F P.1
<i>Fowleria aurita</i> (Valenciennes 1831)	Crosseyed cardinalfish, (M), Ref. 2334. Max. 9 cm TL.	Blenniidae <i>Andamia tetradactylus</i> (Bleeker 1858)	Coml (M), F Kn
<i>Fowleria marmorata</i> (Alleyne & MacLeay 1877)	Marbled cardinalfish, (M), Ref. 1602. Max. 7.5 cm TL. Known from the Manado area, Sulawesi (Celebes) and Maumere Bay, Flores, Allen pers. comm.	<i>Aspidontus dussumieri</i> (Valenciennes 1836)	Lance Kn
<i>Fowleria variegata</i> (Valenciennes 1832)	Variiegated cardinalfish, (M), Ref. 1602. Max. 6.5 cm SL. Known from Komodo I. (Allen, pers. comm.).	<i>Aspidontus taeniatus taeniatus</i> Quoy & Gaimard 1834	False
<i>Gymnapogon urospilotus</i> Lachner 1953	(M), Ref. 1602. Max. 2.7 cm SL. Known from Maumere Bay, Flores (Allen pers. comm.).	<i>Atrosalarias fuscus holomelas</i> (Günther 1872)	Brow Kn
<i>Pseudamia amblyuroptera</i> (Bleeker 1856)	(M, Br), Ref. 526. Max. 8 cm SL. Museum: NTM S.10824-001 (TGT2370). Found from Bali Strait to Timor Sea (Ref. 5978).	<i>Blenniella bilitonensis</i> (Bleeker 1858)	(M), F
<i>Pseudamia gelatinosa</i> Smith 1955	Gelatinous cardinalfish, (M), Ref. 526. Max. 7.9 cm SL.	<i>Blenniella periophthalmus</i> (Valenciennes 1836)	Blue-c SL
<i>Pseudamia hayashii</i> Randall, Lachner & Fraser 1985	Hayashi's cardinalfish, (M), Ref. 526. Max. 6.2 cm SL.	<i>Cirripectes auritus</i> Carlson 1981	Black Knc
<i>Pterapogon kauderni</i> Koumans 1933	(M), Ref. 9936. Apparently restricted to the Benggai Is. off the east coast of central Sulawesi (Celebes), Indonesia. Museum: RMNH 17003.	<i>Cirripectes castaneus</i> (Valenciennes 1836)	Chest Als
<i>Rhabdamia cypselurus</i> Weber 1909	(M), Ref. 1602. Max. 5.1 cm SL. Known from the Manado area, Sulawesi (Celebes), Maumere Bay, Flores, and Komodo I. (Allen, pers. comm.).	<i>Cirripectes filamentosus</i> (Alleyne & Macleay 1877)	Filame
<i>Rhabdamia gracilis</i> (Bleeker 1856)	Luminous cardinalfish, (M), Ref. 8631. Max. 5.1 cm SL. In range Ref.: 1602. Museum: WAM P.28137-00 (TGT2283). Found from southwest Sumatra to Timor Sea (Ref. 5978); including Flores.	<i>Cirripectes gilberti</i> Williams 1988	(M), F off,
<i>Siphamia fistulosa</i> (Weber 1909)	(M), Ref. 1602. Max. 1.6 cm SL.	<i>Cirripectes polyzona</i> (Bleeker 1868)	Barrec
<i>Siphamia fuscolineata</i>	Crown-of-thorns cardinalfish, (M), Ref. 5978. Max. 4 cm	<i>Cirripectes quagga</i> (Fowler & Ball 1924)	Squig
		<i>Cirripectes springeri</i> Williams 1988	Spring Ref
		<i>Cirripectes stigmaticus</i> Strasburg & Schultz 1953	Red-s
		<i>Crossosalarias macrospilus</i> Smith-Vaniz & Springer 1971	Tripple
		<i>Ecsenius bandanus</i> Springer 1971	Banda Mus

Table 2. Continuation.
 [Tabel 2. Sambungan.]

	Banda Sea. Seribu Is., USNM 211988. Karimunjawa, USNM 211979. Bone Betang, BPBM 26720. Kabaena, AMS I.18490-001. Ambon, USNM 209767. Ceram, USNM 209660. Saparua, USNM 209991. Biak, USNM 221236.	(Schneider & Forster 1801) <i>Istiblennius lineatus</i> (Valenciennes 1836) <i>Laiphognathus multimaculatus</i> Smith 1955	Linea Spot Fowl
<i>Ecsenius bathi</i> Springer 1988	Bath's comb-tooth, (M, En), Ref. 5296. Max. 3.6 cm SL. Known from Bali, Toko Toko Rock. Also from Flores (Ref. 8631).	<i>Litobranchius fowleri</i> (Herre 1926) <i>Meiacanthus anema</i> (Bleeker 1852) <i>Meiacanthus atrodorsalis</i> (Günther 1877) <i>Meiacanthus ditrema</i> Smith-Vaniz 1976	(M, E) St Fork Al Ones Mi ar
<i>Ecsenius bicolor</i> (Day 1888)	Bicolor blenny, (M), Ref. 5296. Max. 11 cm TL. Also Ref.: 8631. Known from Bali.	<i>Meiacanthus grammistes</i> (Valenciennes 1836)	Strip- Mi Fl
<i>Ecsenius lividanalis</i> Chapman & Schultz 1952	(M), Ref. 5296. Max. 4 cm TL.	<i>Meiacanthus smithi</i> Klausewitz 1961 <i>Meiacanthus vittatus</i> Smith-Vaniz 1976	Disc W (M),
<i>Ecsenius melarchus</i> McKinney & Springer 1976	Yellow-eyed comb-tooth, (M), Ref. 5296. Max. 5 cm TL. Also Ref.: 8631. Known from Flores.	<i>Nannosalarias nativitatus</i> (Regan 1909)	Pygn
<i>Ecsenius midas</i> Starck 1969	Persian blenny, (M), Ref. 5296. Max. 13 cm TL. Also Ref.: 8631. Known from Flores.	<i>Omobranchus elongatus</i> (Peters 1855) <i>Omobranchus punctatus</i> (Valenciennes 1836)	Clois Muzz
<i>Ecsenius monoculus</i> Springer 1988	(M), Ref. 5296. Max. 5 cm TL.	<i>Omox biporos</i> Springer 1972	Omo:
<i>Ecsenius namiyei</i> (Jordan & Evermann 1903)	Black comb-tooth, (M), Ref. 5296. Max. 9 cm TL. Also Ref.: 8631. Known from Bali.	<i>Petrosirtes breviceps</i> (Valenciennes 1836) <i>Petrosirtes mitratus</i> Rüppell 1830	Strip Me Flora Kn
<i>Ecsenius paroculus</i> Springer 1988	(M), Ref. 5296 Museum: Pulau Tikus, Pulau Pari Group, Pulau Seribu, USNM 260389 (Holotype). Also known from Bawean I. off N central Java.	<i>Petrosirtes thepassii</i> (Bleeker 1853) <i>Petrosirtes variabilis</i> Cantor 1850	Thep Ma Varia Ma
<i>Ecsenius pictus</i> McKinney & Springer 1976	White-lined comb-tooth, (M), Ref. 5296. Max. 5 cm TL. Also Ref.: 8631. Known from Moluccas, Bone Rate islands, and Flores.	<i>Petrosirtes xestus</i> Jordan & Seale 1906 <i>Plagiotremus laudandus</i> (Whitley 1961) <i>Plagiotremus rhinorhynchus</i> (Bleeker 1852)	Xestu SL Bicol Kn Blues
<i>Ecsenius schroederi</i> McKinney & Springer 1976	(M), Ref. 9137. Max. 7 cm TL.	<i>Plagiotremus tapeinosoma</i> (Bleeker 1857) <i>Praealticus amboinensis</i> (Bleeker 1857)	Pian SL Ambo Kn
<i>Ecsenius stigmatura</i> Fowler 1952	(M), Ref. 5296. Max. 6 cm TL.	<i>Salarias ceramensis</i> Bleeker 1852 <i>Salarias fasciatus</i> (Bloch 1786) <i>Salarias guttatus</i> Valenciennes 1836 <i>Salarias sinuosus</i> Snyder 1908	(M), F Jewe Kn Breas Flo Fring
<i>Ecsenius trilineatus</i> Springer 1972	Three-lined blenny, White-spotted comb-tooth, (M), Ref. 5296. Max. 2.7 cm SL. Museum: Moluccas, Saparua, USNM 211926; Banda Islands, USNM 211930, 211941, 211945; presumably Banda Sea, USNM 202477. Kai Islands, USNM 221239. Also from Flores (Ref. 8631).		
<i>Ecsenius yaeyamaensis</i> (Aoyagi 1954)	Yaeyama blenny, (M), Ref. 5296. Max. 5.2 cm SL. Also Ref.: 8631. Known from Flores.		
<i>Enchelyurus kraussi</i> (Klunzinger 1871)	Krauss' blenny, (M), Ref. 1602. Max. 4.5 cm SL.		
<i>Entomacrodus caudofasciatus</i> (Regan 1909)	Tail-barred rockskipper, (M), Ref. 1602. Max. 6.2 cm SL.		
<i>Entomacrodus decussatus</i> (Bleeker 1857)	Wavy-lined blenny, (M), Ref. 1602. Max. 6.7 cm SL.		
<i>Entomacrodus thalassinus</i> (Jordan & Seale 1906)	(M), Ref. 1602. Max. 4 cm SL.		
<i>Exallias brevis</i> (Kner 1868)	Leopard blenny, (M), Ref. 8631. Max. 11 cm SL. Known from Flores.		
<i>Istiblennius chrysospilos</i> (Bleeker 1857)	Redspotted blenny, (M), Ref. 1602. Max. 11 cm SL.		
<i>Istiblennius cyanostigma</i> (Bleeker 1849)	Bluespotted blenny, (M), Ref. 1602. Max. 7.4 cm SL. Known from Moluccas.		
<i>Istiblennius dussumieri</i> (Valenciennes 1836)	Streaky rockskipper, (M, Br), Ref. 2334. Max. 12 cm.		
<i>Istiblennius edentulus</i>	Rippled rockskipper, (M, Fr), Ref. 1602. Max. 14 cm SL.		

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Stanulus seychellensis</i> Smith 1959	Seychelle's blenny, (M), Ref. 1602. Max. 2.7 cm SL.	<i>Pterocaesio trilineata</i> Carpenter 1987	Three TL.
<i>Xiphasia matsubarai</i> Okada & Suzuki 1952	Japanese snake blenny, (M), Ref. 1602. Max. 30 cm SL.	Callionymidae	Drage
<i>Xiphasia setifer</i> Swainson 1839	Hairtail blenny, (M), Ref. 2334. Max. 53 cm. Also Ref.: 8631. Museum: CSIRO CA2282, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Anaora tentaculata</i> Gray 1835	(M), F
Bramidae	Pomfrets	<i>Callionymus filamentosus</i> Valenciennes 1837	Blotch Mu sou
<i>Brama dussumieri</i> Cuvier 1831	Lowfin pomfret, (M), Ref. 5978. Max. 19 cm SL. Museum: NTM S.11119-001. From southwest Sumatra to Bali Strait. In range Ref.: 3326.	<i>Callionymus japonicus</i> Houttuyn 1782	(M), F
<i>Taractes rubescens</i> (Jordan & Evermann 1887)	Black pomfret, (M), Ref. 5978. Max. 70 cm SL. Museum: LPPL JIF50 (TGT2125). From Bali Strait to Timor Sea.	<i>Callionymus meridionalis</i> Suwardji 1965	(M), F (TG)
Caesionidae	Fusiliers	<i>Callionymus semeiophor</i> Fricke 1983	(M), F
<i>Caesio caeruleaurea</i> Lacepède 1801	Blue and gold fusilier, Pisang-pisang, (M, Fi, Bait), Ref. 402. Max. 35 cm TL.	<i>Callionymus superbus</i> Fricke 1983	(M), F
<i>Caesio cuning</i> (Bloch 1791)	Redbelly yellowtail fusilier, Ekor kuning, (M, Fi), Ref. 402. Max. 60 cm TL. Museum: BPBM 29378 (TGT2172). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Callionymus whiteheadi</i> Fricke 1981	(M), F (Su)
<i>Caesio lunaris</i> Cuvier 1830	Lunar fusilier, Pisang-pisang, (M, Fi), Ref. 402. Max. 40 cm TL. Museum: BPBM 29334, from Bali Strait to Timor Sea (Ref. 5978).	<i>Diplogrammus goramensis</i> (Bleeker 1858)	(M), F
<i>Caesio teres</i> Seale 1906	Yellow and blueback fusilier, Ekor kuning pisang, (M, Fi), Ref. 402. Max. 40 cm TL. Museum: BPBM 29367, from Bali Strait to Timor Sea (Ref. 5978).	<i>Synchiropus altivelis</i> (Temminck & Schlegel 1850)	(M), F 011
<i>Caesio varilineata</i> Carpenter 1987	Variable-lined fusilier, (M, Fi, Bait), Ref. 402. Max. 40 cm TL.	<i>Synchiropus morrisoni</i> Schultz 1960	Morris In
<i>Caesio xanthonota</i> Bleeker 1853	Yellowback fusilier, Ekor kuning pisang, (M, Fi), Ref. 402. Max. 40 cm TL. Museum: BPBM 29375, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Synchiropus ocellatus</i> (Pallas 1770)	Ocella
<i>Dipterygonotus balteatus</i> (Valenciennes 1830)	Mottled fusilier, (M, Fi, Bait), Ref. 402. Max. 14 cm TL. Museum: BPBM 29376, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Synchiropus splendidus</i> (Herre 1927)	Mand In r
<i>Gymnocaesio gymnoptera</i> (Bleeker 1856)	Slender fusilier, (M, Fi, Bait), Ref. 402. Max. 18 cm TL. Museum: BPBM 29377, from southwest Sumatra to Timor Sea (Ref. 5978).	Carangidae	Jacks
<i>Pterocaesio chrysozona</i> (Cuvier 1830)	Goldband fusilier, Pisang-pisang, (M, Fi, Bait), Ref. 402. Max. 21 cm TL. Museum: WAM P.26191-003. From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Alectis ciliaris</i> (Bloch 1787)	Africa cm sou
<i>Pterocaesio digramma</i> (Bleeker 1865)	Double-lined fusilier, (M, Fi), Ref. 402. Max. 30 cm TL. Museum: BPBM 29341, from Bali Strait to Timor Sea (Ref. 5978).	<i>Alectis indicus</i> (Rüppell 1830)	Indian 150 Sea
<i>Pterocaesio marri</i> Schultz 1953	Marr's fusilier, (M, Fi, Bait), Ref. 402. Max. 35 cm TL.	<i>Alepes djedaba</i> (Forsskål 1775)	Shrim Ref sou
<i>Pterocaesio pisang</i> (Bleeker 1853)	Banana fusilier, Pisang-pisang, (M, Fi, Bait), Ref. 402. Max. 21 cm TL. Museum: BPBM 29342, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Alepes melanoptera</i> Swainson 1839	Black Mu sou
<i>Pterocaesio randalli</i> Carpenter 1987	Randall's fusilier, (M, Fi), Ref. 402. Max. 25 cm TL.	<i>Alepes vari</i> (Cuvier 1833)	Herrin Mu Stra Clefth
<i>Pterocaesio tessellata</i> Carpenter 1987	One-stripe fusilier, (M, Fi), Ref. 402. Max. 25 cm TL. Museum: BPBM 29368 (TGT1582). From southwest Sumatra to Timor (Ref. 5978).	<i>Atropus atropus</i> (Bloch & Schneider 1801)	Yellow 152 sou
<i>Pterocaesio tile</i> (Cuvier 1830)	Dark-banded fusilier, (M, Fi, Bait), Ref. 402. Max. 30 cm TL.	<i>Atule mate</i> (Cuvier 1833)	Longf Ma
		<i>Carangoides armatus</i> (Rüppell 1830)	Orang Ma
		<i>Carangoides bajad</i> (Forsskål 1775)	

Table 2. Continuation.
[Tabel 2. Sambungan.]

	152114 (TGT3002). Found from Bali Strait to Timor Sea (Ref. 5978).	<i>Caranx melampyus</i> Cuvier 1833	Bluefi cm
<i>Carangoides caeruleopinnatus</i> (Ruppell 1830)	Coastal trevally, (M, Fi, Sport), Ref. 3287. Max. 36 cm FL. Museum: LPPL JIF173 (TGT2396); LPPL JIF220 (TGT 3296) (as <i>C. uii</i>) From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Caranx papuensis</i> Alleyne & MacLeay 1877	Su Brass
<i>Carangoides chrysophrys</i> (Cuvier 1833)	Longnose trevally, (M, Br, Fi), Ref. 3280. Max. 60 cm TL. Museum: LPPL JIF174 (TGT2245). Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Caranx sexfasciatus</i> Quoy & Gaimard 1825	Ma Bigey cm 59
<i>Carangoides dinema</i> Bleeker 1851	Shadow trevally, (M, Br, Fi, Sport), Ref. 3287. Max. 59 cm TL. Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Caranx tille</i> Cuvier 1833	Tille t Mu Tir
<i>Carangoides ferdau</i> (Forsskål 1775)	Blue trevally, (M, Br, Fi, Sport), Ref. 3287. Max. 70 cm TL. Also Ref.: 8631. Museum: LPPL JIF175 (TGT3041). From Bali Strait to Timor Sea (Ref. 5978).	<i>Decapterus kurroides</i> Bleeker 1855	Redt Mu Ba
<i>Carangoides fulvoguttatus</i> (Forsskål 1775)	Yellowspotted trevally, (M, Fi, Sport), Ref. 3287. Max. 100 cm FL. Found from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Decapterus macarellus</i> (Cuvier 1833)	Mack cm Su
<i>Carangoides gymnostethus</i> (Cuvier 1833)	Bludger, (M, Fi, Sport), Ref. 3287. Max. 90 cm TL.	<i>Decapterus macrosoma</i> Bleeker 1851	Short Mu Su
<i>Carangoides hedlandensis</i> (Whitley 1934)	Bumpnose trevally, (M, Fi), Ref. 3287. Max. 28 cm FL. Museum: LPPL JIF176, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Decapterus russelli</i> (Rüppell 1830)	Indiar Mu Ba
<i>Carangoides humerosus</i> (McCulloch 1915)	Duskyshoulder trevally, (M, Fi), Ref. 3132. Max. 25 cm TL.	<i>Elagatis bipinnulata</i> (Quoy & Gaimard 1825)	Raint Al so
<i>Carangoides malabaricus</i> (Bloch & Schneider 1801)	Malabar trevally, Kwee, (M, Fi, Sport), Ref. 3280. Max. 60 cm. Also Ref.: 6567. Museum: LPPL JIF177, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Gnathanodon speciosus</i> (Forsskål 1775)	Gold FL so
<i>Carangoides oblongus</i> (Cuvier 1833)	Coachwhip trevally, (M, Fi, Sport), Ref. 3287. Max. 46 cm TL.	<i>Megalaspis cordyla</i> (Linnaeus 1758)	Torpe Mu Su
<i>Carangoides orthogrammus</i> (Jordan & Gilbert 1882)	Island trevally, (M, Fi, Sport), Ref. 3287. Max. 63 cm FL. Museum: LPPL JIF204, from Bali Strait to Timor Sea (Ref. 5978).	<i>Naucrates ductor</i> (Linnaeus 1758)	Pilotf
<i>Carangoides plagiotaenia</i> Bleeker 1857	Barcheek trevally, (M, Fi, Sport), Ref. 3287. Max. 50 cm. Museum: LPPL JIF178, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Pantolabus radiatus</i> (Macleay 1881)	Fring Ma
<i>Carangoides praeustus</i> (Bennett 1830)	Brownback trevally, (M, Fi, Sport), Ref. 3287. Max. 19.5 cm FL.	<i>Parastromateus niger</i> (Bloch 1795)	Black Re sc
<i>Carangoides talamparoides</i> Bleeker 1852	Imposter trevally, (M, Fi), Ref. 3287. Max. 28 cm FL. Museum: LPPL JIF179, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Scomberoides commersonianus</i> Lacepède 1801	Talar cm 59
<i>Caranx bucculentus</i> Alleyne & Macleay 1877	Bluespotted trevally, (M, Br, Fi, Sport), Ref. 2334. Max. 66 cm.	<i>Scomberoides lysan</i> (Forsskål 1775)	Dout M
<i>Caranx heberi</i> (Bennett 1830)	Blacktip trevally, (M, Br, Fi, Sport), Ref. 4537. Max. 85 cm. Also Ref.: 8631(Bali), 3287 and 3197 (as <i>C. sem</i>).	<i>Scomberoides tala</i> (Cuvier 1832)	Barre M Ti
<i>Caranx ignobilis</i> (Forsskål 1775)	Giant trevally, (M, Fi, Sport), Ref. 3280. Max. 165 cm TL. Museum: LPPL JIF180 (TGT3136). From southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 7050.	<i>Scomberoides tol</i> (Cuvier 1832)	Neec M Fr
<i>Caranx kleinii</i> (Bloch 1793)	Banded scad, (M, Fi), Ref. 4537. Max. 16 cm FL. Museum: LPPL JIF216 (TGT2120). From southwest Sumatra to Timor Sea, as <i>C. para</i> (Ref. 5978). Also Ref.: 3287, as <i>C. para</i> .	<i>Selar boops</i> (Cuvier 1833)	Oxe M m 59
		<i>Selar crumenophthalmus</i>	Bige

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

(Bloch 1793)	SL. Also Ref.: 6567. Museum: LPPL JIF187 (TGT2612). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Owstonia pectinifer</i> (Myers 1939)	(M, S Ba)
<i>Selaroides leptolepis</i> (Cuvier 1833)	Yellowstripe scad, (M, Fi), Ref. 3287. Max. 18.5 cm FL. Museum: ANSP 152026, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Owstonia totoniensis</i> Tanaka 1908	(M, F ML Ba)
<i>Seriola dumerili</i> (Risso 1810)	Greater amberjack, (M, Fi, Aq, Sport, Dan), Ref. 3397. Max. 190 cm TL. Museum: LPPL JIF188, from southwest Sumatra to Bali Strait (Ref. 5978).	Chaetodontidae	Butte
<i>Seriola rivoliana</i> Valenciennes 1833	Almaco jack, (M, Fi, Sport, Dan), Ref. 3287. Max. 110 cm FL. Also Ref.: 8631. Museum: CSIRO CA2467, from Bali Strait to Timor Sea (Ref. 5978).	<i>Chaetodon adiergastos</i> Seale 1910	Philip Als
<i>Seriolina nigrofasciata</i> (Rüppell 1829)	Blackbanded trevally, (M, Fi, Sport), Ref. 3287. Max. 70 cm TL. Museum: LPPL JIF189 (TGT2443): From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Chaetodon assarius</i> Waite 1905	West cm Au up
<i>Trachinotus africanus</i> Smith 1967	Southern pompano, (M, Br, Fi, Sport), Ref. 3287. Max. 92 cm TL.	<i>Chaetodon aureofasciatus</i> Macleay 1878	Goldé Als
<i>Trachinotus baillonii</i> (Lacepède 1801)	Smallspotted dart, (M, Br, Fi), Ref. 3287. Max. 60 cm TL.	<i>Chaetodon auriga</i> Forsskål 1775	Threa Kn
<i>Trachinotus blochii</i> (Lacepède 1801)	Snubnose pompano, (M, Fi, Sport), Ref. 3280. Max. 110 cm FL.	<i>Chaetodon baronessa</i> Cuvier 1831	Easte cm 00f
<i>Trachinotus botla</i> (Shaw 1803)	Largespotted dart, (M, Br, Fi), Ref. 3197. Max. 75 cm TL.	<i>Chaetodon bennetti</i> Cuvier 1831	Bluele Ma Flo
<i>Trachinotus mookalee</i> Cuvier 1832	Indian pompano, (M, Fi, Sport), Ref. 3287. Max. 77 cm FL.	<i>Chaetodon citrinellus</i> Cuvier 1831	Spect ML Tir Redta
<i>Ulua aurochs</i> (Ogilby 1915)	Silvermouth trevally, (M, Fi), Ref. 4537. Max. 50 cm TL. Also Ref.: 3132.	<i>Chaetodon collare</i> Bloch 1787	
<i>Ulua mentalis</i> (Cuvier 1833)	Longrakered trevally, (M, Fi, Sport), Ref. 3287. Max. 100 cm TL. Museum: LPPL JIF190 (TGT1372). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Chaetodon decussatus</i> Cuvier 1831	Indian cm. LPF Sea
<i>Uraspis helvola</i> (Forster 1801)	Whitemouth jack, (M, Fi, Sport), Ref. 3287. Max. 46 cm FL.	<i>Chaetodon ephippium</i> Cuvier 1831	Saddl. Mu: Tim
<i>Uraspis uraspis</i> (Günther 1860)	Whitetongue jack, (M, Fi), Ref. 3287. Max. 28 cm FL. Museum: LPPL JIF191, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Chaetodon guentheri</i> Ahl 1923	Croch Als (TG Sea
Centrogeniidae		<i>Chaetodon guttatissimus</i> Bennett 1823	Peppe Knc
<i>Centrogenys vaigiensis</i> (Quoy & Gaimard 1824)	False scorpionfish, (M, Br), Ref. 8631. Known from Bali.	<i>Chaetodon kleinii</i> Bloch 1790	Sunbu Mus Tim
Centrolophidae		<i>Chaetodon lunulatus</i> Quoy & Gaimard 1825	(M), R
<i>Psenopsis obscura</i> Haedrich 1967	Medusafishes	<i>Chaetodon melannotus</i> Bloch & Schneider 1801	Blackt. Alsc
	Obscure ruff, (M), Ref. 4410. Max. 20 cm. Museum: BMNH 1984.1.1.96, BMNH 1984.1.1.97, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Chaetodon meyeri</i> Bloch & Schneider 1801	Frow Kno
Centropomidae		<i>Chaetodon ocellicaudus</i> Cuvier 1831	Spotta
<i>Lates calcarifer</i> (Bloch 1790)	Snooks	<i>Chaetodon ornatissimus</i> Solander 1831	Ornate Kno
<i>Psammoperca waigiensis</i> (Cuvier 1828)	Barramundi, Kakap, (M, Br, Fr, Fi, Aq, Sport), Ref. 3281. Max. 200 cm TL.		
Cepolidae	Waigieu seaperch, (M, Br), Ref. 9799. Max. 47 cm TL.		
<i>Acanthocephala abbreviata</i> (Valenciennes 1853)	Bandfishes		
<i>Acanthocephala krusensterni</i> (Temminck & Schlegel 1845)	Bandfish, (M), Ref. 5978. Museum: ANSP 152037 (TGT917). From Bali Strait to Timor Sea.		
<i>Cepola schlegelii</i> Bleeker 1854	(M), Ref. 5978. Max. 40 cm TL. Museum: ANSP 152036 (TGT2462). From southwest Sumatra to Bali Strait.		
	(M), Ref. 5978. Max. 50 cm TL. From southwest Sumatra to Bali Strait.		

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Chaetodon oxycephalus</i> Bleeker 1853	Spot-nape butterflyfish, (M), Ref. 8631. Max. 25 cm TL. Known from Bali.	(Cuvier 1829)	
<i>Chaetodon punctatofasciatus</i> Cuvier 1831	Spotband butterflyfish, (M), Ref. 8631. Max. 9.6 cm SL. Also Ref.: 4537. In range Ref.: 1602. Known from Flores.	<i>Parachaetodon ocellatus</i> (Cuvier 1831)	Sixsp Mu Ba
<i>Chaetodon rafflesii</i> Bennett 1830	Latticed butterflyfish, (M), Ref. 8631. Max. 15 cm TL. Known from Bali.	Champsodontidae	
<i>Chaetodon selene</i> Bleeker 1853	Yellowdotted butterflyfish, (M), Ref. 5978. Max. 18 cm TL. Also Ref.: 8631. Museum: LPPL JIF105 (TGT3227). From Bali Strait to Timor Sea.	<i>Champsodon arafurensis</i> Regan 1908	(M), Fr
<i>Chaetodon semeion</i> Bleeker 1855	Pec, (M), Ref. 8631. Max. 25.5 cm TL. Known from Flores.	<i>Champsodon capensis</i> Regan 1908	Gape Mu Fr 66
<i>Chaetodon speculum</i> Cuvier 1831	Mirror butterflyfish, (M), Ref. 4859. Max. 15 cm TL. Also Ref.: 8631.	<i>Champsodon guentheri</i> Regan 1908	(M), Su
<i>Chaetodon triangulum</i> Cuvier 1831	Triangle butterflyfish, (M), Ref. 4858. Max. 15 cm.	<i>Champsodon longipinnis</i> Matsubara & Amaoka 1964	(M), (T)
<i>Chaetodon trifascialis</i> Quoy & Gaimard 1824	Chevron butterflyfish, (M), Ref. 5978. Max. 18 cm TL. Also Ref.: 8631. Museum: LPPL JIF107 (TGT1096). From Bali Strait to Timor Sea.	Chiasmodontidae	
<i>Chaetodon ulietensis</i> Cuvier 1831	Pacific doublesaddle butterflyfish, (M), Ref. 8631. Max. 15 cm TL. In range Ref.: 1602.	<i>Pseudoscopelus altipinnis</i> Parr 1933	(M, S Fr
<i>Chaetodon vagabundus</i> Linnaeus 1758	Vagabond butterflyfish, (M), Ref. 8631. Max. 18 cm TL. Known from Bali.	Cichlidae	Cich
<i>Chaetodon xanthurus</i> Bleeker 1857	Pearlscale butterflyfish, (M), Ref. 8631. Max. 14 cm SL. Known from Flores.	<i>Oreochromis mossambicus</i> (Peters 1852)	Moza 17
<i>Coradion altivelis</i> McCulloch 1916	Highfin coralfish, (M), Ref. 4855. Max. 9.1 cm SL.	Cirrihitidae	Haw
<i>Coradion chrysozonus</i> (Cuvier 1831)	Goldengirdled coralfish, (M), Ref. 5978. Max. 11.5 cm SL. Museum: CSIRO CA998. From southwest Sumatra to Timor Sea.	<i>Amblycirrhitus bimacula</i> (Jenkins 1903)	Twos
<i>Coradion melanopus</i> (Cuvier 1831)	Twospot coralfish, (M), Ref. 4537. Max. 15 cm TL. Flores (Ref. 8631).	<i>Cirrhitichthys aprinus</i> (Cuvier 1829)	Spot Mu (R
<i>Forcipiger flavissimus</i> Jordan & McGregor 1898	Longnose butterfly fish, (M), Ref. 8631. Max. 17.5 cm SL. Known from Flores.	<i>Cirrhitichthys aureus</i> (Temminck & Schlegel 1843)	Yellow Mu Ba
<i>Forcipiger longirostris</i> (Broussonet 1782)	Longnose butterflyfish, (M), Ref. 8631. Max. 17.9 cm SL. Known from Flores.	<i>Cirrhitichthys falco</i> Randall 1963	Dwar Re
<i>Hemitaurichthys polylepis</i> (Bleeker 1857)	Pyramid butterflyfish, (M), Ref. 1602. Max. 12.7 cm SL. Also Ref.: 8631.	<i>Cirrhitichthys oxycephalus</i> (Bleeker 1855)	Cor Kr
<i>Heniochus acuminatus</i> (Linnaeus 1758)	Pennant coralfish, (M), Ref. 4859. Max. 20.5 cm SL. Museum: LPPL JIF108 (TGT3121). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Cirrhitus pinnulatus</i> (Forster 1801)	Stoc
<i>Heniochus chrysostomus</i> Cuvier 1831	Threeband pennantfish, (M), Ref. 5978. Max. 12.5 cm SL. Museum: LPPL JIF109 (TGT1547). From Bali Strait To Timor Sea.	<i>Cyprinocirrhites polyactis</i> (Bleeker 1875)	Swal Re S. Se
<i>Heniochus diphreutes</i> Jordan 1903	False moorish idol, (M), Ref. 5978. Max. 20 cm SL. Museum: CSIRO CA1504. From southwest Sumatra to Timor Sea.	<i>Oxycirrhites typus</i> Bleeker 1857	Long Kr
<i>Heniochus monoceros</i> Cuvier 1831	Masked bannerfish, (M), Ref. 4859. Max. 30 cm TL.	<i>Paracirrhites arcatus</i> (Cuvier 1829)	Arc-e
<i>Heniochus pleurotaenia</i> Ahl 1923.	Phantom bannerfish, (M), Ref. 4855. Max. 16 cm TL. West coast of Sumatra.	<i>Paracirrhites forsteri</i> (Schneider 1801)	Black Re St
<i>Heniochus singularis</i> Smith & Radcliffe 1911	Singular bannerfish, (M), Ref. 5978. Max. 23.7 cm SL. Also Ref.: 8631. Museum: LPPL JIF96 (TGT1025). From Bali Strait to Timor Sea.	Coryphaenidae	Dolp
<i>Heniochus varius</i>	Horned bannerfish, (M), Ref. 1602. Max. 15.2 cm SL.	<i>Coryphaena equiselis</i> Linnaeus 1758	Pomp
		<i>Coryphaena hippurus</i> Linnaeus 1758	Com Ma
		Creediidae	Sanc
		<i>Limnichthys fasciatus</i>	(M)

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

Waite 1904			<i>Lepidocybium flavobrunneum</i>	Escol.
Draconettidae			(Smith 1843)	Mu
<i>Centrodraco insolitus</i>	(M), Ref. 5978. Museum: SF P514-1984-003 (TGT1709).		<i>Nealotus tripes</i>	Bal
(McKay 1971)	From Bali Strait to Timor Sea.		Johnson 1865	Black
Drepanidae	Sicklefishes		<i>Neoepinnula orientalis</i>	Sackf
<i>Drepane longimana</i>	Concertina fish, (M, Br), Ref. 9800. Max. 50 cm TL.		(Gilchrist & von Bonde 1924)	
(Bloch & Schneider 1801)	Museum: NTM S.11016-005 (TGT2461).		<i>Nesiarchus nasutus</i>	Black
	From southwest Sumatra to Timor Sea (Ref. 5978).		Johnson 1862	
<i>Drepane punctata</i>	Spotted sicklefish, (M, Br, Fi), Ref. 9800. Max. 45 cm TL.		<i>Promethichthys prometheus</i>	Roudi
(Linnaeus 1758)	Museum: LPPL JIF93 (TGT2247). From southwest		(Cuvier 1832)	Mu
	Sumatra to Timor Sea (Ref. 5978).			(Re
Echeneidae	Remoras		<i>Rexea bengalensis</i>	Benga
<i>Echeneis naucrates</i>	Live sharksucker, (M, Sport), Ref. 5978. Max. 110 cm.		(Alcock 1894)	
Linnaeus 1758	Also Ref.: 8631. Museum: WAM P.26220-006.		<i>Rexea nakamurai</i>	Nakar
	From southwest Sumatra to Timor Sea.		Parin 1989	
<i>Remora remora</i>	Common remora, (M), Ref. 5978. Max. 86 cm TL.		<i>Rexea prometheoides</i>	Royal
(Linnaeus 1758)	Museum: NTM S.10730-003 (TGT1501). From Bali		(Bleeker 1856)	Mu
	Strait to Timor Sea.			NTI
Eleotridae	Sleepers		<i>Ruvettus pretiosus</i>	Sea
<i>Belobranchus belobranchus</i>	(M, Br, Fr), Ref. 7050. Max. 19.5 cm TL. Known from		Cocco 1834	Oilfish
Valenciennes 1837	Nias, Java, Sulawesi, Lesser Sundas, and		<i>Thyrsitoides marleyi</i>	Black
	Moluccas.		Fowler 1929	Mu:
<i>Butis amboinensis</i>	Olive flathead-gudgeon, (M, Br, Fr), Ref. 7050.			Tim
(Bleeker 1853)	Max. 13.6 cm TL.		Gerreidae	Mojar
<i>Eleotris acanthopoma</i>	(M, Br, Fr), Ref. 7050. Max. 11.8 cm SL.		<i>Gerres abbreviatus</i>	Deep-
Bleeker 1853			Bleeker 1850	SL.
Emmelichthyidae	Rovers			Tim
<i>Erythrocles schlegelii</i>	Japanese rubyfish, (M), Ref. 5978. Max. 50 cm.		<i>Gerres acinaces</i>	Longit
(Richardson 1846)	Museum: BMNH 1984.1.1.68 (TGT (PJPW) 846).		Bleeker 1854	Fro
	From southwest Sumatra to Bali Strait.			Alsi
Ephippidae	Spadefishes, batfishes and scats		<i>Gerres filamentosus</i>	Whippi
<i>Ephippus orbis</i>	Orbfish, (M, Fi), Ref. 5978. Max. 25 cm TL.		Cuvier 1829	cm
(Bloch 1787)	Museum: NTM S.11001-004. From southwest			sou
Sumatra	to Timor Sea.		<i>Gerres kapas</i>	Singaj
<i>Platax batavianus</i>	Humpback batfish, (M), Ref. 5978. Max. 50 cm TL.		Bleeker 1851	Mus
Cuvier 1831	Museum: LPPL JIF94 (TGT2450). From southwest			Stre
	Sumatra to Timor Sea.		<i>Gerres oyena</i>	Comrr
<i>Platax boersii</i>	(M), Ref. 9407. Max. 40 cm TL.		(Forsskål 1775)	Mus
Bleeker 1852				Stre
<i>Platax orbicularis</i>	Batfish, (M, Br), Ref. 5978. Max. 50 cm. Also Ref.:		<i>Gerres poietii</i>	Strong
(Forsskål 1775)	8631. From southwest Sumatra to Bali Strait.		Cuvier 1829	cm.
<i>Platax pinnatus</i>	Dusky batfish, (M), Ref. 8631. Max. 37 cm SL.			Longfi
(Linnaeus 1758)	Also Ref.: 4537. Known from Flores and Bali.		<i>Pentaprion longimanus</i>	656
<i>Platax teira</i>	Longfin batfish, (M, Fi), Ref. 5978. Max. 41 cm SL.		(Cantor 1850)	(PJ
(Bloch & Schneider 1801)	Also Ref.: 8631. Museum: LPPL JIF95 (TGT2440).			(Re
	From southwest Sumatra to Bali Strait; including		Gobiidae	Gobie
	Flores.		<i>Amblyeleotris diagonalis</i>	(M), R
<i>Zabidius novemaculeatus</i>	Ninespine batfish, (M), Ref. 4537. Max. 45 cm TL.		Polunin & Lubbock 1979	
(McCulloch 1916)			<i>Amblyeleotris fasciata</i>	Red-b.
Gempylidae	Snake mackerels		(Herre 1953)	
<i>Diplospinus multistriatus</i>	Striped escolar, (M), Ref. 6181. Max. 33 cm SL.		<i>Amblyeleotris fontanesii</i>	Giant
Maul 1948			(Bleeker 1852)	Alsc
<i>Gempylus serpens</i>	Snake mackerel, (M, Fi), Ref. 6181. Max. 100 cm SL.		<i>Amblyeleotris guttata</i>	Spotte
Cuvier 1829	Museum: NTM S.11118-001, from Bali Strait to Timor			
	Sea (Ref. 5978).			

Table 2. Continuation.
[Tabel 2. Sambungan.]

(Fowler 1938)	Known from Flores.	<i>Cryptocentrus strigiliceps</i> , (Jordan & Seale 1906)	Targe
<i>Amblyeleotris gymnocephala</i> (Bleeker 1853)	Masked shrimp-goby (M), Ref. 6771. Max. 14 cm TL. From Java. Also from Flores (Ref. 8631).	<i>Ctenogobiops aurocingulus</i> (Herre 1935)	Kr Gold
<i>Amblyeleotris periophthalmus</i> (Bleeker 1853)	(M), Ref. 2334. Max. 7.5 cm SL. Also Ref.: 8631. Known from Bali.	<i>Ctenogobiops feroculus</i> Lubbock & Polunin 1977	SL Sanc
<i>Amblyeleotris randalli</i> Hoese & Steene 1978	Randall's prawn-goby, (M), Ref. 1602. Max. 7.3 cm SL. Also Ref.: 8631. Known from Flores and the Moluccas.	<i>Ctenogobiops pomastictus</i> Lubbock & Polunin 1977	Kr Gold
<i>Amblyeleotris steinitzi</i> (Klausewitz 1974)	Steinitz' prawn-goby, (M), Ref. 8631. Max. 8 cm. Known from Flores.	<i>Ctenogobiops tangaroae</i> Lubbock & Polunin 1977	SL Tang;
<i>Amblyeleotris wheeleri</i> (Polunin & Lubbock 1977)	Gorgeous prawn-goby, (M), Ref. 8631. Max. 6.5 cm SL. Known from Bali.	<i>Ctenotrypauchen microcephalus</i> (Bleeker 1860)	Flc (M, B
<i>Amblygobius albimaculatus</i> (Rüppell 1830)	Butterfly goby, (M, Br), Ref. 4343. Max. 18 cm SL.	<i>Eviota afelei</i> Jordan & Seale 1906	Afele
<i>Amblygobius decussatus</i> (Bleeker 1855)	Orange-striped goby, (M), Ref. 1602. Max. 5.3 cm SL. Also Ref.: 8631. Known from Flores.	<i>Eviota bifasciata</i> Lachner & Karanella 1980	Twos Als
<i>Amblygobius nocturnus</i> (Herre 1945)	Nocturn goby, (M), Ref. 8631. Max. 4.1 cm SL. Known from Flores.	<i>Eviota herrei</i> Jordan & Seale 1906	Herre
<i>Amblygobius phalaena</i> (Valenciennes 1837)	Banded goby, (M), Ref. 8631. Max. 12 cm SL. Known from Bali.	<i>Eviota lachdeberei</i> Giltay 1933	Lachc SL
<i>Amblygobius rainfordi</i> (Whitley 1940)	Old glory, (M), Ref. 8631. Max. 6.5 cm TL. Known from Flores.	<i>Eviota melasma</i> Lachner & Karnella 1980	Melas
<i>Amblygobius sphynx</i> (Valenciennes 1837)	Sphinx goby, (M, Br), Ref. 2334. Max. 18 cm.	<i>Eviota nebulosa</i> Smith 1958	Nebul
<i>Apocryptodon madurensis</i> (Bleeker 1849)	(M), Ref. 5218. Max. 7.1 cm SL.	<i>Eviota nigriventris</i> Giltay 1933	(M), F Knc
<i>Bathygobius fuscus</i> (Rüppell 1830)	Dusky frill-goby, (M, Br, Fr), Ref. 1602. Max. 12 cm TL.	<i>Eviota pellucida</i> Larson 1976	Pelluc Knc
<i>Bathygobius padangensis</i> (Bleeker 1851)	(M), Ref. 559.	<i>Eviota prasina</i> (Klunzinger 1871)	Greer
<i>Bathygobius petrophilus</i> (Bleeker 1853)	(M), Ref. 559.	<i>Eviota prasites</i> Jordan & Seale 1906	Prasit Als
<i>Boleophthalmus boddarti</i> (Pallas 1770)	Boddart's goggle-eyed goby, (M, Br, Fr), Ref. 7050. Max. 22 cm TL. Kalimantan, Sumatra, and Java (Ref. 7050).	<i>Eviota punctulata</i> Jewett & Lachner 1983	Peppé
<i>Bryaninops tigris</i> Larson 1985	Black coral goby, (M), Ref. 8631. Max. 5.5 cm SL. Known from Bali.	<i>Eviota queenslandica</i> Whitley 1932	Queer SL
<i>Bryaninops yongei</i> (Davis & Cohen 1969)	Whip coral goby, (M), Ref. 8631. Max. 2.8 cm SL. Known from Flores.	<i>Eviota sebreei</i> Jordan & Seale 1906	Sebre Knc
<i>Callogobius centrolepis</i> Weber 1909	(M), Ref. 1602. Max. 4 cm SL.	<i>Eviota sparsa</i> Jewett & Lachner 1983	(M), R
<i>Callogobius snelli</i> Koumans 1953	(M), Ref. 559.	<i>Eviota spilota</i> Lachner & Karnella 1980	(M), R
<i>Cryptocentroides insignis</i> (Seale 1910)	Insignia prawn-goby, (M, Br), Ref. 1602. Max. 6.9 cm SL.	<i>Eviota storthynx</i> (Rofen 1959)	(M), R
<i>Cryptocentrus cinctus</i> (Herre 1936)	Yellow-prawn goby, (M), Ref. 1602. Max. 7.5 cm. Also Ref. 8631. Known from Flores.	<i>Eviota zonura</i> Jordan & Seale 1906	(M), R
<i>Cryptocentrus fasciatus</i> (Playfair & Günther 1867)	Y-bar shrimp goby, Black shrimp-goby, (M), Ref. 2334. Max. 8 cm TL. From Flores (Ref. 8631).	<i>Exyrias belissimus</i> (Smith 1959)	Mud re
<i>Cryptocentrus leucostictus</i> (Günther 1871)	Saddled prawn-goby, (M), Ref. 2334. Max. 7 cm TL.	<i>Exyrias puntang</i> (Bleeker 1851)	Puntar
<i>Cryptocentrus octofasciatus</i> Regan 1908	Blue-speckled prawn goby, (M), Ref. 8631. Max. 4.5 cm SL. Known from Flores.	<i>Fusigobius longispinus</i> Goren 1978	Orang TL
		<i>Fusigobius neophytus</i> (Günther 1877)	Comm from

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Fusigobius signipinnis</i> Hoese & Obika 1988	(M), Ref. 5261. Max. 4.9 cm SL.	<i>Pleurosicya bilobatus</i> (Koumans 1941)	(M), Ref.
<i>Gnatholepis cauerensis</i> (Bleeker 1853)	Eyebar goby, (M, Br), Ref. 4343. Max. 8 cm TL.	<i>Pleurosicya elongata</i> Larson 1990	Cling gob
<i>Gnatholepis scapulostigma</i> Herre 1953	Shoulderspot goby, Eye-bar sand-goby, (M), Ref. 2334. Max. 5.5 cm TL. From Flores (Ref. 8631).	<i>Priolepis fallacincta</i> Winterbottom & Burrigge 1992	(M), Ref.
<i>Gobiodon histrio</i> (Valenciennes 1837)	Broad-barred goby, (M), Ref. 2334. Max. 3.5 cm.	<i>Priolepis semidoliatus</i> (Valenciennes 1837)	(M), Ref.
<i>Gobiodon okinawae</i> Sawada, Arai & Abe 1972	Okinawa goby, Yellow-speckled cave-goby, (M), Ref. Known from Waigiu, Irian Jaya. Known from Waigiu, Irian Jaya. 1602. Max. 3.5 cm TL.	<i>Scartelaos histophorus</i> (Valenciennes 1837)	Walking
<i>Gobiopsis bravoii</i> (Herre 1940)	(M), Ref. 11081. Known from Waigiu, Irian Jaya. Also Ref.: 1602.	<i>Signigobius biocellatus</i> Hoese & Allen 1977	Twinspot Ref.: 8
<i>Istigobius decoratus</i> (Herre 1927)	Decorated goby, (M), Ref. 420. Max. 13 cm TL. Museum: Moluccas, USNM 254306, 254307.	<i>Stenogobius genivittatus</i> (Valenciennes 1837)	(M, Br, F
<i>Istigobius goldmanni</i> (Bleeker 1852)	(M), Ref. 420. Max. 5 cm TL. Also Ref.: 8631. Museum: Moluccas, USNM 211078.	<i>Stonogobius nematodes</i> Hoese & Randall 1982	(M), Ref
<i>Istigobius nigroocellatus</i> (Günther 1873)	Black-spotted goby, (M), Ref. 8631. Max. 4.9 cm TL. Known from Flores.	<i>Stonogobioops xanthorhinica</i> Hoese & Randall 1982	Yellownc 8631.
<i>Istigobius ornatus</i> (Rüppell 1830)	Ornate goby, (M, Br), Ref. 420. Max. 8 cm SL. Also Ref.: 8631. Known from Sumatra and Java.	<i>Trimma okinawae</i> (Aoyagi 1949)	Nusa Okinawa
<i>Istigobius rigilius</i> (Herre 1953)	Rigilius goby, (M), Ref. 420. Max. 7.9 cm SL. Also Ref.: 8631. Museum: Sulawesi (Celebes), USNM 254295. Also known from Flores.	<i>Valenciennea helsdingenii</i> (Bleeker 1858)	Also F Twostrip Ref.: i
<i>Macrodontogobius wilburi</i> Herre 1936	Large-tooth goby, (M), Ref. 403. Max. 5.2 cm SL.	<i>Valenciennea longipinnis</i> (Lay & Bennett 1839)	Long-fin Also Putri, 1610
<i>Mahidolia mystacina</i> (Valenciennes 1837)	Flagfin prawn goby, (M, Br), Ref. 4343. Max. 6.5 cm SL. Also Ref.: 8631. Known from Flores.	<i>Valenciennea muralis</i> (Valenciennes 1837)	Mural g 8631. 2110
<i>Odontamblyopus rubicundus</i> (Hamilton 1822)	(M, Br), Ref. 7050. Max. 22 cm TL. In range Ref.: 4833.	<i>Valenciennea puellaris</i> (Tomiyama 1955)	149; I Aru Is Maiden Ref.: Kaba P.252
<i>Oligolepis acutipennis</i> (Valenciennes 1837)	Sharptail goby, (M, Br), Ref. 4343. Max. 15 cm TL.	<i>Valenciennea randalli</i> Hoese and Larson 1994	(M), Ref
<i>Oplopomus oplopomus</i> (Valenciennes 1837)	Spinecheek goby, (M), Ref. 8631. Max. 8 cm. Known from Flores.	<i>Valenciennea sexguttata</i> (Valenciennes 1837)	Sixspot 8527 Ment Suma
<i>Oxyurichthys ophthalmonema</i> (Bleeker 1856-57)	Eyebrow goby, (M, Br, Fr), Ref. 5978. Max. 18 cm TL. From Bali Museum: NTM S.10733-002 (TGT1076). Strait to Timor Sea. In range Ref.: 2798.	<i>Valenciennea strigata</i> (Broussonet 1782)	Bluebar Ref.: Sea, Celeb USNI
<i>Papillogobius reichei</i> (Bleeker 1853)	(M, Br, Fr), Ref. 7050. Max. 8.3 cm TL.	<i>Vanderhorstia ambanoro</i> (Fourmanoir 1957)	Ambanc SL. r
<i>Periophthalmodon freycineti</i> Valenciennes 1837	(M), Ref. 5218.	<i>Yongeichthys nebulosus</i> (Forsskål 1775)	Shadow Max. Flora
<i>Periophthalmus argentilineatus</i> Valenciennes 1837	Barred mudskipper, (M, Br), Ref. 5218. Max. 19 cm TL.		
<i>Periophthalmus gracilis</i> Eggert 1935	(M), Ref. 5218.		
<i>Periophthalmus malaccensis</i> Eggert 1935	(M, Br, Fr), Ref. 5218.		
<i>Periophthalmus minutus</i> Eggert 1935	(M), Ref. 5218.		
<i>Platygobiopsis akihito</i> Springer & Randall 1992	(M, En), Ref. 8935. Max. 9.64 cm SL. Museum: USNM 309181 (Maumere Bay off Sao Wisata Resort), AMS I.31467-001, BMNH 1991.5.7.1, BPBM 32817, CAS 760055, NSMT-P 34720, ROM 61628, USNM 309197, NTM S. 13009-001, USNM 316643.		

Table 2. Continuation.
[Tabel 2. Sambungan.]

Haemulidae	Grunts	(Cuvier 1830)	Mu
<i>Diagramma pictum</i> (Thunberg 1792)	Painted sweetlips, (M, Fi, Sport, Dan), Ref. 2112. Max. 100 cm FL. Also Ref.: 6567. Museum: CSIRO CA1649 (conspecific material). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pomadasys maculatus</i> (Bloch 1797)	Su Sado Al
<i>Diagramma punctatum</i> Cuvier 1830	(M), Ref. 5978. Museum: NTM S.11037-001 (TGT2444). From southwest Sumatra to Timor Sea.	<i>Pomadasys opercularis</i> (Playfair & Günther 1866)	(T (R Sma
<i>Plectorhinchus albovittatus</i> (Rüppell 1835)	Two-stripe sweetlips, (M), Ref. 1602. Max. 250 cm SL.	Istiophoridae	Billi
<i>Plectorhinchus celebicus</i> Bleeker 1873	Celebes sweetlips, (M, Fi), Ref. 160. Max. 41 cm SL. Museum: BPBM 29335, from Bali Strait to Timor Sea (Ref. 5978).	<i>Istiophorus platypterus</i> (Shaw & Nodder 1792)	Indo- Ma
<i>Plectorhinchus chaetodonoides</i> Lacepède 1801	Harlequin sweetlips, (M), Ref. 8631. Max. 60 cm SL. In range Ref.: 1602. From southwest Sumatra to Bali Strait (Ref. 5978). Also known from Flores.	<i>Makaira indica</i> (Cuvier 1832)	Black
<i>Plectorhinchus chubbii</i> (Regan 1919)	Dusky rubberlips, (M), Ref. 5978. Max. 75 cm. Museum: LPPL JIF74 (TGT900). From Bali Strait to Timor Sea.	<i>Makaira mazara</i> (Jordan & Snyder 1901)	Indo- Ma
<i>Plectorhinchus flavomaculatus</i> (Ehrenberg 1830)	Lemonfish, (M, Fi), Ref. 5978. Max. 60 cm. From southwest Sumatra to Timor Sea.	<i>Tetrapturus angustirostris</i> Tanaka 1914	Shor TL
<i>Plectorhinchus gaterinoides</i> (Cuvier 1830)	Lined sweetlips, (M), Ref. 8631. Max. 40 cm SL. Known from Flores.	<i>Tetrapturus audax</i> (Philippi 1887)	Strip TL
<i>Plectorhinchus gibbosus</i> (Lacepède 1802)	Harry hotlips, (M, Br, Fr, Sport), Ref. 5978. Max. 75 cm. Museum: WAM P26218-004. From southwest Sumatra to Timor Sea. Also Ref.: 7050.	Kuhliidae	Ahol
<i>Plectorhinchus goldmanni</i> (Bleeker 1853)	Goldman's sweetlips, (M), Ref. 1602. Max. 60 cm SL. Museum: LPPL JIF75, from Bali Strait to Timor Sea (Ref. 5978).	<i>Kuhlia marginata</i> (Cuvier 1829)	Dark cm
<i>Plectorhinchus lineatus</i> (Cuvier 1830)	Yellowbanded sweetlips, (M, Br, Fr), Ref. 5978. Max. 30 cm SL. Museum: BPBM 29350 (TGT2123). From Bali Strait to Timor Sea.	<i>Kuhlia mugil</i> (Schneider 1801)	Barre cm
<i>Plectorhinchus obscurum</i> (Günther 1871)	Giant sweetlips, (M), Ref. 5978. Max. 83 cm SL. Museum: BPBM 29349 (TGT1239). From Bali Strait to Timor Sea.	<i>Kuhlia rupestris</i> (Lacepède 1802)	Rock
<i>Plectorhinchus orientalis</i> (Bloch 1793)	Oriental sweetlips, (M, Sport), Ref. 5978. Max. 72 cm SL. Also Ref.: 8631. Museum: BPBM 29336 (TGT1357). From Bali Strait to Timor Sea. Also known from Flores.	Kurtidae	Nurs
<i>Plectorhinchus picus</i> (Cuvier 1830)	Painted sweetlip, (M, Fi), Ref. 3412. Max. 70 cm SL. Museum: QM I.20291, from Bali Strait to Timor Sea (Ref. 5978).	<i>Kurtus indicus</i> Bloch 1786	India TL
<i>Plectorhinchus polytaenia</i> (Bleeker 1852)	Ribbanded sweetlips, (M), Ref. 5978. Max. 40 cm. Also Ref.: 8631. Museum: LPPL JIF72. From Bali Strait to Timor Sea (Ref. 5978).	Kyphosidae	Sea
<i>Pomadasys argenteus</i> (Forsskål 1775)	Silver grunt, (M, Br, Fi), Ref. 5978. Max. 52 cm. Museum: BPBM 29343. From southwest Sumatra to Timor Sea. Also Ref.: 7050.	<i>Kyphosus cinerascens</i> (Forssål 1775)	Blue Al
<i>Pomadasys argyreus</i> (Valenciennes 1833)	Bluecheek silver grunt, (M), Ref. 5978. Max. 40 cm. Museum: LPPL JIF73. From southwest Sumatra to Timor Sea.	<i>Kyphosus vaigiensis</i> (Quoy & Gaimard 1825)	Str Al Se
<i>Pomadasys furcatus</i> (Bloch & Schneider 1801)	Banded grunt, (M, Fi), Ref. 5978. Max. 50 cm. Museum: BPBM 29348 (TGT3028). From Bali Strait to Timor Sea.	Labridae	Wras
<i>Pomadasys kaakan</i>	Javelin grunter, (M, Br), Ref. 5978. Max. 80 cm.	<i>Anampses caeruleopunctatus</i> Rüppell 1829	Blues Kr
		<i>Anampses geographicus</i> Valenciennes 1840	Geog Kr
		<i>Anampses lineatus</i> Randall 1972	Line Ma
		<i>Anampses melanurus</i> Bleeker 1857	White Al
		<i>Anampses meleagrides</i> Valenciennes 1840	Yellow Al
		<i>Anampses neoguinaicus</i> Bleeker 1878	New
		<i>Anampses twistii</i> Bleeker 1856	Yellow Kr
		<i>Bodianus anthioides</i> (Bennett 1832)	Lyret Ma
		<i>Bodianus axillaris</i> (Bennett 1832)	Axils

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Bodianus bilunulatus</i> (Lacepède 1801)	Tarry hogfish, (M, Sport), Ref. 5978. Max. 50 cm SL. Museum: NTM S.10735-003 (TGT1363). From Bali Strait to Timor Sea. Also Ref.: 8631.	<i>Coris dorsomacula</i> Fowler 1908	Pale
<i>Bodianus bimaculatus</i> Allen 1973	Twospot hogfish, Yellow hogfish, (M), Ref. 1602 Max. 10 cm TL. Flores (Ref. 8631).	<i>Coris gaimard gaimard</i> (Quoy & Gaimard 1824)	Yellow
<i>Bodianus diana</i> (Lacepède 1801)	Diana's hogfish, (M), Ref. 8631. Max. 25 cm SL. Bali.	<i>Coris pictoides</i> Randall & Kuitert 1982	Black
<i>Bodianus mesothorax</i> (Bloch & Schneider 1801)	Splitlevel hogfish, Black-belt hogfish, (M), Ref. 559. Max. 20 cm. Flores (Ref. 8631).	<i>Coris variegata</i> (Rüppell 1835)	Dappled
<i>Cheilinus chlorourus</i> (Bloch 1791)	Floral wrasse, (M), Ref. 2334. Max. 36 cm SL.	<i>Diproctacanthus xanthurus</i> (Bleeker 1856)	Yellow
<i>Cheilinus fasciatus</i> (Bloch 1791)	Redbreasted wrasse, (M), Ref. 1602. Max. 40 cm SL.	<i>Epibulus insidiator</i> (Pallas 1770)	Sling
<i>Cheilinus oxycephalus</i> Bleeker 1853	Snooty wrasse, Point-head maori, (M), Ref. 8631. Max. 14 cm SL. Known from Bali. In range Ref.: 1602, 2334.	<i>Gomphosus varius</i> Lacepède 1801	Bird
<i>Cheilinus trilobatus</i> Lacepède 1801	Tripletail wrasse, Triple-tail maori, (M), Ref. 5978. Max. 40 cm SL. Museum: LPPL JIF119 (TGT909). From Bali Strait to Timor Sea. Also in Flores (Ref. 8631).	<i>Halichoeres argus</i> (Bloch & Schneider 1801)	Argus
<i>Cheilinus undulatus</i> Rüppell 1835	Humphead wrasse, (M, Fi, Dan), Ref. 1602. Max. 229 cm SL.	<i>Halichoeres binotopsis</i> (Bleeker 1849)	(M), F
<i>Cheilio inermis</i> (Forsskål 1775)	Cigar wrasse, (M), Ref. 5978. Max. 50 cm SL. Also Ref.: 8631. Museum: NTM S.10748-015 (TGT1004). From Bali Strait to Timor Sea.	<i>Halichoeres biocellatus</i> Schultz 1960	Redline
<i>Choerodon anchorago</i> (Bloch 1791)	Orange-dotted tuskfish, White-belly tuskfish, (M), Ref. 2334. Max. 38 cm SL. Museum: NTM S.10741-001, from Bali Strait to Timor Sea (Ref. 5978). Also Ref.: 8631.	<i>Halichoeres chloropterus</i> (Bloch 1791)	Paste
<i>Choerodon cephalotes</i> (Castelnau 1875)	Purple tuskfish, (M), Ref. 2334. Max. 38 cm TL.	<i>Halichoeres chrysurus</i> Randall 1981	Almond
<i>Choerodon robustus</i> (Günther 1862)	Robust tuskfish, (M), Ref. 5978. Max. 30 cm TL. From southwest Sumatra to Bali Strait.	<i>Halichoeres dussumieri</i> (Valenciennes 1839)	Bubble
<i>Choerodon schoenleinii</i> (Valenciennes 1839)	Blackspot tuskfish, (M, Sport), Ref. 5978. Max. 100 cm TL. From southwest Sumatra to Timor Sea.	<i>Halichoeres hartzfeldii</i> (Bleeker 1852)	Hartzfeld
<i>Choerodon zamboangae</i> (Seale & Bean 1907)	Purple eyebrowed tuskfish, (M), Ref. 5978. Museum: NTM S.10752-014 (TGT3238). From Bali Strait to Timor Sea.	<i>Halichoeres hortulanus</i> (Lacepède 1801)	Checkered
<i>Cirrhilabrus cyanopleura</i> (Bleeker 1851)	Blueside wrasse, (M), Ref. 2745. Max. 15 cm SL. Also Ref.: 8631. Known from Flores.	<i>Halichoeres margaritaceus</i> (Valenciennes 1839)	Pink-t
<i>Cirrhilabrus exquisitus</i> Smith 1957	Exquisite wrasse, (M), Ref. 5278. Max. 12 cm SL. Also Ref.: 8631.	<i>Halichoeres marginatus</i> Rüppell 1835	Spine
<i>Cirrhilabrus filamentosus</i> (Klausewitz 1976)	Whip-fin wrasse, (M), Ref. 5978. Max. 8 cm. Also Ref.: 8631. Museum: NTM S.10744-001 (TGT955). From Bali Strait to Timor Sea.	<i>Halichoeres melanurus</i> (Bleeker 1851)	Tail-sp
<i>Cirrhilabrus lubbocki</i> Randall & Carpenter 1980	Lubbock's wrasse, (M), Ref. 9823. Max. 7 cm SL. Recorded from Celebes.	<i>Halichoeres melasmapomus</i> Randall 1980	Ref
<i>Cirrhilabrus rubrimarginatus</i> Randall 1992	(M), Ref. 5278. Max. 12.2 cm SL. Museum: off Sulawesi, Tukanbesi Group, Moromaho I., BPBM 34199. Bali, small bay NE of Padangbai, BPBM 30184, 31577.	<i>Halichoeres miniatus</i> (Valenciennes 1839)	Ocella
<i>Cirrhilabrus temminckii</i> Bleeker 1853	Threadfin wrasse, (M), Ref. 8631. Max. 9.9 cm SL. Known from Bali.	<i>Halichoeres nebulosus</i> (Valenciennes 1839)	Almond
<i>Coris aygula</i> Lacepède 1801	Clown wrasse, (M), Ref. 1602. Max. 120 cm TL.	<i>Halichoeres omatissimus</i> (Garrett 1863)	Mus
		<i>Halichoeres podostigma</i> (Bleeker 1854)	Sea
		<i>Halichoeres prosopoeion</i> (Bleeker 1853)	(M), R
			Kno
			Axil sp
			Also
			Flora
			Twotor
			Also

Table 2. Continuation.
[Tabel 2. Sambungan.]

	<i>Halichoeres purpurascens</i> (Bloch & Schneider 1801)	Silty wrasse, (M), Ref. 6023. Max. 12 cm TL. From Flores (Ref. 8631).	<i>Novaculichthys taeniourus</i> (Lacepède 1801)	Roc A (T S
	<i>Halichoeres richmondi</i> Fowler & Bean 1928	Richmond's wrasse, (M), Ref. 1602. Max. 13 cm SL. Also Ref.: 8631. Known from Moluccas and Flores.		Two M S
	<i>Halichoeres scapularis</i> (Bennett 1832)	Zigzag wrasse, (M), Ref. 8631. Max. 15.6 cm SL. Known from Bali.	<i>Oxycheilinus bimaculatus</i> (Valenciennes 1840)	Cele M Che
	<i>Halichoeres solorensis</i> (Bleeker 1853)	Green wrasse, (M), Ref. 6023. Max. 18 cm TL. From Flores (Ref. 8631).	<i>Oxycheilinus celebicus</i> (Bleeker 1853)	Ori M S
	<i>Halichoeres timorensis</i> (Bleeker 1852)	Timor wrasse, (M), Ref. 2136. Max. 7.8 cm SL. Also Ref.: 8631. Known from Bali.	<i>Oxycheilinus digrammus</i> (Lacepède 1801)	Rin T
	<i>Halichoeres trimaculatus</i> (Quoy & Gaimard 1824)	Threespot wrasse, (M), Ref. 1602. Max. 22 cm SL.	<i>Oxycheilinus orientalis</i> (Günther 1862)	Pink M S
	<i>Halichoeres zeylonicus</i> Bennett 1832	Goldstripe wrasse, (M), Ref. 2334. Max. 20 cm.	<i>Oxycheilinus unifasciatus</i> (Streets 1877)	Ring T
	<i>Hemigymnus fasciatus</i> (Bloch 1792)	Barred thicklip, (M), Ref. 8631. Max. 50 cm SL. Known from Bali and Flores.	<i>Paracheilinus carpenteri</i> Randall & Lubbock 1981	Par M S
	<i>Hemigymnus melapterus</i> (Bloch 1791)	Blackeye thicklip, (M), Ref. 5978. Max. 71 cm SL. Also Ref.: 8631. Museum: LPPL JIF121 (TGT1551). From Bali Strait to Timor Sea.	<i>Paracheilinus filamentosus</i> Allen 1974	Fila A
	<i>Hologymnosus annulatus</i> (Lacepède 1801)	Ring wrasse, (M), Ref. 2334. Max. 40 cm TL.	<i>Pseudocheilinus ataenia</i> Schultz 1960	Pel A C
	<i>Hologymnosus doliatus</i> (Lacepède 1801)	Pastel ringwrasse, (M), Ref. 8631. Max. 38 cm TL. Known from Flores.	<i>Pseudocheilinus evanidus</i> Jordan & Evermann 1903	Stria K
	<i>Hologymnosus rhodonotus</i> Randall & Yamakawa 1988	Redback longface wrasse, (M), Ref. 5277. Max. 27.4 cm SL. From Bali Strait to Timor (Ref. 5978). Museum: Bali, BPBM 31973.	<i>Pseudocheilinus hexataenia</i> (Bleeker 1857)	Pyja Fl
251	<i>Labrichthys unilineatus</i> (Guichenot 1847)	Tubelip wrasse, (M), Ref. 2747. Max. 17.5 cm TL. Flores (Ref. 8631).	<i>Pseudocheilinus octotaenia</i> Jenkins 1901	Eigh K
	<i>Labroides bicolor</i> Fowler & Bean 1928	Bicolor cleaner wrasse, (M), Ref. 1602. Max. 9.8 cm SL.	<i>Pseudocoris heteroptera</i> (Bleeker 1857)	Torp
	<i>Labroides dimidiatus</i> (Valenciennes 1839)	Bluestreak cleaner wrasse, (M), Ref. 1602. Max. 11.5 cm TL. Also Ref.: 8631. Museum: NTM S.10736-001 (TGT1757). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pseudocoris philippina</i> (Fowler & Bean 1928)	Phili K
	<i>Labroides pectoralis</i> Randall & Springer 1975	Blackspot cleaner wrasse, (M), Ref. 8631. Max. 6 cm SL. Known from Flores.	<i>Pseudocoris yamashiroi</i> (Schmidt 1930)	Red K
	<i>Labropsis alleni</i> Randall 1981	Allen's tubelip, (M), Ref. 2137. Max. 8.2 cm SL. Also Ref.: 8631. Known from Flores. Museum: Molucca Is., Ambon, BPBM 19303.	<i>Pseudodax moluccanus</i> (Valenciennes 1840)	Chis R (F
	<i>Labropsis manabei</i> Schmidt 1930	Northern tubelip, (M), Ref. 8631. Max. 11.7 cm SL. Known from Flores.	<i>Pteragogus amboinensis</i> (Bleeker 1856)	(M),
	<i>Labropsis xanthonota</i> Randall 1981	Yellowback tubelip, (M), Ref. 2137. Max. 13 cm TL. Also Ref.: 8631. Known from Flores. Museum: Java, Seribu Is., Pulau Putri, BPBM 19522.	<i>Pteragogus cryptus</i> Randall 1981	Cryp M
	<i>Leptojulius cyanopleura</i> (Bleeker 1853)	Shoulder-spot wrasse, (M), Ref. 5978. Max. 13 cm TL. Also Ref.: 8631, 9823. Museum: AMS I.22806-019. From Bali Strait to Timor Sea. In range Ref.: 3132.	<i>Pteragogus flagellifer</i> (Valenciennes 1839)	Coch M S
	<i>Macropharyngodon negrosensis</i> Herre 1932	Yellowspotted wrasse, (M), Ref. 8631. Max. 10.2 cm SL. Known from Flores.	<i>Pteragogus guttatus</i> (Fowler & Bean 1928)	(M),
	<i>Macropharyngodon ornatus</i> Randall 1978	Ornate wrasse, False leopard, (M), Ref. 6192. Max. 12 cm TL. Also from Flores (Ref. 8631). Museum: Molucca Is., Ambon, BPBM 18546 (Holotype).	<i>Stethojulis bandanensis</i> (Bleeker 1851)	Red
	<i>Novaculichthys macrolepidotus</i> (Bloch 1791)	Seagrass wrasse, (M), Ref. 1602. Max. 12 cm SL.	<i>Stethojulis interrupta</i> (Bleeker 1851)	Cutr M Ti

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Stethojulis strigiventer</i> (Bennett 1832)	Three-ribbon rainbowfish, (M), Ref. 8631. Max. 15 cm TL. Known from Flores.	<i>Leiognathus blochi</i> (Valenciennes 1835)	From s Twobloto
<i>Stethojulis trilineata</i> (Bloch & Schneider 1801)	Threeline rainbowfish, (M, Fi), Ref. 2334. Max. 15 cm TL. Also Ref.: 8631. Known from Flores.	<i>Leiognathus decorus</i> (De Vis 1844)	TL. (M, Br), F 1984.
<i>Thalassoma amblycephalum</i> (Bleeker 1856)	Bluntheaded wrasse, (M), Ref. 8631. Max. 16 cm SL. Known from Bali.	<i>Leiognathus dussumieri</i> (Valenciennes 1835)	Sumat Dussumi
<i>Thalassoma hardwickii</i> (Bennett 1828-30)	Sixbar wrasse, (M), Ref. 8631. Max. 17 cm SL. Known from Bali.	<i>Leiognathus elongatus</i> (Günther 1874)	Max. 1 Slender p Museu
<i>Thalassoma janseni</i> (Bleeker 1856)	Jansen's wrasse, (M), Ref. 8631. Max. 17 cm SL. Known from Bali and Flores.	<i>Leiognathus equulus</i> (Forsskål 1775)	southw Commor
<i>Thalassoma lunare</i> (Linnaeus 1758)	Moon wrasse, (M), Ref. 8631. Max. 30 cm. Known from Bali.	<i>Leiognathus fasciatus</i> (Lacepède 1803)	TL. M Sumat
<i>Thalassoma purpureum</i> (Forsskål 1775)	Surge wrasse, (M), Ref. 8631. Max. 46 cm. Known from Bali.	<i>Leiognathus leuciscus</i> (Günther 1860)	Striped p Museu to Tim
<i>Thalassoma trilobatum</i> (Lacepède 1801)	Christmas wrasse, (M), Ref. 1602. Max. 24 cm SL.	<i>Leiognathus rapsoni</i> Munro 1964	Whipfin p Museu From :
<i>Wetmorella albofasciata</i> Schultz & Marshall 1954	Whitebanded sharpnose wrasse, (M), Ref. 1602. Max. 4.6 cm SL.	<i>Leiognathus smithursti</i> (Ramsay & Ogilby 1886)	Rapson's Museu From :
<i>Wetmorella nigropinnata</i> (Seale 1901)	Sharpnose wrasse, (M), Ref. 2138. Max. 6.5 cm SL. Also Ref.: 8631. Known from Flores. Museum: Pulau Seribu, Java, USNM 231377. Marsegoe Bay, Moluccas, AMS I.18469-082. Ceram, USNM 209661, 210030. Kabaena I., USNM 236520. Huruku I., Moluccas, USNM 209587.	<i>Leiognathus splendens</i> (Cuvier 1829)	Smithurs Museu From :
<i>Xiphocheilus typus</i> Bleeker 1856	Blue-banded wrasse, (M), Ref. 5978. Max. 12 cm SL. Museum: NTM S.11016-004 (TGT2473). From southwest Sumatra to Bali Strait.	<i>Leiognathus stercorarius</i> Evermann & Seale 1907	Splendid Also F southw
<i>Xyrichtys aneitensis</i> (Günther 1862)	Yellowblotch razorfish, (M), Ref. 1602. Max. 20 cm SL. Also Ref.: 8631. Known from Bali.	<i>Secutor indicus</i> Monkolprasit 1973	Oblong s Museu to Bali
<i>Xyrichtys celebicus</i> (Bleeker 1856)	Celebes razorfish, (M), Ref. 1602. Max. 16.1 cm SL.	<i>Secutor insidiator</i> (Bloch 1787)	(M, Br), I JIF21: Sea. F 7050).
<i>Xyrichtys dea</i> Temminck & Schlegel 1845	(M), Ref. 5978. Max. 30 cm TL. Museum: CSIRO CA2145. From Bali Strait to Timor Sea.	<i>Secutor ruconius</i> (Hamilton 1822)	Pugnose SL. M southw Ref.: 6
<i>Xyrichtys pavo</i> Valenciennes 1840	Peacock wrasse, (M, Sport), Ref. 5978. Max. 35 cm TL. Museum: NTM S.10745-002 (TGT1888). From southwest Sumatra to Timor Sea.		Deep pu Museu Suma
<i>Xyrichtys pentadactylus</i> (Linnaeus 1758)	Five-finger wrasse, (M), Ref. 5978. Max. 25 cm TL. Also Ref.: 8631. Museum: LPPL JIF123 (TGT1592). From Bali Strait to Timor Sea.		Empero
<i>Xyrichtys twistii</i> (Bleeker 1856)	(M), Ref. 559. Max. 20 cm SL.		Striped li 30 cm southw
Lactariidae	False trevallies	Lethrinidae	
<i>Lactarius lactarius</i> (Bloch & Schneider 1801)	False trevally, (M, Fi), Ref. 6567. Max. 40 cm. Museum: LPPL JIF46 (TGT2503). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Gnathodentex aureolineatus</i> (Lacepède 1802)	Forktail l TL. M Suma
Leiognathidae	Slimys, slipmouths, or ponyfishes	<i>Gymnocranius elongatus</i> Senta 1973	Yellowsn Max. : Blue-line cm TL
<i>Gazza aklamys</i> Jordan and Starks 1917	Smalltoothed ponyfish, (M, Br, Fi), Ref. 5978. Max. 15 cm SL. Museum: NTM S.11040-001 (TGT2504). From southwest Sumatra to Timor Sea.	<i>Gymnocranius frenatus</i> Bleeker 1873	
<i>Gazza minuta</i> (Bloch 1797)	Toothpony, (M, Br, Fi), Ref. 5978. Max. 21 cm FL. Museum: NTM S.11031-003 (TGT2615). From southwest Sumatra to Timor Sea. Also Ref.: 7050.	<i>Gymnocranius grandoculis</i> (Valenciennes 1830)	
<i>Leiognathus bindus</i> (Valenciennes 1835)	Orangefin ponyfish, (M, Br), Ref. 5978. Max. 11 cm. Museum: NTM S.10733-042 (TGT2108).		

Table 2. Continuation.
[Tabel 2. Sambungan.]

		from southwest Sumatra to Timor Sea (Ref. 5978). Grey large-eye bream, (M, Fi), Ref. 2295. Max. 35 cm TL. Museum: LPPL JIF195 (TGT2369). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Wattisia mossambica</i> (Smith 1957)	Moz Ma (R
	<i>Gymnocranius griseus</i> (Temminck & Schlegel 1843)			
	<i>Gymnocranius microdon</i> (Bleeker 1851)	Blue-spotted large-eye bream, (M, Fi), Ref. 2295. Max. 45 cm TL.	Lobotidae <i>Lobotes surinamensis</i> (Bloch 1790)	Tripl Atlan Ma
	<i>Lethrinus amboinensis</i> Bleeker 1854	Ambon emperor, (M, Fi), Ref. 2295. Max. 70 cm TL.	Lutjanidae <i>Aphareus furca</i> (Lacepède 1801)	Snape Sma Ma
	<i>Lethrinus atkinsoni</i> Seale 1910	Pacific yellowtail emperor, (M, Fi, Sport), Ref. 2295. Max. 45 cm TL.	<i>Aphareus rutilans</i> Cuvier 1830	Rust Ma to
	<i>Lethrinus conchylatus</i> (Smith 1959)	Redaxil emperor, (M, Fi), Ref. 2295. Max. 76 cm TL.	<i>Aprion virescens</i> Valenciennes 1830	Gre SL (T (R
	<i>Lethrinus erythracanthus</i> Valenciennes 1830	Orange-spotted emperor, (M, Fi, Dan), Ref. 2295. Max. 70 cm TL.		
	<i>Lethrinus erythropterus</i> Valenciennes 1830	Longfin emperor, (M, Fi), Ref. 2295. Max. 50 cm TL. From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Etelis carbunculus</i> Cuvier 1828	Ruby Ma Se
	<i>Lethrinus genivittatus</i> Valenciennes 1830	Longspine emperor, (M, Br, Fi), Ref. 2295. Max. 25 cm TL. From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Etelis coruscans</i> Valenciennes 1862	Flam
	<i>Lethrinus harak</i> (Forsskål 1775)	Thumbprint emperor, (M, Fi), Ref. 2295. Max. 50 cm TL.	<i>Etelis radiosus</i> Anderson 1981	Scar Ma Su
	<i>Lethrinus laticaudis</i> Alleyne & Macleay 1877	Grass emperor, (M, Br, Fi, Sport), Ref. 2295. Max. 56 cm TL.	<i>Lipocheilus carnolabrum</i> (Chah 1970)	Tang
	<i>Lethrinus lentjan</i> (Lacepède 1802)	Pink ear emperor, (M, Fi), Ref. 2295. Max. 50 cm TL. Museum: WAMRL Leth Bali 6/83 (TGT2406). From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Lutjanus argentimaculatus</i> (Forsskål 1775)	Man cn (T (R
253	<i>Lethrinus microdon</i> Valenciennes 1830	Smalltooth emperor, (M, Fi), Ref. 2295. Max. 70 cm TL. Museum: LPPL JIF198, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Lutjanus bengalensis</i> (Bloch 1790)	Ben M Su
	<i>Lethrinus nebulosus</i> (Forsskål 1775)	Spangled emperor, (M, Fi, Sport), Ref. 2295. Max. 80 cm TL. From Bali Strait to Timor Sea (Ref. 5978).	<i>Lutjanus biguttatus</i> (Valenciennes 1830)	Two- TL
	<i>Lethrinus obsoletus</i> (Forsskål 1775)	Orange-striped emperor, (M, Fi), Ref. 2295. Max. 60 cm TL.	<i>Lutjanus bitaeniatus</i> (Valenciennes 1830)	Indo M Su
	<i>Lethrinus olivaceus</i> Valenciennes 1830	Longface emperor, (M, Fi, Dan), Ref. 2295. Max. 100 cm TL.	<i>Lutjanus bohar</i> (Forsskål 1775)	Two- 75 (T (F
	<i>Lethrinus ornatus</i> Valenciennes 1830	Ornate emperor, (M, Fi), Ref. 2295. Max. 40 cm TL. Museum: LPPL JIF199, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Lutjanus bouton</i> (Lacepède 1802)	Mol R Se
	<i>Lethrinus reticulatus</i> Valenciennes 1830	Red snout emperor, (M, Fi), Ref. 2295. Max. 40 cm TL.	<i>Lutjanus carponotatus</i> (Richardson 1842)	Spar M
	<i>Lethrinus rubrioperculatus</i> Sato 1978	Spotcheek emperor, (M, Fi), Ref. 2295. Max. 50 cm TL. Museum: WAMRL Leth Bali 10/83, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Lutjanus decussatus</i> (Cuvier 1828)	Chec Al fr in- Sunt
	<i>Lethrinus semicinctus</i> Valenciennes 1830	Black blotch emperor, (M, Fi), Ref. 2295. Max. 35 cm TL. Museum: CSIRO CA948, from Bali Strait to Timor Sea (Ref. 5978).	<i>Lutjanus dodecacanthoides</i> (Bleeker 1854)	Blac AI
	<i>Lethrinus variegatus</i> Valenciennes 1830	Slender emperor, (M, Fi), Ref. 2295. Max. 20 cm TL. Museum: CSIRO CA968, from Bali Strait to Timor Sea (Ref. 5978).	<i>Lutjanus ehrenbergii</i> (Peters 1869)	
	<i>Lethrinus xanthochilus</i> Klunzinger 1870	Yellowlip emperor, (M, Fi, Sport), Ref. 2295. Max. 60 cm TL.		
	<i>Monotaxis grandoculis</i> (Forsskål 1775)	Humpnose big-eye bream, (M, Fi, Dan), Ref. 2295. Max. 60 cm TL. Museum: LPPL JIF200, from southwest Sumatra to Timor Sea (Ref. 5978).		

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

254

<i>Lutjanus erythropterus</i> Bloch 1790	Crimson snapper, Bambang, (M, Fi), Ref. 55. Max. 60 cm TL. Also Ref.: 6567, Museum: LPPL JIF68 (TGT2258). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Lutjanus semicinctus</i> Quoy & Gaimard 1824	Black-b...
<i>Lutjanus fulviflamma</i> (Forsskål 1775)	Blackspot snapper, (M, Fi), Ref. 55. Max. 35 cm TL. Museum: LPPL JIF69, from Bali Strait to Timor Sea (Ref. 5978).	<i>Lutjanus timorensis</i> (Quoy & Gaimard 1824)	Timor s Ref.: Sum Flora
<i>Lutjanus fulvus</i> (Schneider 1801)	Blacktail snapper, (M, Fi, Sport, Dan), Ref. 55. Max. 40 cm TL. Also Ref.: 8631. Museum: LPPL JIF214 (TGT1015). From Bali Strait to Timor Sea (Ref. 5978).	<i>Lutjanus vitta</i> (Quoy & Gaimard 1824)	Browns TL. M Timor
<i>Lutjanus gibbus</i> (Forsskål 1775)	Humpback snapper, (M, Fi, Sport, Dan), Ref. 55. Max. 50 cm TL. Also Ref.: 8631. From Bali Strait to Timor Sea (Ref. 5978).	<i>Macolor macularis</i> Fowler 1931	Midnigh Also
<i>Lutjanus johnii</i> (Bloch 1792)	John's snapper, (M, Fi, Sport), Ref. 55. Max. 70 cm TL. Also Ref.: 8631. From southwest Sumatra to Timor Sea (Ref. 5978); including Flores.	<i>Macolor niger</i> (Forsskål 1775)	Black a cm S Bali
<i>Lutjanus kasmira</i> (Forsskål 1775)	Common bluestripe snapper, (M, Fi, Sport), Ref. 55. Max. 40 cm TL. Also Ref.: 8631. Known from Bali.	<i>Paracaesio kusakarii</i> Abe 1960	Saddle- Muse to Tim
<i>Lutjanus lemniscatus</i> (Valenciennes 1828)	Yellowstreaked snapper, (M, Fi), Ref. 55. Max. 65 cm TL. Museum: LPPL JIF54; from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Paracaesio sordida</i> Abe & Shinohara 1962	Dirty or
<i>Lutjanus lunulatus</i> (Park 1797)	Lunartail snapper, (M, Fi), Ref. 55. Max. 35 cm TL. Museum: BPBM 29338, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Paracaesio xanthura</i> (Bleeker 1869)	Yelloww GME From
<i>Lutjanus lutjanus</i> Bloch 1790	Bigeye snapper, (M, Fi), Ref. 55. Max. 30 cm TL. Also Ref.: 8631. Museum: LPPL JIF55 (TGT2359). From southwest Sumatra to Timor Sea (Ref. 5978); including Flores.	<i>Pinjalo lewisi</i> Randall, Allen, & Anderson 1987	Slender
<i>Lutjanus madras</i> (Valenciennes 1831)	Indian snapper, (M, Fi), Ref. 55. Max. 30 cm TL. Museum: LPPL JIF56, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pinjalo pinjalo</i> (Bleeker 1850)	Pinjalo, JIF6 5978
<i>Lutjanus malabaricus</i> (Bloch & Schneider 1801)	Malabar blood snapper, (M, Fi, Sport), Ref. 55. Max. 100 cm TL. Also Ref.: 6567, 9987. Museum: LPPL JIF57, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pristipomoides argyrogrammicus</i> (Valenciennes 1832)	Ornate
<i>Lutjanus mizenkoi</i> Allen & Talbot 1985	Samoan snapper, (M, Fi), Ref. 55. Max. 30 cm TL.	<i>Pristipomoides auricilla</i> (Jordan, Evermann & Tanaka 1927)	Goldfla
<i>Lutjanus monostigma</i> (Cuvier 1828)	One-spot snapper, (M, Fi, Dan), Ref. 55. Max. 60 cm TL. Also Ref.: 8631. Museum: LPPL JIF58 (TGT2405). From southwest Sumatra to Timor Sea (Ref. 5978); also Bali.	<i>Pristipomoides filamentosus</i> (Valenciennes 1830)	Crimso Muse (Ref.)
<i>Lutjanus quinquelineatus</i> (Bloch 1790)	Five-lined snapper, (M, Fi, Sport), Ref. 55. Max. 38 cm TL. Museum: BPBM 29339, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pristipomoides flavipinnis</i> Shinohara 1963	Golden
<i>Lutjanus rivulatus</i> (Cuvier 1828)	Blubberlip snapper, (M, Fi), Ref. 55. Max. 65 cm TL. Also Ref.: 8631. Museum: LPPL JIF60 (TGT2344). From southwest Sumatra to Bali Strait (Ref. 5978); including Flores.	<i>Pristipomoides multidentis</i> (Day 1870)	Goldba Muse Timor
<i>Lutjanus russelli</i> (Bleeker 1849)	Russell's snapper, (M, Fi), Ref. 55. Max. 50 cm TL. Also Ref.: 8631. Museum: LPPL JIF61, from Bali Strait to Timor Sea (Ref. 5978). Also known from Flores.	<i>Pristipomoides sieboldii</i> (Bleeker 1857)	Lavend SL. M Sea.
<i>Lutjanus sebae</i> (Cuvier 1816)	Emperor red snapper, (M, Fi, Sport), Ref. 55. Max. 100 cm TL. Museum: LPPL JIF62, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pristipomoides typus</i> Bleeker 1852	Sharpt Ref.: Sum
		<i>Pristipomoides zonatus</i> (Valenciennes 1830)	Oblique Max.
		<i>Symphorichthys spilurus</i> (Günther 1874)	Sailfin s Ref.: Bali
		<i>Symphorus nematophorus</i> (Bleeker 1860)	Chinam SL. A (TGT inclu

Table 2. Continuation.
[Tabel 2. Sambungan.]

<p>Malacanthidae <i>Branchiostegus australiensis</i> Dooley & Kailola 1988 <i>Branchiostegus gloerfelti</i> Dooley & Kailola 1988</p> <p><i>Hoplaltilus cuniculus</i> Randall & Dooley 1974 <i>Hoplaltilus fourmanoiri</i> Smith 1964 <i>Hoplaltilus luteus</i> Allen & Kuitert 1989 <i>Hoplaltilus marcosi</i> Burgess 1977 <i>Hoplaltilus starcki</i> Randall & Dooley 1974</p> <p><i>Malacanthus brevirostris</i> Guichenot 1848</p> <p><i>Malacanthus latovittatus</i> (Lacepède 1801)</p>	<p>Tilefishes (M), Ref. 9069. Max. 26.6 cm SL. From southwest Sumatra to Bali Strait (Ref. 5978). (M), Ref. 9069. Max. 24 cm SL. From southwest Sumatra to Bali Strait. Museum: WAM P.28304-001 (Holotype). Dusky tilefish, (M), Ref. 8631. Max. 12.9 cm SL. Known from Flores. Yellow-spotted tilefish, (M), Ref. 8991. Max. 13.7 cm TL. Yellow tilefish, Yellow-spotted tilefish, (M), Ref. 8989. Max. 10 cm. From Flores (Ref. 8631). Redback sand tilefish, (M), Ref. 9137. Max. 10 cm TL. Stark's tilefish, (M), Ref. 8631. Max. 11.6 cm SL. Also Ref.: 1602. Known from Flores and the Moluccas. Quakerfish, (M), Ref. 5978. Max. 30 cm TL. Also Ref.: 8991. Museum: BMNH uncatalogued: (PJPW2043). ZMA 111.185; RMNH 184; AMS IA.7011; AMS IB.129. From southwest Sumatra to Bali Strait. Blue blanquillo, (M), Ref. 5978. Max. 45 cm SL. Also Ref.: 8631, 8991. Museum: LPPL JIF45 (TGT3033). ZMA 111.184; RMNH 16012; USNM 216683 - 4. From Bali Strait to Timor Sea; including Flores.</p>	<p><i>Ptereleotris microlepis</i> (Bleeker 1856) <i>Ptereleotris monopectera</i> Randall & Hoese 1985 <i>Ptereleotris uroditaenia</i> Randall & Hoese 1985 <i>Ptereleotris zebra</i> (Fowler 1938)</p> <p>Monodactylidae <i>Monodactylus argenteus</i> (Linnaeus 1758)</p> <p>Mugilidae <i>Crenimugil heterocheilus</i> (Bleeker 1855) <i>Liza alata</i> (Steindachner 1892) <i>Liza parrata</i> (Cantor 1849) <i>Liza subviridis</i> (Valenciennes 1836) <i>Liza tade</i> (Forsskål 1775) <i>Valamugil cunnesius</i> (Valenciennes 1836) <i>Valamugil speigleri</i> (Bleeker 1858)</p>
<p>Menidae <i>Mene maculata</i> (Bloch & Schneider 1801)</p>	<p>Moonfish Moonfish, (M, Br, Fi), Ref. 5978. Max. 30 cm TL. Museum: LPPL JIF49 (TGT2223). From southwest Sumatra to Timor Sea.</p>	<p>Mullidae <i>Mulloidichthys flavolineatus</i> (Lacepède 1801)</p>
<p>Microdesmidae <i>Gunnellichthys curiosus</i> Dawson 1968 <i>Gunnellichthys monostigma</i> Smith 1958 <i>Gunnellichthys pleurotaenia</i> Bleeker 1858 <i>Nemateleotris decora</i> Randall & Allen 1973 <i>Nemateleotris magnifica</i> Fowler 1938 <i>Parioglossus nudus</i> Rennis & Hoese 1985 <i>Parioglossus palustris</i> (Herre 1945) <i>Parioglossus raoi</i> (Herre 1939) <i>Ptereleotris evides</i> (Jordan & Hubbs 1925) <i>Ptereleotris grammica</i> Randall & Lubbock 1982 <i>Ptereleotris hanae</i> (Jordan & Snyder 1901) <i>Ptereleotris heteroptera</i> (Bleeker 1855)</p>	<p>Wormfishes Curious wormfish, Neon worm-goby, (M), Ref. 2334 Max. 11.5 cm TL. From Flores (Ref. 8631). Onespot wormfish, Black-spot worm-goby, (M), Ref. 2334 Max. 11 cm TL. From Bali (Ref. 8631). Onestripe wormfish, (M), Ref. 1602. Max. 7.4 cm SL. Also Ref.: 8631. Known from Java and Flores. Elegant firefish, (M), Ref. 8631. Max. 7.5 cm TL. Known from Flores. Fire goby, (M), Ref. 8631. Max. 9 cm TL. Known from Flores. (M), Ref. 1602. (M, Br), Ref. 1602. (M, Br), Ref. 1602. Max. 3.1 cm SL. Blackfin dartfish, (M), Ref. 528. Max. 12 cm TL. Also Ref.: 8631. Known from Flores. Lined dartfish, (M), Ref. 528. Max. 10 cm. Blue hana goby, (M), Ref. 8631. Max. 10 cm SL. Known from Bali. Blacktail goby, (M), Ref. 528. Max. 14 cm TL. Also Ref.: 8631. Known from Flores.</p>	<p><i>Mulloidichthys vanicolensis</i> (Valenciennes 1831) <i>Parupeneus barberinoides</i> (Bleeker 1852)</p> <p><i>Parupeneus barberinus</i> (Lacepède 1801)</p> <p><i>Parupeneus bifasciatus</i> (Lacepède 1801)</p> <p><i>Parupeneus chrysopleuron</i> (Temminck & Schlegel 1843)</p> <p><i>Parupeneus ciliatus</i> (Lacepède 1802) <i>Parupeneus cyclostomus</i> (Lacepède 1801)</p> <p><i>Parupeneus heptacanthus</i> (Lacepède 1801)</p> <p><i>Parupeneus indicus</i> (Shaw 1803)</p>

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	Sumatra to Timor Sea.		
<i>Parupeneus macronema</i> (Lacepède 1801)	Long-barbel goatfish, (M), Ref. 5978. Max. 35 cm. Also Ref.: 8631. In range Ref.: 5405. Museum: LPPL JIF83 (TGT2307). From southwest Sumatra to Timor Sea; including Bali.	<i>Nemipterus bathybius</i> Snyder 1911	Timor Yellow
<i>Parupeneus multifasciatus</i> (Quoy & Gaimard 1824)	Manybar goatfish, (M, Fi), Ref. 8631. Max. 30 cm TL. Known from Flores.	<i>Nemipterus bipunctatus</i> (Ehrenberg 1830)	Malay (Timor) (R)
<i>Parupeneus pleurostigma</i> (Bennett 1831)	Sidespot goatfish, (M, Fi, Sport), Ref. 5978. Max. 25 cm SL. Also Ref.: 8631. Museum: LPPL JIF85 (TGT963). From Bali Strait to Timor Sea; including Flores.	<i>Nemipterus celebicus</i> (Bleeker 1854)	Delagoa SL Celeb SL Tim
<i>Parupeneus rubescens</i> (Lacepède 1801)	Rosy goatfish, (M), Ref. 5978. Max. 42 cm. Museum: LPPL JIF86 (TGT3320). From Bali Strait to Timor Sea.	<i>Nemipterus furcosus</i> (Valenciennes 1830)	Fork- cm Su
<i>Parupeneus spilurus</i> (Bleeker 1854)	(M), Ref. 9947. Max. 36 cm TL.	<i>Nemipterus gracilis</i> (Bleeker 1873)	Grace Me (Timor) (R)
<i>Upeneus asymmetricus</i> Lachner 1954	Asymmetrical goatfish, (M), Ref. 3132. Max. 30 cm TL.	<i>Nemipterus hexodon</i> (Quoy & Gaimard 1824)	Ornat SL Tim
<i>Upeneus bensasi</i> (Temminck & Schlegel 1843)	Bensasi goatfish, (M, Fi), Ref. 5978. Max. 20 cm. Museum: NTM S.10995-003 (TGT2591). From southwest Sumatra to Timor Sea.	<i>Nemipterus isacanthus</i> (Bleeker 1873)	Teard SL
<i>Upeneus luzonius</i> Jordan & Seale 1907	Darkbarred goatfish, (M), Ref. 5978. Also Ref.: 8631. Museum: CSIRO CA2057 (conspecific material). From Bali Strait to Timor Sea.	<i>Nemipterus japonicus</i> (Bloch 1791)	Japan Ma Ba
<i>Upeneus moluccensis</i> (Bleeker 1855)	Goldband goldfish, Bidji nangka, (M, Fi), Ref. 2110. Max. 30 cm. Also Ref.: 6567. Museum: LPPL JIF87 (TGT2432). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Nemipterus marginatus</i> (Valenciennes 1830)	Red f 15 sou
<i>Upeneus quadrilineatus</i> Cheng & Wang 1963	(M), Ref. 5978. Max. 17 cm SL. Museum: NTM S.10748-012 (TGT1017). From southwest Sumatra to Timor Sea.	<i>Nemipterus mesoprion</i> (Bleeker 1853)	Mauv cm Su
<i>Upeneus sulphureus</i> Cuvier 1829	Sulphur goatfish, (M, Br, Fi), Ref. 5978. Max. 23 cm. Museum: LPPL JIF88. From southwest Sumatra to Timor Sea.	<i>Nemipterus nematophorus</i> (Bleeker 1853)	Doub cm Su
<i>Upeneus sundaicus</i> (Bleeker 1855)	Ochre-banded goatfish, Kunir, (M, Br), Ref. 4899. Max. 22 cm.	<i>Nemipterus nematopus</i> (Bleeker 1851)	Yellow 17 Str
<i>Upeneus tragula</i> Richardson 1846	Freckled goatfish, (M, Br, Fi), Ref. 5978. Max. 33 cm TL. Also Ref.: 8631. Museum: LPPL JIF89 (TGT1061). From southwest Sumatra to Bali Strait; including Flores.	<i>Nemipterus nemurus</i> (Bleeker 1857)	Redsq cm
<i>Upeneus vittatus</i> (Forsskål 1775)	Yellowstriped goatfish, (M), Ref. 5978. Max. 30 cm. Also Ref.: 8631. Museum: LPPL JIF90. From southwest Sumatra to Timor Sea; including Flores.	<i>Nemipterus peronii</i> (Valenciennes 1830)	Notch 26 Fro
Nemipteridae	Threadfin breams, Whiptail breams	<i>Nemipterus tambuloides</i> (Bleeker 1853)	Fivelin cm 00 59
<i>Nemipterus aurora</i> Russell 1993	Dawn threadfin bream, (M), Ref. 9785. Max. 20 cm SL.	<i>Nemipterus thosaporni</i> Russell 1991	Palefi cm
<i>Nemipterus balinensis</i> (Bleeker 1858-9)	Balinese threadfin bream, (M, Fi, En), Ref. 3810. Max. 18 cm SL. Museum: NTM S.10824-002, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Nemipterus virgatus</i> (Houttuyn 1782)	Gold SL Su
<i>Nemipterus balinensoides</i> (Popta 1918)	Dwarf threadfin bream, (M, Fi), Ref. 3810. Max. 12.5 cm SL. Museum: NTM S.11001-002 (TGT2560) as <i>Nemipterus</i> sp.2; NTM S.10733-006 as <i>Nemipterus</i> sp. 4. From southwest Sumatra to	<i>Nemipterus zysron</i> (Bleeker 1856-57)	Slend Ma (Timor)
		<i>Parascalopsis eriomma</i>	Rosy

Table 2. Continuation.
[Tabel 2. Sambungan.]

(Jordan & Richardson 1909)	25.5 cm SL. Museum: NTM S.10995-001 (TGT2592). From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Scolopsis margaritifer</i> (Cuvier 1830)	Pearly Ma S.1 597
<i>Parascolopsis inermis</i> (Temminck & Schlegel 1843)	Unarmed dwarf monocle bream, (M, Fi), Ref. 3810. Max. 18 cm SL. Museum: NTM S.10752-005, from Bali Strait to Timor Sea (Ref. 5978).	<i>Scolopsis monogramma</i> (Kuhl & Van Hasselt 1830)	Monogram 381 Tim
<i>Parascolopsis qantasi</i> Russell & Gloerfelt-Tarp 1984	Slender dwarf monocle bream, (M, En), Ref. 3810. Max. 10.3 cm SL. Museum: NTM S.10996-001 (TGT2323). From southwest Sumatra to Bali Strait (Ref. 5978).	<i>Scolopsis taeniopterus</i> (Kuhl & Van Hasselt 1830)	Latticed Ma (TG
<i>Parascolopsis tanyactis</i> Russell 1986	Long-rayed dwarf monocle bream, (M, Fi), Ref. 3810. Max. 21 cm SL. Museum: NTM S.10739-002 (TGT1782), AMS I.24278-001 (Ref. 9923). From southwest Sumatra to Timor (Ref. 5978).	<i>Scolopsis temporalis</i> (Cuvier 1830)	Bald-s cm Three
<i>Parascolopsis tosensis</i> (Kamohara 1938)	Tosa dwarf monocle bream, (M, Fi), Ref. 3810. Max. 10 cm SL. Museum: NTM S.10760-003, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Scolopsis trilineatus</i> Kner 1868	381 from
<i>Pentapodus bifasciatus</i> (Bleeker 1848)	White-shouldered whiptail, (M, Fi), Ref. 3810. Max. 15 cm SL.	<i>Scolopsis vosmeri</i> (Bloch 1792)	White 381 S.1 Sea
<i>Pentapodus caninus</i> (Cuvier 1830)	Small-toothed whiptail, Gurisi, (M, Fi), Ref. 3810. Max. 18.5 cm SL. Also Ref.: 8631. Museum: NTM S.10660-001, from Bali Strait to Timor Sea (Ref. 5978). Also known from Flores.	<i>Scolopsis xenochrous</i> Günther 1872	Obliqu Ma and
<i>Pentapodus emeryii</i> (Richardson 1843)	Double whiptail, (M, Fi), Ref. 3810. Max. 24.5 cm SL. Also Ref.: 8631. Museum: NTMS.10731-004, from Bali Strait to Timor Sea (Ref. 5978).	Nomeidae <i>Cubiceps kotlyari</i> Agafonova 1988	Driftfi (M), F
<i>Pentapodus nagasakiensis</i> (Tanaka 1915)	Japanese whiptail, (M, Fi), Ref. 3810. Max. 15 cm SL. Museum: AMS I.22831-033, from Bali Strait to Timor Sea (Ref. 5978).	<i>Cubiceps pauciradiatus</i> Günther 1872	Bigey Ma sou
<i>Pentapodus porosus</i> (Valenciennes 1830)	Northwest Australian whiptail, (M, Fi), Ref. 3810. Max. 23 cm SL.	<i>Psenes arafurensis</i> Günther 1889	Bande Mu sou
<i>Pentapodus setosus</i> (Valenciennes 1830)	Butterfly whiptail, Krisi, (M, Fi), Ref. 3810. Max. 17.5 cm SL. Museum: NTM S.10749-004, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Psenes pellucidus</i> Lütken 1880	Bluefi Ma (PJ Stra
<i>Pentapodus trivittatus</i> (Bloch 1791)	Three-striped whiptail, Krisi, (M, Fi), Ref. 3810. Max. 28 cm TL. Also Ref.: 8631. Museum: NTM S.10733-020, from Bali Strait to Timor Sea (Ref. 5978).	<i>Psenes whiteleggii</i> Waite 1894	Shado Mu Fro 441
<i>Scolopsis affinis</i> Peters 1877	Peters' monocle bream, (M, Fi), Ref. 3810. Max. 20 cm SL. Also Ref.: 8631. Museum: NTM S.10763-002, from Bali Strait to Timor Sea (Ref. 5978).	Opistognathidae <i>Opistognathus castelnaui</i> Bleeker 1871	Jawfi (M), F S.1 Sea
<i>Scolopsis auratus</i> (Park 1797)	Yellowstripe monocle bream, (M, Fi), Ref. 3810. Max. 21 cm SL. Museum: NTM S.10771-001, from southwest Sumatra to Bali Strait (Ref. 5978).	Pempheridae <i>Parapriacanthus ransonneti</i> Steindachner 1870	Swee Pigmy Ma S.1 to T
<i>Scolopsis bilineatus</i> (Bloch 1793)	Two-lined monocle bream, Ija puti, (M, Fi), Ref. 3810. Max. 20 cm SL. Also Ref.: 8631. Museum: NTM S.10732-001, from Bali Strait to Timor Sea (Ref. 5978). Also known from Flores.	<i>Pempheris moluca</i> Cuvier 1831	(M), F sou
<i>Scolopsis ciliatus</i> (Lacepède 1802)	Saw-jawed monocle bream, Ija putilo ote, (M, Fi), Ref. 3810. Max. 13.5 cm SL. Also Ref.: 8631. Museum: NTM S.10733-005, from Bali Strait to Timor Sea (Ref. 5978).	<i>Pempheris oualensis</i> Cuvier 1831	Silver Kne
<i>Scolopsis lineatus</i> Quoy & Gaimard 1824	Striped monocle bream, Pasir-pasir, (M, Fi), Ref. 3810. Max. 20 cm SL. Museum: NTM S.11002-001, from Bali Strait to Timor Sea (Ref. 5978).		

Table 2. Marine and brackishwater fishes of Indonesia.

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<p><i>Pempheris schwenkii</i> Bleeker 1855 <i>Pempheris vanicolensis</i> Cuvier 1831</p>	<p>Schwenk's sweeper, (M), Ref. 2334. Max. 15 cm TL. Vanikoro sweeper, (M), Ref. 5978 Max. 20 cm TL. Museum: NTM S.10732-005 (PAN5). From Southwest Sumatra to Timor Sea.</p>	<p>Polynemidae <i>Eleutheronema tetradactylum</i> (Shaw 1804) <i>Eleutheronema tridactylum</i> (Bleeker 1849) <i>Filimanus heptadactyla</i> (Cuvier 1829) <i>Filimanus hexanema</i> (Cuvier 1829)</p>	<p>Threacidae Fourfin Threafin Sevenfin Javane Jaka Stra Splend Pad Eightfin Yellowf 198 Sum Indian Small-f Also Fro Borr Seven Black-f TL Striped Mus Fro Sixting Blacks Max Stra Dwarf Kno of a traw Feb Angeli Griffis Pho veril Threes Alsc (TG Golder Acc occl trad Bicolor Alsc Twosp FL.</p>
<p>Pentacerotidae <i>Histiopertus typus</i> Temminck & Schlegel 1844</p>	<p>Armorheads Sailfin armourhead, (M, Fi), Ref. 5978. Max. 35 cm SL. Museum: CSIRO CA2257. From southwest Sumatra to Bali Strait.</p>	<p><i>Filimanus perplexa</i> Feltes 1991 <i>Filimanus sealei</i> (Jordan & Richardson 1910) <i>Filimanus xanthonema</i> (Valenciennes 1831)</p>	
<p>Percophidae <i>Bembrops curvatura</i> Okada & Suzuki 1952</p>	<p>Duckbills (M), Ref. 5978. Max. 16 cm SL. Museum: NTM S.10740-004 (TGT1630). From Bali Strait to Timor Sea.</p>	<p><i>Polydactylus indicus</i> (Shaw 1804) <i>Polydactylus microstoma</i> (Bleeker 1851)</p>	
<p><i>Bembrops filodorsalia</i> Okada & Suzuki 1952</p>	<p>Sharpnosed duckbill, (M), Ref. 5978. Max. 20 cm TL. Museum: AMS I.22808-001. From Bali Strait to Timor Sea. In range Ref.: 3132.</p>	<p><i>Polydactylus multiradiatus</i> (Günther 1860) <i>Polydactylus nigripinnis</i> Munro 1964</p>	
<p><i>Bembrops philippinus</i> Fowler 1939 <i>Chironema chlorotaenia</i> McKay 1971</p>	<p>(M), Ref. 5978. Museum: NTM S.10998-002 (TGT2512). From Bali Strait to Timor Sea. (M), Ref. 5978 Max. 22 cm SL. Museum: NTM S.10760- 009 (TGT1727). From Bali Strait to Timor Sea.</p>	<p><i>Polydactylus sextarius</i> (Bloch & Schneider 1801)</p>	
<p>Pholidichthyidae <i>Pholidichthys leucotaenia</i> Bleeker 1856</p>	<p>Convict blenny Convict blenny, (M), Ref. 8631. Max. 15 cm TL. Also Ref.: 6201. Known from Flores. Museum: BPBM 18050.</p>	<p><i>Polynemus verekeri</i> (Saville-Kent 1889)</p>	
<p>Pinguipedidae <i>Parapercis alboguttata</i> (Günther 1872)</p>	<p>Sandperches Blue-nosed grubfish, (M), Ref. 3132. Max. 22 cm TL.</p>	<p><i>Centropyge aurantius</i> (Lacepède 1831)</p>	
<p><i>Parapercis clathrata</i> Ogilby 1910</p>	<p>Latticed sandperch, (M), Ref. 8631. Max. 15.3 cm SL. Known from Flores.</p>	<p><i>Centropyge bicolor</i> (Bloch 1787) <i>Centropyge bispinosus</i> (Günther 1860)</p>	
<p><i>Parapercis cylindrica</i> (Bloch 1792)</p>	<p>Cylindrical sandperch, (M), Ref. 2334. Max. 13 cm SL.</p>	<p><i>Centropyge trimaculatus</i> (Lacepède 1831)</p>	
<p><i>Parapercis hexophtalma</i> (Cuvier 1829)</p>	<p>Spotted sandmelt, (M), Ref. 8631. Max. 23 cm TL. Known from Flores.</p>	<p><i>Centropyge tetracantha</i> (Lacepède 1802)</p>	
<p><i>Parapercis millepunctata</i> (Günther 1860)</p>	<p>Blackdotted sand perch, (M), Ref. 8631. Max. 17 cm SL. Known from Bali.</p>	<p><i>Centropyge xanthozona</i> (Bleeker 1849)</p>	
<p><i>Parapercis mimaseana</i> (Kamohara 1937)</p>	<p>(M), Ref. 5978. Max. 20 cm SL. Museum: CSIRO CA1441. From southwest Sumatra to Bali Strait.</p>	<p><i>Calloptlesiops altivelis</i> (Steindachner 1903)</p>	
<p><i>Parapercis multiplicata</i> Randall 1984</p>	<p>Double-stitch grubfish, (M), Ref. 2334. Max. 12 cm TL.</p>	<p><i>Calloptlesiops somaliensis</i> Schultz 1968</p>	
<p><i>Parapercis sexfasciata</i> (Temminck & Schlegel 1843)</p>	<p>(M), Ref. 5978. Max. 12 cm TL. Museum: BMNH 1984.1.1.88 (TGT (PJPW) 560). From southwest Sumatra to Bali Strait.</p>	<p><i>Parapercis snyderi</i> Jordan & Starks 1905</p>	
<p><i>Parapercis snyderi</i> Jordan & Starks 1905</p>	<p>U-mark sandperch, (M), Ref. 2334. Max. 10 cm TL.</p>	<p><i>Parapercis somaliensis</i> Schultz 1968</p>	
<p><i>Parapercis tetraacantha</i> (Lacepède 1802)</p>	<p>Weeping sandmelt, (M), Ref. 5978. Max. 20 cm TL. Museum: NTM S.10746-004 (TGT1342). From Bali Strait to Timor Sea. This is a new record for southern Indonesia.</p>	<p><i>Parapercis xanthozona</i> (Bleeker 1849)</p>	
<p><i>Parapercis xanthozona</i> (Bleeker 1849)</p>	<p>White-blotched sandperch, (M), Ref. 8631. Max. 22 cm SL. Known from Flores.</p>	<p><i>Calloptlesiops altivelis</i> (Steindachner 1903)</p>	
<p>Plesiopidae <i>Calloptlesiops altivelis</i> (Steindachner 1903)</p>	<p>Yellowbar sandperch, Peppered grubfish, (M), Ref. 2334 Max. 23 cm TL. From Bali (Ref. 8631).</p>	<p><i>Calloptlesiops altivelis</i> (Steindachner 1903)</p>	
<p></p>	<p>Roundheads Comet, (M), Ref. 2334. Max. 16 cm TL. Flores (Ref. 8631).</p>	<p></p>	

Table 2. Continuation.
 [Tabel 2. Sambungan.]

<i>Centropyge colini</i> Smith-Vaniz & Randall 1974	Cocos-Keeling angelfish, (M), Ref. Max. 7 cm SL. Museum: BPBM 34195.	<i>Abudefduf septemfasciatus</i> (Cuvier 1830)	Ban-
<i>Centropyge eibli</i> Klausewitz 1963	Blacktail angelfish, (M), Ref. 4859. Max. 15 cm. Also Ref.: 8631. Known from Bali.	<i>Abudefduf sexfasciatus</i> (Lacepède 1801)	Scis
<i>Centropyge flavicauda</i> Fraser-Brunner 1933	Whitetail angelfish, (M), Ref. 4537. Max. 5.7 cm SL. Also Ref.: 8631. Known from Flores.	<i>Abudefduf sordidus</i> (Forsskål 1775)	Blac
<i>Centropyge nox</i> (Bleeker 1853)	Midnight angelfish, (M), Ref. 4859. Max. 9 cm TL.	<i>Abudefduf vaigiensis</i> (Quoy & Gaimard 1825)	Indo
<i>Centropyge tibicen</i> (Cuvier 1831)	Keyhole angelfish, (M), Ref. 4537. Max. 14.9 cm SL.	<i>Acanthochromis polyacanthus</i> (Bleeker 1855)	Spin R
<i>Centropyge vroliki</i> (Bleeker 1853)	Pearlscale angelfish, (M, Fi), Ref. 1602. Max. 7.5 cm SL. Also Ref.: 8631. Known from Flores.	<i>Amblyglyphidodon aureus</i> (Cuvier 1830)	pe Golc
<i>Chaetodontoplus chrysocephalus</i> Bleeker 1854	Orangeface angelfish, Orange-faced angelfish, (M), Ref. 4858. Max. 22 cm.	<i>Amblyglyphidodon curacao</i> (Bloch 1787)	AI Stag
<i>Chaetodontoplus duboulayi</i> (Günther 1867)	Scribbled angelfish, (M), Ref. 2334. Max. 28 cm TL.	<i>Amblyglyphidodon leucogaster</i> (Bleeker 1847)	Yellc KI
<i>Chaetodontoplus melanosoma</i> (Bleeker 1853)	Black-velvet angelfish, (M), Ref. 5978. Max. 18 cm TL. Also Ref.: 559, 8631. Museum: NTM S.10744-002 (TGT951). From Bali Strait to Timor Sea.	<i>Amblyglyphidodon ternatensis</i> (Bleeker 1853)	Terri. AI
<i>Chaetodontoplus mesoleucus</i> (Bloch 1787)	Vermiculated angelfish, (M), Ref. 1602. Max. 15.1 cm SL.	<i>Amblypomacentrus breviceps</i> (Schlegel & Müller 1839)	Blac SI
<i>Genicanthus lamarck</i> (Lacepède 1802)	Blackstriped angelfish, (M), Ref. 2334. Max. 22 cm TL. Also Ref.: 8631. Known from Flores.	<i>Amphiprion akallopisos</i> Bleeker 1853	Skur AI
<i>Genicanthus melanospilus</i> (Bleeker 1857)	Spotbreast angelfish, (M), Ref. 1602. Max. 14.9 cm SL. Also Ref.: 8631. Known from Flores.	<i>Amphiprion chrysopterus</i> Cuvier 1830	Orar SI
<i>Paracentropyge multifasciatus</i> (Smith & Radcliffe 1911)	Barred angelfish, (M), Ref. 4859. Max. 9.3 cm SL. Also Ref.: 8631.	<i>Amphiprion clarkii</i> (Bennett 1830)	Yello AI
<i>Pomacanthus annularis</i> (Bloch 1787)	Bluering angelfish, (M), Ref. 5978. Max. 35 cm TL. Also Ref.: 8631. From southwest Sumatra to Timor Sea.	<i>Amphiprion ephippium</i> (Bloch 1790)	Sadc AI
<i>Pomacanthus imperator</i> (Bloch 1787)	Emperor angelfish, (M), Ref. 5978. Max. 40 cm. Also Ref.: 8631. Museum: LPPL JIF97 (TGT3275). From Bali Strait to Timor Sea.	<i>Amphiprion frenatus</i> Brevoort 1856	Tomr AI
<i>Pomacanthus navarchus</i> (Cuvier 1831)	Bluegirdled angelfish, (M), Ref. 1602. Max. 30 cm TL. Also Ref.: 8631. Known from Flores.	<i>Amphiprion melanopus</i> Bleeker 1852	Fire AI
<i>Pomacanthus semicirculatus</i> (Cuvier 1831)	Semicircle angelfish, (M), Ref. 4859. Max. 29 cm SL. Also Ref.: 8631. Museum: LPPL JIF98, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Amphiprion ocellaris</i> Cuvier 1830	Clow AI
<i>Pomacanthus sexstriatus</i> (Cuvier 1831)	Sixbar angelfish, (M), Ref. 4859. Max. 46 cm TL. Museum: LPPL JIF99 (TGT3242). From Bali Strait to Timor Sea (Ref. 5978).	<i>Amphiprion perideraion</i> Bleeker 1855	Pink
<i>Pomacanthus xanthometopon</i> (Bleeker 1853)	Yellowface angelfish, (M, Fi), Ref. 8631. Max. 32 cm SL. Known from Bali.	<i>Amphiprion polymnus</i> (Linnaeus 1758)	Sadc
<i>Pygoplites diacanthus</i> (Boddaert 1772)	Regal angelfish, (M), Ref. 8631. Max. 20.9 cm SL. Known from Bali and Flores.	<i>Amphiprion sandaracinos</i> Allen 1972	Yello Kr
Pomacentridae	Damselfishes	<i>Amphiprion sebae</i> Bleeker 1853	Brow
<i>Abudefduf bengalensis</i> (Bloch 1787)	Bengal sergeant, (M, Br), Ref. 4966. Max. 14 cm SL. Also Ref.: 8631. Known from Jakarta.	<i>Cheiloprion labiatus</i> (Day 1877)	Big-li AI
<i>Abudefduf lorentzi</i> Hensley & Allen 1977	Black-tail sergeant, (M), Ref. 1602. Max. 15 cm SL. Also Ref.: 8631. Known from Flores.	<i>Chromis alpha</i> Randall 1988	Yello Kr
<i>Abudefduf notatus</i> (Day 1870)	Yellowtail sergeant, (M), Ref. 7247. Max. 14 cm SL.	<i>Chromis amboinensis</i> (Bleeker 1873)	Ambc Kr
<i>Abudefduf saxatilis</i> (Linnaeus 1758)	Sergeant major, (M, Fi), Ref. 5978. Max. 21.5 cm TL. Museum: LPPL JIF110 (TGT1576). From Bali Strait to Timor Sea.	<i>Chromis analis</i> (Cuvier 1830)	Yello AI
		<i>Chromis atripectoralis</i> Welander & Schultz 1951	Black AI
		<i>Chromis atripes</i> Fowler & Bean 1928	Dark Kr

Table 2. Marine and brackishwater fishes of Indonesia .

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Chromis caudalis</i> Randall 1988	Blue-axil chromis, (M), Ref. 1602. Ref.: 8631. Known from Flores. Max. 7.5 cm SL. Also in	<i>Chrysiptera oxycephala</i> (Bleeker 1877)	Blue-
<i>Chromis cinerascens</i> (Cuvier 1830)	Green chromis, (M), Ref. 7247. Max. 10 cm SL.	<i>Chrysiptera parasema</i> (Fowler 1918)	Goldt
<i>Chromis delta</i> Randall 1988	Deep-reef chromis, (M), Ref. 1602. Max. 5 cm SL. Also Ref.: 8631. Known from Flores.	<i>Chrysiptera rex</i> (Snyder 1909)	Kn
<i>Chromis elerae</i> Fowler & Bean 1928	Twinspot chromis, (M), Ref. 8631. Max. 5.5 cm SL. Known from Flores.	<i>Chrysiptera rollandi</i> (Whitley 1961)	King
<i>Chromis flavipectoralis</i> Randall 1989	Malayan chromis, (M), Ref. 7247. Max. 5.5 cm SL. Museum: Java Sea, Pulau Seribu, Pulau Putri, BPBM 18569 (Holotype) (Ref.10591).	<i>Chrysiptera springeri</i> (Allen & Lubbock 1976)	Alis
<i>Chromis fumea</i> (Tanaka 1917)	Smokey chromis, (M), Ref. 3132. Max. 10 cm SL. Known from Komodo I. (Allen pers. comm.).	<i>Chrysiptera talboti</i> (Allen 1975)	an
<i>Chromis lepidolepis</i> Bleeker 1877	Scaly chromis, (M), Ref. 8631. Max. 6.5 cm SL. Known from Flores.	<i>Chrysiptera unimaculata</i> (Cuvier 1830)	Talbo
<i>Chromis lineata</i> Fowler & Bean 1928	Lined chromis, (M), Ref. 7247. Max. 4 cm SL. Also Ref.: 8631. Known from Flores.	<i>Dascyllus aruanus</i> (Linnaeus 1758)	Alis
<i>Chromis margaritifer</i> Fowler 1946	Bicolor chromis, (M), Ref. 8631. Max. 6.5 cm SL. Known from Flores.	<i>Dascyllus carneus</i> Fischer 1885	White
<i>Chromis nigroanalis</i> Randall 1989	Kenyan chromis, (M), Ref. 10591. Max. 9 cm SL. Ranges east to the western Java Sea. Museum: Pulau Seribu, Pulau Pari Grp, Pulau Tikus, USNM 270852.	<i>Dascyllus melanurus</i> Bleeker 1854	Alis
<i>Chromis retrofasciata</i> Weber 1913	Black-bar chromis, (M), Ref. 1602. Max. 4 cm SL. Also Ref.: 8631. Known from Flores.	<i>Dascyllus reticulatus</i> (Richardson 1846)	Cloud
<i>Chromis scotochiloptera</i> Fowler 1918	Philippines chromis, (M), Ref. 7247. Max. 12 cm SL. Also Ref.: 8631, 9137. Known from Flores.	<i>Dascyllus trimaculatus</i> (Rüppell 1828)	Jav
<i>Chromis ternatensis</i> (Bleeker 1856)	Ternate chromis, (M), Ref. 1602. Max. 7.4 cm SL. Known from the Manado area, Sulawesi (Celebes), Maumere Bay, Flores, and Komodo I. (Allen, pers. comm.).	<i>Dischistodus chrysopoecilus</i> (Schlegel & Müller 1839)	Black
<i>Chromis viridis</i> (Cuvier 1830)	Blue-green damselfish, (M), Ref. 8631. Max. 7 cm SL. Known from Flores.	<i>Dischistodus fasciatus</i> (Cuvier 1830)	Alis
<i>Chromis weberi</i> Fowler & Bean 1928	Weber's chromis, (M), Ref. 8631. Max. 9.5 cm SL. Known from Flores.	<i>Dischistodus melanotus</i> (Bleeker 1858)	Black
<i>Chromis xanthochira</i> (Bleeker 1851)	Yellow-axil chromis, (M), Ref. 1602. Max. 10 cm SL. Also Ref.: 8631. Known from Flores.	<i>Dischistodus perspicillatus</i> (Cuvier 1830)	White
<i>Chromis xanthura</i> (Bleeker 1854)	Paletail chromis, (M), Ref. 8631. Max. 12 cm SL. Known from Flores.	<i>Dischistodus prosopotaenia</i> (Bleeker 1852)	from
<i>Chrysiptera biocellata</i> (Quoy & Gaimard 1825)	Twinspot damselfish, (M), Ref. 8631. Max. 8 cm SL. Known from Flores.	<i>Dischistodus pseudochrysopoecilus</i> (Allen & Robertson 1974)	Hone
<i>Chrysiptera bleekeri</i> (Fowler & Bean 1928)	Bleeker's damsel, (M), Ref. 7247. Max. 6.5 cm SL. Also Ref.: 8631. Known from Timor and Flores.	<i>Hemiglyphidodon plagiometopon</i> (Bleeker 1852)	Alis
<i>Chrysiptera caeruleolineata</i> (Allen 1973)	Blueline demoiselle, (M), Ref. 8631. Max. 4 cm SL. Known from Flores.	<i>Lepidozygus tapeinosoma</i> (Bleeker 1856)	Mon
<i>Chrysiptera cyanea</i> (Quoy & Gaimard 1825)	Sapphire devil, (M), Ref. 7247. Max. 6 cm SL.	<i>Neoglyphidodon bonang</i> (Bleeker 1852)	from
<i>Chrysiptera glauca</i> (Cuvier 1830)	Grey demoiselle, (M), Ref. 1602. Max. 8 cm SL. Known from the Manado area, Sulawesi (Celebes) and Maumere Bay, Flores (Allen, pers. comm.).	<i>Neoglyphidodon crossi</i> Allen 1991	Lago
<i>Chrysiptera hemicyanea</i> (Weber 1913)	Azure demoiselle, (M), Ref. 4966. Max. 5 cm SL.	<i>Neoglyphidodon melas</i> (Cuvier 1830)	Re
<i>Chrysiptera leucopoma</i> (Lesson 1830)	Surge damselfish, (M), Ref. 8631. Max. 6 cm SL. Known from Flores.	<i>Neoglyphidodon nigroris</i> (Cuvier 1830)	Fusili
		<i>Neoglyphidodon oxyodon</i> (Bleeker 1858)	Kn
			Ko
			cor
			Ocell
			Cross
			Bowti
			Kn
			Black
			Alis
			Blues
			Alis

Table 2. Continuation.
[Tabel 2. Sambungan.]

	<i>Neoglyphidodon thoracotaeniatus</i> (Fowler & Bean 1928)	Barhead damsel, Bar-cheek damsel, (M), Ref. 4966 Max. 8.5 cm SL. From Flores (Ref. 8631).	<i>Pomacentrus cuneatus</i> Allen 1991	Wed
	<i>Neopomacentrus anabatoides</i> (Bleeker 1847)	Silver demoiselle, Brown demoiselle, (M), Ref. 4966 Max. 8 cm SL.	<i>Pomacentrus emarginatus</i> Cuvier 1830	Oute
	<i>Neopomacentrus azysron</i> (Bleeker 1877)	Yellowtail demoiselle, Yellow-tail demoiselle (M), Ref. 4966 Max. 6 cm SL. Also Ref.: 8631. Known from Flores.	<i>Pomacentrus grammorhynchus</i> Fowler 1918	Blue
	<i>Neopomacentrus cyanomos</i> (Bleeker 1856)	Regal demoiselle, (M), Ref. 8631. Max. 7 cm SL. Known from Bali.	<i>Pomacentrus javanicus</i> Allen 1991	Java
	<i>Neopomacentrus filamentosus</i> (Macleay 1882)	Brown demoiselle, (M), Ref. 7247. Max. 6 cm SL.	<i>Pomacentrus lepidogenys</i> Fowler & Bean 1928	Scal Ri
	<i>Neopomacentrus nemurus</i> (Bleeker 1857)	Coral demoiselle, (M), Ref. 7247. Max. 5.5 cm SL. Also Ref.: 8631. Known from Flores.	<i>Pomacentrus littoralis</i> Cuvier 1830	Smo
	<i>Neopomacentrus taeniurus</i> (Bleeker 1856)	Freshwater demoiselle, (M, Br, Fr), Ref. 4966. Max. 8.5 cm SL.	<i>Pomacentrus melanochir</i> Bleeker 1877	Indo Al
	<i>Neopomacentrus violascens</i> (Bleeker 1848)	Violet demoiselle, (M), Ref. 1602. Max. 5 cm SL. Also Ref.: 8631. Known from Bali.	<i>Pomacentrus moluccensis</i> Bleeker 1853	Lem Re
	<i>Plectroglyphidodon dickii</i> (Liénard 1839)	Blackbar devil, (M), Ref. 7247. Max. 8.5 cm SL. Known from the Manado area, Sulawesi (Celebes), Komodo I., and Maumere Bay, Flores (Allen, pers. comm.).	<i>Pomacentrus nagasakiensis</i> Tanaka 1917	Naga fr
	<i>Plectroglyphidodon johnstonianus</i> Fowler & Ball 1924	Johnston Island damsel, (M), Ref. 8631. Max. 7 cm SL. Known from Flores.	<i>Pomacentrus nigromanus</i> Weber 1913	Gold
	<i>Plectroglyphidodon lacrymatus</i> (Quoy & Gaimard 1825)	Whitespotted devil, (M), Ref. 7247. Max. 8.5 cm SL. Known from the Manado area, Sulawesi (Celebes), Komodo I., and Maumere Bay, Flores (Allen, pers. comm.).	<i>Pomacentrus nigromarginatus</i> Allen 1973	Blac Al
261	<i>Plectroglyphidodon leucozona</i> (Bleeker 1859)	Singlebar devil, (M), Ref. 7247. Max. 9 cm SL. Known from the Manado area, Sulawesi (Celebes), Komodo I., and Maumere Bay, Flores (Allen, pers. comm.).	<i>Pomacentrus pavo</i> (Bloch 1787)	Sapp Kr
	<i>Plectroglyphidodon phoenixensis</i> (Schultz 1943)	Phoenix devil, (M), Ref. 7247. Max. 7 cm SL. Known from Komodo I. (Allen pers. comm.).	<i>Pomacentrus philippinus</i> Evermann & Seale 1907	Philip Kr
	<i>Pomacentrus adelus</i> Allen 1991	Obscure damsel, (M), Ref. 7247. Max. 6.5 cm SL.	<i>Pomacentrus reidi</i> Fowler & Bean 1928	Reid Re
	<i>Pomacentrus alexanderae</i> Evermann & Seale 1907	Alexander's damsel, (M), Ref. 4966. Max. 7 cm SL. Also Ref.: 8631. Known from Flores.	<i>Pomacentrus simsiang</i> Bleeker 1856	Blue Re
	<i>Pomacentrus amboinensis</i> Bleeker 1868	Ambon damsel, (M), Ref. 1602. Max. 10.5 cm FL. Also Ref.: 8631. Known from Flores.	<i>Pomacentrus smithi</i> Fowler & Bean 1928	Smith Re
	<i>Pomacentrus auriventris</i> Allen 1991	Goldbelly damsel, (M), Ref. 7247. Max. 5.5 cm SL. Known from the Manado area, Sulawesi (Celebes), Komodo I., and Maumere Bay, Flores (Allen, pers. comm.).	<i>Pomacentrus taeniometopon</i> Bleeker 1852	Brac Kr an
	<i>Pomacentrus azuremaculatus</i> Allen 1991	Bluespotted damsel, (M), Ref. 7247. Max. 8 cm SL.	<i>Pomacentrus tripunctatus</i> Cuvier 1830	Thre Kr
	<i>Pomacentrus bankanensis</i> Bleeker 1853	Speckled damselfish, (M), Ref. 7247. Max. 7 cm SL. Also Ref.: 8631. Known from Flores.	<i>Pomacentrus vaiuli</i> Jordan & Seale 1906	Ocel Ma Fl
	<i>Pomacentrus brachialis</i> Cuvier 1830	Charcoal damsel, (M), Ref. 1602. Max. 8 cm SL.	<i>Pomacentrus xanthosternus</i> Allen 1991	Yell SL
	<i>Pomacentrus burroughi</i> Fowler 1918	Burrough's damsel, (M), Ref. 1602. Max. 6.5 cm SL. Known from the Manado area, Sulawesi (Celebes), Komodo I., and Maumere Bay, Flores (Allen, pers. comm.).	<i>Premnas biaculeatus</i> (Bloch 1790)	Spin SL
	<i>Pomacentrus chrysurus</i> Cuvier 1830	Whitetail damsel, (M), Ref. 8631. Max. 7 cm SL. Known from Bali and Flores.	<i>Pristotis jerdoni</i> (Day 1873)	Gulf Mu Str
	<i>Pomacentrus coelestis</i> Jordan & Starks 1901	Neon damselfish, (M), Ref. 7247. Max. 7 cm SL. Also Ref.: 8631. Known from Flores.	<i>Stegastes albifasciatus</i> (Schlegel & Müller 1839)	White Al
			<i>Stegastes fasciolatus</i> (Ogilby 1889)	Pacif
			<i>Stegastes lividus</i> (Bloch & Schneider 1801)	Blunt
			<i>Stegastes nigricans</i> (Lacepède 1802)	Dusk Al

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Stegastes obreptus</i> (Whitley 1948)	Western gregory, (M), Ref. 510. Max. 12 cm SL. Molucca Islands and Seribu Islands, off Java.	<i>Pseudochromis polynemus</i> Fowler 1931	Longf Kn
Priacanthidae	Bigeyes or catalufas	<i>Pseudochromis porphyreus</i> Lubbock & Goldman 1974	Mage
<i>Cookeolus japonicus</i> (Cuvier 1829)	Longfinned bullseye, (M), Ref. 5978 Max. 60 cm TL. Museum: USNM 263762 (TGT1414). From southwest Sumatra to Bali Strait.	<i>Pseudochromis quinquentatus</i> McCulloch 1926	Spiny 00
<i>Heteropriacanthus cruentatus</i> (Lacepède 1801)	Glasseye, Red big-eye, (M, Fi), Ref. 5978 Max. 34 cm TL. Museum: NTM S.10732-007 (PAN20). From Bali Strait to Timor Sea. Also Ref. 8631. In range Ref.: 2334.	<i>Pseudochromis splendens</i> Fowler 1931	Splen fro
<i>Priacanthus blochii</i> Bleeker 1853	Silver big-eye, (M), Ref. 5403. Max. 23.5 cm SL. Museum: USNM 263758 (TGT2531). Found from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Pseudochromis steenei</i> Gill & Randall 1992	Lyrete on or r
<i>Priacanthus fitchi</i> Sarnes 1988	(M), Ref. 5403. Max. 18.5 cm SL. Museum: USNM 263760 (TGT2454). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Pseudochromis tapeinosoma</i> Bleeker 1853	(M), F
<i>Priacanthus hamrur</i> (Forsskål 1775)	Moontail bullseye, (M, Fi), Ref. 5978. Max. 45 cm. Museum: USNM 263761 (TGT2414). From southwest Sumatra to Timor Sea.	<i>Pseudoplesiops annae</i> (Weber 1913)	(M), F
<i>Priacanthus macracanthus</i> Cuvier 1829	Red bigeye, (M, Fi), Ref. 5978. Max. 29 cm SL. Museum: USNM 263759 (TGT2557). From southwest Sumatra to Timor Sea.	<i>Pseudoplesiops knighti</i> Allen 1987	(M), F
<i>Priacanthus sagittarius</i> Sarnes 1988	(M, Fi), Ref. 5403. Max. 29 cm SL. From southwest Sumatra to Timor Sea (Ref. 5978). Museum: Sumatra, USNM 285042 (Holotype); USNM 263757 (TGT2291).	<i>Pseudoplesiops multisquamatus</i> Allen 1987	Fine-s Ma
262 <i>Priacanthus tayenus</i> Richardson 1846	Purple-spotted bigeye, (M, Fi), Ref. 3414. Max. 25 cm Museum: LPPL JIF207 (TGT2587). SL. Found from southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 6567.	<i>Pseudoplesiops rosae</i> Schultz 1943	(M), F
<i>Pristigenys meyeri</i> (Günther 1872)	(M), Ref. 5403. Max. 22.5 cm SL. Sulawesi and New Guinea.	<i>Pseudoplesiops typus</i> Bleeker 1858	Hidde
<i>Pristigenys niphonica</i> (Cuvier 1829)	Japanese bigeye, (M), Ref. 5978. Max. 27.4 cm SL. Museum: CSIRO B.2174. From southwest Sumatra to Bali Strait.	Rachycentridae	Cobia
Pseudochromidae	Dottybacks	<i>Rachycentron canadum</i> (Linnaeus 1766)	Cobia Alsa Bal 109
<i>Congrogadus malayanus</i> (Weber 1909)	(M), Ref. 531. Max. 7.1 cm SL. Known from Aru Islands, Arafura Sea.	Scaridae	Parro
<i>Congrogadus subducens</i> (Richardson 1843)	Carpet eel-blenny, (M), Ref. 531. Max. 45 cm TL.	<i>Bolboretomon muricatum</i> (Valenciennes 1840)	Green cm
<i>Haliophis aethiopus</i> Winterbottom 1985	(M), Ref. 531. Max. 5 cm SL. Museum: Bali, Sanur, off Alit's Beach Bungalows, BPBM 20920 (Holotype); ROM 38419 (Paratype).	<i>Calotomus carolinus</i> (Valenciennes 1840)	Caroli in A
<i>Labracinus cyclophthalmus</i> (Müller & Troschel 1849)	Fire-tail devil, (M), Ref. 8631. Max. 20 cm TL. From Flores.	<i>Calotomus spinidens</i> (Quoy & Gaimard 1824)	Spiny Mu: See
<i>Pseudochromis bitaeniatus</i> (Fowler 1931)	Double-striped dottyback, (M), Ref. 2334. Max. 7 cm TL. Flores (Ref. 8631).	<i>Hipposcarus longiceps</i> (Valenciennes 1840)	Pacific SL.
<i>Pseudochromis cyanotaenia</i> Bleeker 1857	Surge dottyback, (M), Ref. 1602. Max. 4.5 cm SL.	<i>Leptoscarus vaigiensis</i> (Quoy & Gaimard 1824)	Marble Mu: (Re
<i>Pseudochromis fuscus</i> Müller & Troschel 1849	Brown dottyback, (M), Ref. 8631. Max. 9 cm. In range Ref.: 2334. Known from Bali and Flores.	<i>Scarus atropectoralis</i> Schultz 1958	Red p Knc
<i>Pseudochromis marshallensis</i> (Schultz 1953)	Marshall Is. dottyback, (M), Ref. 8631. Max. 6.5 cm SL. Known from Flores.	<i>Scarus bleekeri</i> (De Beaufort 1940)	Bleek Ref
<i>Pseudochromis paccagnellae</i> Axelrod 1973	Royal dottyback, (M), Ref. 2334. Max. 7 cm TL. Flores (Ref. 8631).	<i>Scarus bowersi</i> (Snyder 1909)	Bower
		<i>Scarus dimidiatus</i> Bleeker 1859	Yellow
		<i>Scarus festivus</i> Valenciennes 1840	Festive
		<i>Scarus flavipectoralis</i> Schultz 1958	Yellow Knc
		<i>Scarus forsteni</i> (Bleeker 1861)	Forste

Table 2. Continuation.
 [Tabel 2. Sambungan.]

263	<i>Scarus frenatus</i> Lacepède 1802	Bridled parrotfish, (M), Ref. 1602. Max. 47 cm TL.	<i>Aspericorvina jubata</i> (Bleeker 1855)	Pric...
	<i>Scarus ghobban</i> Forsskål 1775	Blue-barred parrotfish, (M, Br, Fi), Ref. 5978. Max. 90 cm. Also Ref.: 8631. From southwest Sumatra to Timor Sea.	<i>Atrobucca kyushini</i> Sasaki & Kailola 1988	Blac...
	<i>Scarus gibbus</i> Rüppell 1828	Heavybeak parrotfish, (M), Ref. 2689. Max. 50 cm SL.	<i>Atrobucca nibe</i> (Jordan & Thompson 1911)	Blac...
	<i>Scarus globiceps</i> Valenciennes 1840	Globehead parrotfish, (M), Ref. 1602. Max. 26 cm SL.	<i>Austronibeia oedogenys</i> Trewavas 1977	O...
	<i>Scarus hypselopterus</i> (Bleeker 1853)	Yellow-tail parrotfish, (M), Ref. 8631. Max. 26 cm SL. Also Ref.: 1602.	<i>Bahaba polykladiskos</i> (Bleeker 1852)	Yellc...
	<i>Scarus japonensis</i> (Bloch 1789)	Palecheek parrotfish, (M), Ref. 2689. Max. 25 cm SL. Also Ref.: 9137.	<i>Chrysochir aureus</i> (Richardson 1846)	Spin...
	<i>Scarus javanicus</i> Bleeker 1854	Java parrotfish, (M), Ref. 1602. Max. 26 cm SL.	<i>Dendrophysa russelii</i> (Cuvier 1829)	Be...
	<i>Scarus microrhinos</i> Bleeker 1854	Steephead parrotfish, (M, Dan), Ref. 2334. Max. 70 cm TL. Also Ref.: 8631. Known from Flores.	<i>Johnius amblycephalus</i> (Bleeker 1855)	Bea...
	<i>Scarus niger</i> Forsskål 1775	Dusky parrotfish, (M), Ref. 1602. Max. 35 cm TL.	<i>Johnius australis</i> (Günther 1880)	M...
	<i>Scarus oviceps</i> Valenciennes 1840	Darkcapped parrotfish, (M), Ref. 1602. Max. 30 cm TL.	<i>Johnius belangerii</i> (Cuvier 1830)	sc...
	<i>Scarus prasiognathos</i> Valenciennes 1840	Singapore parrotfish, (M), Ref. 8631. Max. 50 cm SL. Known from Bali.	<i>Johnius borneensis</i> (Bleeker 1851)	Bottl...
	<i>Scarus psittacus</i> Forsskål 1775	Common parrotfish, (M), Ref. 1602. Max. 30 cm TL.	<i>Johnius carouna</i> (Cuvier 1830)	W...
	<i>Scarus pyrrhurus</i> (Jordan & Seale 1906)	(M), Ref. 2689. Max. 25 cm SL.	<i>Johnius coitor</i> (Hamilton 1822)	Bel...
	<i>Scarus quoyi</i> Valenciennes 1840	Quoy's parrotfish, (M), Ref. 8631. Max. 17 cm SL. Known from Flores.	<i>Johnius heterolepis</i> Bleeker 1873	Mi...
	<i>Scarus rivulatus</i> Valenciennes 1840	Rivulated parrotfish, (M), Ref. 2689. Max. 40 cm SL. Museum: LPPL JIF125, from Bali Strait to Timor Sea (Ref. 5978).	<i>Johnius hypostoma</i> (Bleeker 1853)	Fr...
	<i>Scarus rubroviolaceus</i> (Bleeker 1847)	Ember parrotfish, (M), Ref. 1602. Max. 70 cm TL.	<i>Johnius latifrons</i> Sasaki 1992	7C...
	<i>Scarus schlegeli</i> (Bleeker 1861)	Yellowband parrotfish, (M), Ref. 1602. Max. 40 cm TL.	<i>Johnius macropterus</i> (Bleeker 1853)	Shar...
	<i>Scarus sordidus</i> Forsskål 1775	Daisy parrotfish, (M), Ref. 1602. Max. 40 cm TL.	<i>Johnius macrorhynchus</i> (Mohan 1975)	Ma...
	<i>Scarus spinus</i> (Kner 1868)	Greensnout parrotfish, (M), Ref. 1602. Max. 30 cm TL.	<i>Johnius novaeguineae</i> (Nichols 1950)	sa...
	<i>Scarus strongylocephalus</i> Bleeker 1854	Indian Ocean steephead parrotfish, (M), Ref. 2334 Max. 50 cm.	<i>Johnius pacificus</i> Hardenberg 1941	Caro...
	<i>Scarus tricolor</i> Bleeker 1847	Tricolour parrotfish, (M), Ref. 8631. Max. 40 cm SL. Known from Bali.	<i>Johnius plagiotoma</i> (Bleeker 1850)	Coitc...
	<i>Scarus troschellii</i> Bleeker 1853	Troschel's parrotfish, (M), Ref. 2689. Max. 34.6 cm TL. Known from Seribu Islands.	<i>Johnius trachycephalus</i> (Bleeker 1851)	Larg...
	<i>Scarus viridifurcatus</i> (Smith 1956)	Round-head parrotfish, (M), Ref. 9232. Max. 26 cm SL.	<i>Johnius weberi</i> Hardenberg 1936	Smal...
	Scatophagidae	Scats		Broa...
	<i>Scatophagus argus</i> (Linnaeus 1766)	Spotted scat, (M, Fi), Ref. 1602. Max. 38 cm TL.		Larg...
	Sciaenidae	Drums or croakers		Mu...
<i>Argyrosomus amoyensis</i> (Bleeker 1863)	Amoy croaker, (M, Fi), Ref. 5978. Max. 40 cm SL. Museum: BMNH 1984.1.1.75 (TGT (PJPW) 747. Off Java. In range Ref.: 3490.		St...	

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	Sumatera, NMW 39989. Java, ZMH 13892. Bali, AMNH 15008. Lombok, BPBM 30075. Flores, BPBM 32186. Bonarate, RMNH 29818. Sulawesi, AMNH 19877. Borneo, FMNH 22518. Also Ref.: 4787, 8631.	<i>Epinephelus miliaris</i> (Valenciennes 1830)	Netfin
<i>Epinephelus faveatus</i> (Valenciennes 1828)	Barred-chest grouper, (M, Fi), Ref. 5222. Max. 32 cm TL. Also Ref.: 4787. Museum: Bali, BPBM 28953. Lombok, BPBM 29921.	<i>Epinephelus morrhua</i> (Valenciennes 1833)	Also RMH Comet
<i>Epinephelus flavocaeruleus</i> (Lacepède 1802)	Blue and yellow grouper, (M, Fi), Ref. 5222. Max. 80 cm TL. Also Ref.: 4787. Museum: Boelveweh, Sabang Island (off Sumatera), RMNH 12064.	<i>Epinephelus ongus</i> (Bloch 1790)	White- cm (Hol BMH Kari 300 BPE
<i>Epinephelus fuscoguttatus</i> (Forsskål 1775)	Brown-marbled grouper, (M, Fi, Dan), Ref. 5222. Max. 120 cm. Also Ref.: 2114, 4787. Museum: Sumatera, Padang, NMW 40522; ZMB 17741. Java RMNH 19; RMNH 2160. Batavia (Jakarta), NMW 40521. Sulawesi (Celebes), Janeponto, BPBM 26789. Molucca is., Halmahera, USNM 228042. Irian Jaya, USNM 245356.	<i>Epinephelus polyphekadion</i> (Bleeker 1849)	Camou Max Mus 548 170 Dara BPE
<i>Epinephelus heniochus</i> Fowler 1904	Bridled grouper, (M, Fi), Ref. 5222. Max. 35 cm SL. Also Ref.: 4787. Museum: Sumatera (Sumatra), Padang, ANSP 27557 (Holotype, Mentawe Strait), NTM S.10998-066, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Epinephelus quoyanus</i> (Valenciennes 1830)	Longfin Also A.5 Sum Jawa AMS BPE
<i>Epinephelus lanceolatus</i> (Bloch 1790)	Giant grouper, (M, Br, Fi, Dan), Ref. 5222. Max. 270 cm. Also Ref.: 4787, 7050. Museum: ZMB 169 (Holotype); ZMUC 71. Sumatra, NMW 40582-83. Java Sea, NMW 40578. Java, RMNH 132 (Holotype of <i>Serranus geographicus</i>). Batavia (Jakarta), ANSP 90467.	<i>Epinephelus retouti</i> Bleeker 1868	Red-tip Bali USN Tim
266 <i>Epinephelus longispinis</i> (Kner 1864)	Longspine grouper, (M, Fi), Ref. 5222. Max. 55 cm. Also Ref.: 4787. Museum: Lombok, BPBM 29853, 30126. Watubela Group, Uran Is., BPBM 34509. From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Epinephelus rivulatus</i> (Valenciennes 1830)	Halfm Also 313 rhy BPE
<i>Epinephelus macrospilos</i> (Bleeker 1855)	Snubnose grouper, (M, Fi), Ref. 5222. Max. 51 cm TL. Also Ref.: 4787. Museum: BMNH 1880.4.21.14; RMNH 31247; USNM 272427. Molluca Is., Batjan, RMNH 5503 (Holotype of <i>S. macrospilos</i>).	<i>Epinephelus sexfasciatus</i> (Valenciennes 1828)	Sixbar Also Jawa BPE Pan Is., 597
<i>Epinephelus maculatus</i> (Bloch 1790)	Highfin grouper, (M, Fi, Dan), Ref. 5222. Max. 60.5 cm TL. Also Ref.: 4787, 8631. Museum: Ambon, USNM 210459. CSIRO CA893, from Bali Strait to Timor Sea (Ref. 5978); including Flores.	<i>Epinephelus spilotoceps</i> Schultz 1953	Fours Also USN 466
<i>Epinephelus malabaricus</i> (Bloch & Schneider 1801)	Malabar grouper, (M, Br, Fi, Sport), Ref. 5222. Max. 234 cm. Also Ref.: 4787. Museum: BMNH 1880.4.21.3; MNHN 7332 (Holotype of <i>S. crapao</i>); BPBM 29852, 30025, 30125; ZMH 13950. LPPL JIF34, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Epinephelus stictus</i> Randall & Allen 1987	Black- In J
<i>Epinephelus melanostigma</i> Schultz 1953	One-blotch grouper, (M, Fi), Ref. 5222. Max. 33 cm TL. Also Ref.: 4787, 5978. Museum: north of Sumbawa, BPBM 28950.	<i>Epinephelus undulosus</i> (Quoy & Gaimard 1824)	Wavy- Also Sul RMH 546 Pul
<i>Epinephelus merra</i> Bloch 1793	Honeycomb grouper, (M, Fi), Ref. 5222. Max. 30 cm SL. Also Ref.: 4787. Museum: RMNH 8138. Timor, MNHN 7381. Banda Is., NMW 40644. Ambon, NMW 40641. Ternate, MNHN 5908. Sulawesi, MNHN 6431. Jawa, Jakarta, ANSP 90498; MNHN 7382. Sumatera, ZMB 7660. Puloweh, Sabang, ZMH 13924.		

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Gracila albomarginata</i> (Fowler & Bean 1930)	Masked grouper, (M, Fi), Ref. 5222. Max. 38 cm TL. Also Ref.: 4787, 8631. Museum: USNM 89985 (Holotype, Borneo, Kalimantan, Danawan Is., vicinity of Sibuko Bay). Also known from Flores.	<i>Pseudanthias bicolor</i> (Randall 1979)	Frc Bicolc Knr
<i>Grammistes sexlineatus</i> (Thunberg 1792)	Goldenstriped soapfish, (M), Ref. 5978. Max. 28 cm. Also Ref.: 8631. Museum: LPPLJIF38 (TGT1048). Known from Bali Strait to Timor Sea.	<i>Pseudanthias bimaculatus</i> (Smith 1955)	Two-s Ret
<i>Holanthias rhodopeplus</i> (Günther 1871)	(M), Ref. 5978. From southwest Sumatra to Bali Strait.	<i>Pseudanthias cichlops</i> (Bleeker 1853)	(M), F
<i>Liopropoma mitratum</i> Lubbock & Randall 1978	Pinstriped basslet, (M), Ref. 6180. Max. 7.2 cm SL. Museum: Ceram, USNM 210037.	<i>Pseudanthias cooperi</i> (Regan 1902)	Red-b Ret
<i>Liopropoma susumi</i> (Jordan & Seale 1906)	Meteor perch, (M), Ref. 6180. Max. 7.6 cm SL. Museum: Banda Is., USNM 243032. Sulawesi, Kabaena I., USNM 285947.	<i>Pseudanthias dispar</i> (Herre 1955)	Peach fror
<i>Liopropoma swalesi</i> (Fowler & Bean 1930)	(M), Ref. 6180. Museum: Sulawesi, USNM 89983 (Holotype of <i>Chorististium swalesi</i>); USNM 93393 (Paratype). Molucca I., Saparua, USNM 209922.	<i>Pseudanthias fasciata</i> (Kamohara 1954)	(M), F (TC nev
<i>Luzonichthys taeniatus</i> Randall & McCosker 1992	(M), Ref. 8524. Museum: Banda Sea, Lucipara Is., Penyu Group, Kadola I., BPBM 32336 (Holotype). Penyu Group, CAS 62527 and USNM 323627 (Paratypes).	<i>Pseudanthias huchtii</i> (Bleeker 1857)	Red-c Alsc
<i>Luzonichthys waitei</i> (Fowler 1931)	Waite's splitfin, (M), Ref. 2334. Max. 7 cm. Also Ref.: 8524. Museum: Widi Is., Dodoro I., BPBM 34218. Molucca Is., Suparua Reef (off Kulor), USNM 210395. Great Banda I., USNM 218812. Flores Sea, Kakabea I., BPBM 32360. Flores, Pulau Besar, CAS 62480. Maumere (off fuel dock), CAS 62496.	<i>Pseudanthias hypselosoma</i> Bleeker 1878	Stocky Flor
<i>Plectranthias japonicus</i> (Steindachner 1884)	Japanese perchlet, (M), Ref. 5978. Max. 15 cm TL. Museum: BPBM 28951 (TGT3149). From Bali Strait to Timor Sea.	<i>Pseudanthias lori</i> (Lubbock & Randall 1976)	Lori's . 160
<i>Plectranthias longimanus</i> (Weber 1913)	Longfin perchlet, (M), Ref. 1602. Max. 2.8 cm SL.	<i>Pseudanthias luzonensis</i> (Katayama & Masuda 1983)	Yellow Max Bali
<i>Plectranthias sagamiensis</i> (Katayama 1963)	(M), Ref. 5978. Max. 6 cm SL. Museum: RUSI 19990 (TGT3126). From Bali Strait to Timor Sea.	<i>Pseudanthias pleurotaenia</i> (Bleeker 1857)	Squar Alsc
<i>Plectranthias wheeleri</i> Randall 1980	(M), Ref. 7300. Museum: Celebes, BPBM 22401 (Holotype).	<i>Pseudanthias randalli</i> (Lubbock & Allen 1978)	Randa Alsc
<i>Plectranthias winniensis</i> (Tyler 1966)	(M), Ref. 7300. Max. 4 cm SL.	<i>Pseudanthias rubrizonatus</i> (Randall 1983)	Red-br Mus Tim
<i>Plectropomus areolatus</i> Rüppell 1830	Squartail coralgroup, (M, Fi, Dan), Ref. 5222 Max. 60 cm SL.	<i>Pseudanthias smithvanizi</i> (Randall & Lubbock 1981)	Prince
<i>Plectropomus laevis</i> (Lacepède 1801)	Blacksaddled coralgroup, (M, Fi, Dan), Ref. 8631 Max. 100 cm SL. Known from Bali.	<i>Pseudanthias squamipinnis</i> (Peters 1855)	Sea gc 863 002
<i>Plectropomus leopardus</i> (Lacepède 1802)	Leopard coralgroup, (M, Fi, Sport, Dan), Ref. 5222 Max. 68 cm SL. Known from Bali Strait to Timor Sea (Ref. 5978); including Lombok and Flores. Also Ref.: 2114, 4787, 8631.	<i>Pseudanthias tuka</i> (Herre & Montalban 1927)	Yellow: TL .
<i>Plectropomus maculatus</i> (Bloch 1790)	Spotted coralgroup, (M, Fi, Aq), Ref. 5222. Max. 100 cm SL. Also Ref.: 4787.	<i>Pseudanthias ventralis</i> (Randall 1979)	Long-fi
<i>Plectropomus oligacanthus</i> (Bleeker 1854)	Highfin coralgroup, (M, Fi, Dan), Ref. 5222. Max. 75 cm TL. Also Ref.: 8631. Known from Flores.	<i>Serranocirrhites latus</i> Watanabe 1949	Hawkfi Kno
<i>Plectropomus pessuliferus</i> Fowler 1904	Roving coralgroup, (M, Fi, Sport), Ref. 5222. Max. 120 cm TL. Museum: AMS I.24048-001, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Variola albimarginata</i> Baissac 1952	White-r Max RM Molu S.11 (Ref
<i>Plectropomus punctatus</i> Quoy & Gaimard 1824	Marbled coralgroup, (M, Fi), Ref. 5978. Max. 96 cm TL. Museum: BMNH 1984.1.1.18 (TGT (PJPW)818a).	<i>Variola louti</i> (Forsskål 1775)	Yellow- Max (TG Also
		Siganidae	Rabbit
		<i>Siganus argenteus</i> (Quoy & Gaimard 1825)	Stream Max

Table 2. Marine and brackishwater fishes of Indonesia .

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

	1984.1.1.95, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Sillago macrolepis</i> Bleeker 1859	Large-s cm SL
<i>Siganus canaliculatus</i> (Park 1797)	White-spotted spinefoot, (M, Br, Fi, Aq, Dan), Ref. 1419 Max. 25 cm TL. Also Ref.: 8631. Museum: LPPL JIF135, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Sillago nierstraszi</i> Hardenberg 1941	Rough s
<i>Siganus corallinus</i> (Valenciennes 1835)	Blue-spotted spinefoot, (M, Fi, Dan), Ref. 1419. Max. 30 cm TL. Also Ref.: 8631. Museum: LPPL JIF136, from southwest Sumatra to Timor Sea (Ref. 5978).	<i>Sillago sihama</i> (Forsskål 1775)	Silver sil Museu Timor In ran
<i>Siganus doliatus</i> Cuvier 1830	Barred spinefoot, (M, Dan), Ref. 1419. Max. 25 cm TL.	Sparidae	Porgies
<i>Siganus fuscescens</i> (Houttuyn 1782)	Mottled spinefoot, (M, Fi, Dan), Ref. 1419. Max. 40 cm TL.	<i>Acanthopagrus berda</i> (Forsskål 1775)	Picnic se TL. Al
<i>Siganus guttatus</i> (Bloch 1787)	Orange-spotted spinefoot, (M, Br, Fi, Dan), Ref. 1419 Max. 45 cm TL. Also Ref.: 8631. Including Irian Jaya. Museum: LPPL JIF137, from Bali Strait to Timor Sea (Ref. 5978).	<i>Acanthopagrus bifasciatus</i> (Forsskål 1775)	Twobar FL.
<i>Siganus javus</i> (Linnaeus 1766)	Streaked spinefoot, (M, Br, Fi, Dan), Ref. 1419 Max. 55 cm TL. Also Ref.: 8631. Known from Bali.	<i>Acanthopagrus latus</i> (Houttuyn 1782)	Yellowfir Also I
<i>Siganus labyrinthos</i> (Bleeker 1853)	Labyrinth spinefoot, (M, Dan), Ref. 1419. Max. 25 cm TL. Java and Moluccas.	<i>Argyrops spinifer</i> (Forsskål 1775)	King sol Muse Timor
<i>Siganus lineatus</i> (Valenciennes 1835)	Golden-lined spinefoot, (M, Br, Fi, Dan), Ref. 1419 Max. 45 cm TL.	<i>Rhabdosargus sarba</i> (Forsskål 1775)	Goldline Max.
<i>Siganus puellus</i> (Schlegel 1852)	Masked spinefoot, (M, Fi, Dan), Ref. 1419 Max. 38 cm TL. Also Ref.: 8631. Known from Flores.	<i>Taius tumifrons</i> (Temminck & Schlegel 1843)	Yellowb SL. M south
208 <i>Siganus punctatissimus</i> Fowler & Bean 1929	Peppered spinefoot, (M), Ref. 1419. Max. 35 cm TL. Northern Indonesia.	Sphyraenidae	Barracu
<i>Siganus punctatus</i> (Schneider 1801)	Goldspotted spinefoot, (M, Fi, Dan), Ref. 1419 Max. 45 cm SL. Also Ref.: 8631. Museum: LPPL JIF138, from Bali Strait to Timor Sea (Ref. 5978).	<i>Sphyraena barracuda</i> (Walbaum 1792)	Great b 200 c JIF20 5978
<i>Siganus spinus</i> (Linnaeus 1758)	Little spinefoot, (M, Fi, Dan), Ref. 1419. Max. 23 cm SL. Also Ref.: 8631. Museum: NTM S.11044-001, from Bali Strait to Timor Sea (Ref. 5978).	<i>Sphyraena forsteri</i> Cuvier 1829	Bigeye Also south Heller's
<i>Siganus vermiculatus</i> (Valenciennes 1835)	Vermiculated spinefoot, (M, Br, Fi, Dan), Ref. 1419 Max. 45 cm TL. Also Ref.: 8631. Known from Bali.	<i>Sphyraena helleri</i> Jenkins 1901	Pickhar 150 c
<i>Siganus virgatus</i> (Valenciennes 1835)	Barhead spinefoot, (M, Br, Fi, Dan), Ref. 1419 Max. 33 cm TL. Also Ref.: 8631. Museum: NTM S.1075-004, from southwest Sumatra to Bali Strait (Ref. 5978). Includes one record for western Irian Jaya (Ref. 1419).	<i>Sphyraena jello</i> Cuvier 1829	Obtuse Max. JIF11
<i>Siganus vulpinus</i> (Schlegel & Müller 1845)	Foxface, (M, Fi, Dan), Ref. 1419. Max. 25 cm TL. Also Ref.: 8631. Known from Flores.	<i>Sphyraena obtusata</i> Cuvier 1829	Sawtoo cm T Fror
Sillaginidae	Smelt-whittings	<i>Sphyraena putnamiae</i> Jordan & Seale 1905	Butterf
<i>Sillaginopsis panijus</i> (Hamilton 1822)	Flathead sillago, (M, Br, Fi), Ref. 6205. Max. 35 cm . Also Ref.: 7050.	Stromateidae	Silver p Muse south Ref.:
<i>Sillago aeolus</i> Jordan & Evermann 1902	Oriental sillago, (M, Fi), Ref. 6205. Max. 30 cm SL.	<i>Pampus argenteus</i> (Euphrasen 1788)	Chines Max. south
<i>Sillago burrus</i> Richardson 1842	Western trumpeter sillago, (M, Br, Fi), Ref. 6205 Max. 36 cm SL. Museum: NTM S.10754-002 (TGT1179). From southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 7050.	<i>Pampus chinensis</i> (Euphrasen 1788)	
<i>Sillago chondropus</i> Bleeker 1849	Clubfoot sillago, (M, Br, Fi), Ref. 6205. Max. 35 cm SL.	Symphysanodontidae	Longtai Mus- to Ti
		<i>Symphysanodon maunaloae</i> Anderson 1970	

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Symphysanodon typus</i> Bleeker 1878	Insular shelf beauty, (M, Fi), Ref. 5978. Max. 17 cm SL. Museum: GMBL81-162 (TGT1461). From Bali Strait to Timor Sea.	Trichonotidae <i>Trichonotus setiger</i> (Bloch & Schneider 1801)	Sanc Spot 23
Terapontidae	Grunters or tigerperches, thornfishes	Tripterygiidae	Thre
<i>Mesopristes argenteus</i> (Cuvier 1829)	(M, Br, Fr), Ref. 7050. Max. 27.5 cm SL. Sumatra, Java, Sulawesi, and Irian Jaya.	<i>Helcogramma ellioti</i> (Herre 1944)	(M), 16
<i>Pelates quadrilineatus</i> (Bloch 1790)	Fourlined terapon, (M, Br, Fi), Ref. 559. Max. 24 cm TL. Museum: NTM S.10733-033 (TGT1282). From Bali Strait to Timor Sea (Ref. 5978), to New Guinea (Ref. 559). Also Ref.: 7050.	<i>Helcogramma hudsoni</i> (Jordan & Seale 1906)	(M)
<i>Terapon jarbua</i> (Forsskål 1775)	Jarbua terapon, (M, Br, Fr, Fi), Ref. 5978. Max. 33 cm TL. Also Ref.: 7050, 8631. Museum: LPPL JIF40 (TGT2614). From southwest Sumatra to Timor Sea (Ref. 5978).	<i>Helcogramma obtusirostris</i> (Klunzinger 1871)	Hotlij
<i>Terapon puta</i> (Cuvier 1829)	Small-scaled terapon, (M, Br, Fi), Ref. 5978. Max. 16 cm TL. Museum: NTM S.10744-006 (TGT926). From Bali Strait to Timor Sea. Also Ref.: 7050.	<i>Helcogramma springeri</i> Hansen 1986	(M), (h 21
<i>Terapon theraps</i> (Cuvier 1829)	Largescaled terapon, (M, Br, Fi), Ref. 5978. Max. 30 cm SL. Also Ref.: 8631. Museum: LPPL JIF39 (TGT2213). Known from southwest Sumatra to Timor Sea; including Flores.	<i>Helcogramma striata</i> Hansen 1986	(M), M 22
Toxotidae	Archerfishes	<i>Helcogramma trigloides</i> (Bleeker 1858)	(M),
<i>Toxotes chatareus</i> (Hamilton 1822)	Spotted archerfish, (M, Br, Fr, Fi), Ref. 7294. Max. 40 cm TL. Sumatra, Kalimantan, and Irian Jaya. Also Ref.: 7050.	<i>Helcogramma vulcanum</i> Randall & Clark 1993	(M, E ar C,
Trichiuridae	Cutlassfishes	Uranoscopidae	Star
<i>Assurger anzac</i> (Alexander 1917)	Razorback scabbardfish, (M), Ref. 6181. Max. 250 cm SL.	<i>Gnathagnus elongatus</i> (Temminck & Schlegel 1843)	(M), 0C Se
<i>Benthodesmus macrophthalmus</i> Parin & Becker 1970	Bigeye frostfish, (M), Ref. 6181. Max. 50 cm SL.	<i>Uranoscopus cognatus</i> Cantor 1849	Two- M to
<i>Benthodesmus neglectus</i> Parin 1976	Neglected frostfish, (M), Ref. 6181. Max. 23 cm SL.	<i>Uranoscopus kaianus</i> Günther 1880	Kai s BI sc
<i>Benthodesmus tenuis</i> (Günther 1877)	Slender frostfish, (M), Ref. 6181. Max. 230 cm SL.	<i>Uranoscopus oligolepis</i> Bleeker 1878	(M), (T
<i>Benthodesmus tuckeri</i> Parin & Becker 1970	Tucker's frostfish, (M), Ref. 6181. Max. 77 cm SL. Known from the Molucca Islands and south of Java (Ref. 6181).	<i>Uranoscopus sulphureus</i> Valenciennes 1832	Whit M
<i>Benthodesmus vityazi</i> Parin & Becker 1970	Vityaz' frostfish, (M), Ref. 6181. Max. 77 cm SL.	Xiphiidae	Swc Swc
<i>Eupleurogrammus glossodon</i> (Bleeker 1860)	Longtooth hairtai I, (M, Fi), Ref. 6181 Max. 50 cm TL.	<i>Xiphias gladius</i> Linnaeus 1758	Swc Swc
<i>Eupleurogrammus muticus</i> (Gray 1831)	Smallhead hairtail, (Ni, Fi), Ref. 6181 Max. 70 cm TL.	Zanclidae	Moc Moo R B
<i>Lepturacanthus savala</i> (Cuvier 1829)	Savalani hairtail, (M, Fi), Ref. 6181 Max. 100 cm SL.	<i>Zanclus cornutus</i> (Linnaeus 1758)	Eelp
<i>Tentoriceps cristatus</i> (Klunzinger 1884)	Crested hairtail, (M, Fi), Ref. 6181. Max. 90 cm TL. Museum: NTM S.11034-001, from southwest Sumatra to Fimor Sea (Ref. 5978).	Zoarcidae	(M), I: Ti
<i>Trichiurus auriga</i> Klunzinger 1884	Pearly hairtail, (M), Ref. 6181. Max. 35 cm TL.	<i>Lycodes agulhensis</i> Andriashev 1959	
<i>Trichiurus lepturus</i> Linnaeus 1758	Largehead hairtail, (N 1, Br, Fi, Sport), Ref. 6181 Max. 150 cm SL. Museum: NTM S. 1 1005-005 (TGT2214). From southwest Sumatra to Timor Sea (Ref. 5978). Also Ref.: 6567.	Gobiesociformes (clingfishes)	Clin Pale G
		Gobiesocidae	
		<i>Lepadichthys caritus</i> Briggs 1969	

Table 2. Marine and brackishwater fishes of Indonesia.
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Lepadichthys minor</i> Briggs 1955	(M), Ref. 1602. Known from Moluccas.	<i>Psettina profunda</i> (Weber 1913)	(M), F Jav
Pleuronectiformes (flatfishes)		Citharidae	Cithar
Bothidae	Lefteye flounders	<i>Brachypleura novaezeelandiae</i> Günther 1862	Yellow Re
<i>Amoglossus aspidos</i> (Bleeker 1851)	Spotless lefteye flounder, (M), Ref. 9824. Max. 19 cm TL.	<i>Lepidoblepharon ophthalmolepis</i> Weber 1913	Scale Kne 979
<i>Amoglossus daigleishi</i> (von Bonde 1922)	East coast flounder, (M), Ref. 5978. Max. 18.5 cm TL. Museum: BMNH 1984.1.1.103 (PJPW28). From southwest Sumatra to Bali Strait.	Cynoglossidae	Tong
<i>Amoglossus debilis</i> (Gilbert 1905)	Weak lefteye flounder, (M), Ref. 5978. Max. 17 cm TL. Museum: NTM S.10760-007 (TGT1745). From Bali Strait to Timor Sea.	<i>Cynoglossus abbreviatus</i> (Gray 1835)	Three Mu sou
<i>Amoglossus elongatus</i> Weber 1913	Long lefteye flounder, (M), Ref. 9824. Max. 11 cm TL. Known from Madura.	<i>Cynoglossus arel</i> (Bloch & Schneider 1801)	Large TL. sou
<i>Amoglossus polyspilus</i> (Günther 1880)	Many-spotted lefteye flounder, (M), Ref. 9824. Max. 24 cm TL.	<i>Cynoglossus bilineatus</i> (Bloch 1787)	Fourli
<i>Amoglossus tapinosoma</i> (Bleeker 1865)	Large-crested lefteye flounder, (M), Ref. 4900 Max. 13 cm TL. Reported from Sumatra (Padang), Java Sea.	<i>Cynoglossus borneensis</i> (Bleeker 1858)	(M), F
<i>Asterorhombus intermedius</i> (Bleeker 1865)	Intermediate flounder, (M), Ref. 3132. Max. 15 cm TL. Known from the Java Sea (Ref. 1602).	<i>Cynoglossus cynoglossus</i> (Hamilton 1822)	Beng TL.
<i>Bothus myriaster</i> (Temminck & Schlegel 1846)	Oval flounder, (M), Ref. 5978. Max. 27 cm TL. Museum: BMNH 1984.1.1.104 (TGT (PJPW)694). From southwest Sumatra to Bali Strait.	<i>Cynoglossus kopsii</i> (Bleeker 1851)	(M), F
<i>Bothus pantherinus</i> (Rüppell 1830)	Leopard flounder, (M, Fi), Ref. 5978. Max. 27.5 cm TL. Museum: NTM S.10733-013 (TGT1100). From Bali Strait to Timor Sea.	<i>Cynoglossus lida</i> (Bleeker 1851)	Rough cm
<i>Chascanopsetta lugubris</i> Alcock 1894	Pelican flounder, (M, Fi), Ref. 5978. Max. 38 cm SL. Museum: BMNH 1984.1.1.106. From southwest Sumatra to Bali Strait.	<i>Cynoglossus lingua</i> Hamilton 1822	Long
<i>Crossorhombus azureus</i> (Alcock 1889)	Blue flounder, (M), Ref. 9824. Max. 18 cm TL. Known from the Aru Islands.	<i>Cynoglossus monopus</i> (Bleeker 1849)	(M), F
<i>Engyprosopon grandisquama</i> (Temminck & Schlegel 1846)	Largescale flounder, (M, Fi), Ref. 5978. Max. 15 cm. Museum: CSIRO CA3066. From southwest Sumatra to Bali Strait.	<i>Cynoglossus puncticeps</i> (Richardson 1846)	Speck cm
<i>Engyprosopon mogkii</i> (Bleeker 1854)	(M), Ref. 9824. Max. 11 cm SL.	<i>Cynoglossus suyeni</i> Fowler 1934	(M), F
<i>Grammatobothus polyophthalmus</i> (Bleeker 1865)	Threespot flounder, (M), Ref. 4900. Max. 21 cm TL.	<i>Paraplagusia bilineatus</i> (Bloch 1787)	Rock
<i>Kamoharaia megastoma</i> (Kamohara 1936)	Wide-mouthed flounder, (M), Ref. 9824. Max. 22.5 cm TL.	Paralichthyidae	Large
<i>Laeops guentheri</i> Alcock 1890	Günther's flounder, (M), Ref. 5978. Max. 14 cm TL. Museum: NTM S.10751-003 (TGT1674). From Bali Strait to Timor Sea.	<i>Pseudorhombus argus</i> Weber 1913	Peacc
<i>Neolaeops microphthalmus</i> (von Bonde 1922)	Crosseyed flounder, (M), Ref. 5978. Max. 21 cm SL. Museum: NTM S.10998-004 (TGT2526). From southwest Sumatra to Bali Strait.	<i>Pseudorhombus arsius</i> (Hamilton 1822)	Large Ma
<i>Parabothus kiensis</i> (Tanaka 1918)	(M), Ref. 9824. Max. 20.3 cm SL. Known from southern Indonesia.	<i>Pseudorhombus diplospilus</i> Norman 1926	Four t
<i>Psettina brevirictis</i> (Alcock 1890)	(M), Ref. 4417. Max. 8 cm SL. Known from Celebes (Ref. 9824).	<i>Pseudorhombus dupliocellatus</i> Regan 1905	Ocella Mu Fro
<i>Psettina gigantea</i> Amaoka 1963	Rough-scaled flounder, (M), Ref. 9824. Max. 13 cm SL.	<i>Pseudorhombus elevatus</i> Ogilby 1912	Deep Ma Ref
<i>Psettina iijimae</i> (Jordan & Starks 1904)	(M), Ref. 9824. Max. 8.5 cm SL. Known from southern Indonesia.	<i>Pseudorhombus javanicus</i> (Bleeker 1853)	Javan Mu sou wes
		<i>Pseudorhombus malayanus</i> Bleeker 1865	Malay Mu sou

Table 2. Continuation.
[Tabel 2. Sambungan.]

<i>Pseudorhombus megalops</i> Fowler 1934	(M), Ref. 2688. Max. 20 cm SL. Museum: Indian Ocean (south coasts of Sumatra, Java, and Lombok), HUMZ 111768 - 69; NTM 10760-006. Bali Strait, NTM S.11022-002. Arafura Sea, CSIRO CA2526.	<i>Euryglossa orientalis</i> (Bloch & Schneider 1801)	Orie
<i>Pseudorhombus neglectus</i> Bleeker 1865	(M), Ref. 5978. Max. 25 cm SL. Museum: BMNH 1984.1.1.105. From southwest Sumatra to Timor Sea.	<i>Liachirus melanospilus</i> (Bleeker 1854)	(M)
<i>Pseudorhombus pentophthalmus</i> Günther 1862	Fivespot flounder, (M, Fi), Ref. 559. Max. 18 cm SL. Known from the Java Sea northwards (Ref. 9774).	<i>Pardachirus pavoninus</i> (Lacepède 1802)	Pea M
<i>Pseudorhombus polyspilus</i> (Bleeker 1853)	(M), Ref. 4900	<i>Solea ovata</i> Richardson 1846	Ova S
<i>Pseudorhombus quinquocellatus</i> Weber & de Beaufort 1929	Five-eyed flounder, (M), Ref. 9774. Max. 20 cm SL.	<i>Synaptura commersoniana</i> (Lacepède 1802)	(M)
<i>Pseudorhombus triocellatus</i> (Bloch & Schneider 1801)	Three spotted flounders, (M, Fi), Ref. 5978. Max. 15 cm SL. Museum: BMNH 1984.1.1.107 (TGT (PJPW)750). From southwest Sumatra to Bali Strait.	<i>Zebrias quagga</i> Kaup 1858	Fri
Pleuronectidae	Righteye flounders	<i>Zebrias zebra</i> (Bloch & Schneider 1787)	Zeb
<i>Nematops grandisquama</i> Weber & de Beaufort 1929	Large-scale righteye flounder, (M), Ref. 9792. Max. 9 cm TL. Known from Bali.	Tetraodontiformes (puffers and filefishes)	
<i>Nematops macrochirus</i> Norman 1931	Long-fin righteye flounder, (M), Ref. 9792. Max. 15 cm TL. Known from Bali Strait.	Balistidae	Trig
<i>Poecilopsetta colorata</i> Günther 1880	Coloured righteye flounder, (M), Ref. 5978. Max. 17 cm TL. Museum: NTM S.10760-008 (TGT1746). From Bali Strait to Timor Sea.	<i>Abalistes stellaris</i> (Bloch & Schneider 1801)	Sta T
<i>Poecilopsetta praelonga</i> Alcock 1894	Alcock's narrow-body righteye flounder, (M), Ref. 5978 Max. 17.5 cm TL. Museum: NTM S.10742-001, WAM P.26208-021. From southwest Sumatra to Timor Sea.	<i>Balistapus undulatus</i> (Park 1797)	Ora A F
<i>Psammodiscus ocellatus</i> Günther 1862	(M), Ref. 3132. Max. 15 cm TL.	<i>Balistoides conspicillum</i> (Bloch & Schneider 1801)	Clo M S
<i>Samaris cristatus</i> Gray 1831	Cockatoo righteye flounder, (M), Ref. 5978. Max. 22 cm TL. Museum: WAM P.26200-007. From Bali Strait to Timor Sea.	<i>Balistoides viridescens</i> (Bloch & Schneider 1801)	Tita A F
<i>Samariscus huysmani</i> Weber 1913	Huysman's righteye flounder, (M), Ref. 5978 Max. 11.5 cm TL. Museum: BMNH 1984.1.1.108 (PJPW27). From southwest Sumatra to Bali Strait.	<i>Canthidermis maculatus</i> (Bloch 1786)	Sp T (
<i>Samariscus maculatus</i> (Günther 1880)	Spotted righteye flounder, (M), Ref. 9792. Max. 10 cm TL. Known from Kei Islands.	<i>Melichthys indicus</i> Randall & Klauswitz 1973	Indi E
<i>Samariscus sunieri</i> Weber & de Beaufort 1929	Sunier's righteye flounder, (M), Ref. 9792. Max. 13 cm TL. Known from Bali.	<i>Melichthys vidua</i> (Solander 1844)	Pin H
Psettodidae	Psettodids	<i>Odonus niger</i> (Rüppell 1836)	Rec A F
<i>Psettodes erumei</i> (Bloch & Schneider 1801)	Indian spiny turbot, (M, Fi), Ref. 5978. Max. 64 cm. Also Ref.: 3415, 9987. Museum: LPPL JIF145. From southwest Sumatra to Timor Sea.	<i>Pseudobalistes flavimarginatus</i> (Rüppell 1829)	Yell M S
Soleidae	Soles	<i>Pseudobalistes fuscus</i> (Bloch & Schneider 1801)	Yell T S
<i>Aesopia cornuta</i> Kaup 1858	Unicorn sole, (M), Ref. 5978. Max. 20 cm SL. Museum: CSIRO CA2097, CSIRO CA1494. From southwest Sumatra to Bali Strait.	<i>Rhinecanthus aculeatus</i> (Linnaeus 1758)	Wh M T
<i>Aseraggodes cyaneus</i> (Alcock 1890)	(M), Ref. 5978. Max. 8.3 cm SL. Museum: NTM S.10745-005 (TGT1889). From Bali Strait to Timor Sea.	<i>Rhinecanthus rectangulus</i> (Bloch & Schneider 1801)	We
<i>Dexillus muelleri</i> (Steindachner 1879)	Tufted sole, (M), Ref. 561. Max. 18 cm.	<i>Rhinecanthus verrucosus</i> (Linnaeus 1758)	Bla

Table 2. Marine and brackishwater fishes of Indonesia .

[Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Sufflamen bursa</i> (Bloch & Schneider 1801)	Boomerang triggerfish, (M, Thr), Ref. 8631. Max. 25 cm TL. Known from Flores.	(Hollard 1854)	
<i>Sufflamen chrysopterus</i> (Bloch & Schneider 1801)	Halfmoon triggerfish, (M), Ref. 8631. Max. 22 cm TL. Known from Bali.	<i>Cantherhines fronticinctus</i> (Günther 1867)	Spe
<i>Sufflamen fraenatus</i> (Latreille 1804)	Masked triggerfish, (M), Ref. 5978. Max. 50 cm. Museum: LPPL JIF159 (TGT2415). From southwest Sumatra to Timor Sea.	<i>Cantherhines pardalis</i> (Rüppell 1837)	Hor
<i>Xanthichthys auromarginatus</i> (Bennett 1832)	Gilded triggerfish, (M), Ref. 8631. Max. 22 cm TL. Known from Flores.	<i>Chaetodermis penicilligera</i> (Cuvier 1817)	Pric
<i>Xanthichthys caeruleolineatus</i> Randall, Matsuura & Zama 1978	Blue-line triggerfish, (M), Ref. 1602. Max. 35 cm TL. Museum: BPBM 29340, from southwest Sumatra to Bali Strait (Ref. 5978).	<i>Monacanthus chinensis</i> (Osbeck 1765)	Far
Diodontidae	Porcupinefishes	<i>Oxymonacanthus longirostris</i> (Bloch & Schneider 1801)	Lon
<i>Chilomycterus reticulatus</i> (Linnaeus 1758)	Spotfin burrfish, (M, Dan), Ref. 5978. Max. 75 cm SL. Also Ref.: 8631. Museum: AMS I.22555-001 (TGT1370). From southwest Sumatra to Timor Sea.	<i>Paraluteres prionurus</i> (Bleeker 1851)	Fals
<i>Cyclichthys orbicularis</i> (Bloch 1785)	Birdbeak burrfish, (M, Sport), Ref. 5978. Max. 15 cm SL. Also Ref.: 8631. Museum: AMS I.23670-001 (TGT1509). From Bali Strait to Timor Sea.	<i>Paramonacanthus cryptodon</i> (Bleeker 1855)	(M)
<i>Cyclichthys spilostylus</i> (Leis & Randall 1982)	Spotbase burrfish, (M), Ref. 5978. Max. 35 cm SL. Museum: LPPL JIF208 (TGT1656). From southwest Sumatra to Timor Sea. In range Ref.: 4423.	<i>Paramonacanthus japonicus</i> (Tilesius 1810)	(M)
<i>Diodon holocanthus</i> Linnaeus 1758	Long-spine porcupinefish, (M), Ref. 5978. Max. 50 cm TL. Also Ref.: 8631. Museum: LPPL JIF168 (TGT2301). From southwest Sumatra to Timor Sea.	<i>Pervagor janthinosoma</i> (Bleeker 1854)	Blac
272 <i>Diodon hystrix</i> Linnaeus 1758	Spot-fin porcupinefish, (M, Dan), Ref. 5978. Max. 91 cm TL. Also Ref.: 8631. Museum: LPPL JIF169 (TGT2290). From southwest Sumatra to Timor Sea.	<i>Pervagor melanocephalus</i> (Bleeker 1853)	Rec
<i>Diodon lituosus</i> Shaw 1804	Black-blotched porcupinefish, (M), Ref. 8631. Max. 55 cm SL. Known from Flores.	<i>Pervagor nigrolineatus</i> (Herre 1927)	Blac
<i>Lophodiodon calori</i> (Bianconi 1854)	Four-bar porcupinefish, (M), Ref. 5978. Max. 30 cm SL. Museum: LPPL JIF170 (TGT3328). From Bali Strait to Timor Sea. Known from Java (Ref. 9680).	<i>Pseudalutarius nasicomis</i> (Temminck & Schlegel 1850)	Rhi
Molidae	Molas	<i>Pseudomonacanthus macrurus</i> (Bleeker 1857)	S
<i>Mola mola</i> (Linnaeus 1758)	Sunfish, (M, Fi), Ref. 4424. Max. 305 cm TL.	<i>Pseudomonacanthus peroni</i> (Hollard 1854)	Str
Monacanthidae	Filefishes	<i>Rudarius minutus</i> Tyler 1970	M
<i>Acreichthys radiatus</i> (Popta 1900)	Radial leatherjacket, (M), Ref. 559. Max. 7 cm TL.	<i>Thamnaconus striatus</i> (Kotthaus 1979)	Mar
<i>Acreichthys tomentosus</i> (Linnaeus 1758)	Bristle-tail file-fish, (M), Ref. 5978. Max. 8 cm SL. Also Ref.: 8631. Museum: WAM P.28199-001 (TGT1281). From Bali Strait to Timor Sea.	<i>Thamnaconus tessellatus</i> (Günther 1880)	P
<i>Aluterus monoceros</i> (Linnaeus 1758)	Unicorn leatherjacket, (M), Ref. 5978. Max. 75 cm TL. Also Ref.: 8631. Museum: LPPL JIF151 (TGT2261). From southwest Sumatra to Timor Sea.	Ostraciidae	(M)
<i>Aluterus scriptus</i> (Osbeck 1765)	Scrawled filefish, (M, Sport, Dan), Ref. 5978. Max. 110 cm. Also Ref.: 8631. Museum: LPPL JIH152 (TGT3252). From southwest Sumatra to Timor Sea.	<i>Lactoria comuta</i> (Linnaeus 1758)	C
<i>Amanses scopas</i> (Cuvier 1829)	Broom filefish, (M), Ref. 8631. Max. 20 cm TL. Known from Flores.	<i>Lactoria fornasini</i> (Bianconi 1846)	Box
<i>Anacanthus barbatus</i> Gray 1830	Bearded leatherjacket, (M), Ref. 5978. Museum: LPPL CSIRO CA1418 (conspecific material). From southwest Sumatra to Bali Strait.	<i>Ostracion cubicus</i> Linnaeus 1758	Lon
<i>Cantherhines dumerili</i>	Whitespotted filefish, (M), Ref. 583. Max. 35 cm TL.	<i>Ostracion meleagris</i> Shaw & Nodder 1796	M
		<i>Ostracion nasus</i> Bloch 1785	Tho
		<i>Ostracion rhinorhynchus</i>	M
			S
			Yell
			F
			E
			Whi
			S
			Sho
			F
			Hor

Table 2. Continuation.
 [Tabel 2. Sambungan.]

Bleeker 1852	From southwest Sumatra to Timor Sea.	<i>Lagocephalus inermis</i>	Smoc
<i>Ostracion solorensis</i>	Reticulate boxfish, (M, Dan), Ref. 1602. Max. 10 cm SL.	(Temminck & Schlegel 1850)	Mu
Bleeker 1853			Su
<i>Tetrosomus gibbosus</i>	Humpback turretfish, (M, Dan), Ref. 5978. Max. 30 cm	<i>Lagocephalus lagocephalus</i>	Ocea
(Linnaeus 1758)	TL. Museum: LPPL JIF161 (TGT2448). From	(Linnaeus 1758)	Mu
	southwest Sumatra to Timor Sea.		sol
<i>Tetrosomus reipublicae</i>	Smallspine turretfish, (M, Dan), Ref. 5978 Max. 30 cm	<i>Lagocephalus lunaris</i>	Greer
(Ogilby 1913)	TL. Museum: NTM S.10995-004 (TGT2611). From	(Bloch & Schneider 1801)	Ma
	southwest Sumatra to Timor Sea.		Frc
Tetraodontidae	Puffers	<i>Lagocephalus sceleratus</i>	Silver
<i>Arothron caeruleopunctatus</i>	(M), Ref. 9184. Max. 80 cm TL. From Bali Strait to Timor	(Gmelin 1789)	TL.
Matsuura 1994	Sea (Ref. 5978).		sol
<i>Arothron hispidus</i>	White-spotted puffer, (M, Br, Dan), Ref. 1602. Max. 45	<i>Lagocephalus spadiceus</i>	Half-s
(Linnaeus 1758)	cm SL. From southwest Sumatra to Timor Sea (Ref.	(Richardson 1845)	Mu
	5978).	<i>Sphoeroides pachygaster</i>	Blunth
<i>Arothron immaculatus</i>	Narrow-lined toadfish, (M, Br), Ref. 5170. Max. 22.2 cm	(Müller & Troschel 1848)	Mu.
(Bloch & Schneider 1801)	SL. Also Ref.: 8631. Museum: LPPL JIF166, from		to 1
	Bali Strait to Timor Sea (Ref. 5978).	<i>Torquigener brevipinnis</i>	(M, Da
<i>Arothron manilensis</i>	Narrow-lined puffer, (M, Br), Ref. 5170. Max. 33 cm TL.	(Regan 1903)	(TC
(de Procè 1822)	Also Ref.: 8631. Museum: CSIRO CA2234, from	<i>Torquigener hicksi</i>	Hick's
	southwest Sumatra to Timor Sea (Ref. 5978).	Hardy 1983	Mu:
<i>Arothron meleagris</i>	Guineafowl puffer, (M, Dan), Ref. 1602. Max. 40 cm SL.		Strä
(Bloch & Schneider 1801)		<i>Torquigener hypselogeneion</i>	Orang
<i>Arothron nigropunctatus</i>	Blackspotted puffer, (M, Dan), Ref. 8631. Max. 27 cm	(Bleeker 1852)	
(Bloch & Schneider 1801)	SL. Known from Flores.	<i>Torquigener parcuspinus</i>	Yellow
<i>Arothron reticularis</i>	Reticulated pufferfish, Reticulated puffer, (M, Br), Ref.	Hardy 1983	TL.
(Bloch & Schneider 1801)	9407. Max. 40 cm SL.		sou
<i>Arothron stellatus</i>	Starry toadfish, (M), Ref. 5978. Max. 84 cm SL.	<i>Torquigener tuberculiferus</i>	(M), R
(Bloch & Schneider 1801)	Museum: LPPL JIF167 (TGT2268). From southwest	(Ogilby 1912)	Fro
	Sumatra to Timor Sea.	<i>Tylerius spinosissimus</i>	Spiny
<i>Canthigaster compressa</i>	Compressed toby, (M, Br), Ref. 5978. Max. 8.7 cm SL.	(Regan 1908)	
(de Procè 1822)	Also Ref.: 8631. Museum: NTM S.10749-005	Triacanthidae	Triple:
	(TGT1816). From Bali Strait to Timor Sea.	<i>Pseudotriacanthus strigilifer</i>	Long-s
<i>Canthigaster coronata</i>	Crowned puffer, (M), Ref. 5978. Max. 13.5 cm TL.	(Cantor 1849)	Max
(Vaillant & Sauvage 1875)	Museum: NTM S.0747-001 (TGT3312). From Bali		Fro
	Strait to Timor Sea.	<i>Triacanthus biaculeatus</i>	Short-t
<i>Canthigaster epilampra</i>	Lantern toby, (M), Ref. 8631. Max. 10.9 cm. Known from	(Bloch 1786)	cm
(Jenkins 1903)	Bali.		Tim
<i>Canthigaster investigatoris</i>	(M), Ref. 5978. Museum: BMNH 1984.1.1.111,	<i>Triacanthus nieuhofii</i>	Silver
(Annandale & Jenkins 1910)	1984.1.1.112. From southwest Sumatra to Bali Strait.	Bleeker 1852	Mus
<i>Canthigaster janthinoptera</i>	Spotted puffer, (M), Ref. 8631. Max. 6.2 cm SL.		Sun
(Bleeker 1855)	Known from Bali.	<i>Tripodichthys angustifrons</i>	Black-
<i>Canthigaster leoparda</i>	Leopard sharpnose puffer, (M), Ref. 1602. Max. 5.6 cm	(Hollard 1854)	
Lubbock & Allen 1979	SL.	<i>Tripodichthys blochii</i>	Long-t
<i>Canthigaster ocellicincta</i>	Shy toby, Circle-barred puffer, (M), Ref. 2334	(Bleeker 1852)	TL.
Allen & Randall 1977	Max. 6.5 cm TL. From Flores (Ref. 8631).	<i>Tripodichthys oxycephalus</i>	Short-t
<i>Canthigaster solandri</i>	Spotted sharpnose, (M), Ref. 1602. Max. 11 cm TL.	(Bleeker 1851)	
(Richardson 1844)	Also Ref.: 8631. Known from Flores.	<i>Triphichthys weberi</i>	Blackti
<i>Canthigaster valentini</i>	Valentinni's sharpnose puffer, (M, Dan), Ref. 8631	(Chaudhuri 1910)	Mus
(Bleeker 1853)	Max. 11 cm TL. Known from Flores.		Sum
<i>Chelonodon patoca</i>	Milkspotted puffer, (M, Br, Fr), Ref. 3131. Max. 38 cm.	Triacanthodidae	Spiker
(Hamilton 1822)		<i>Atrophacanthus japonicus</i>	(M), R
<i>Lagocephalus gloveri</i>	(M, Dan), Ref. 5978. Max. 35 cm SL. Museum: NMNZ	(Kamohara 1941)	Fro
Abe & Tabeta 1983	P.15095 (TGT2117). From southwest Sumatra to	<i>Halimochirurgus alcocki</i>	(M), R
	Timor Sea.	Weber 1913	

Table 2. Marine and brackishwater fishes of Indonesia
 [Tabel 2. Ikan-ikan laut dan payau Indonesia.]

<i>Halimochirurgus centriscoides</i> Alcock 1899	Longsnout spikefish, (M), Ref. 5978. Max. 15 cm TL. Museum: BMNH 1984.1.1.109 (TGT (PJPW) 470). From southwest Sumatra to Bali Strait. Also Ref.: 3132.
<i>Macrorhamphosodes platycheilus</i> Fowler 1934	Trumpetsnout spikefish, (M), Ref. 5978. Max. 13 cm TL. Museum: NTM S.10998-009 (TGT2516). From southwest Sumatra to Bali Strait.
<i>Triacanthodes ethiops</i> Alcock 1894	Shortsnout spikefish, (M), Ref. 5978. Max. 8.5 cm SL. Museum: NTM S.10761-003 (TGT (PJPW) 515). From Bali Strait to Timor Sea. In range Ref. 6660.
<i>Tydemanina navigatoris</i> Weber 1913	Fleshy-lipped spikefish, (M), Ref. 6660. Max. 12 cm SL. Museum: BMNH 1984.1.1.110 (TGT (PJPW) 515). Found from southwest Sumatra to Timor Sea.
Triodontidae	Three-toothed puffer
<i>Triodon macropterus</i> Lesson 1830	Threetooth puffer, (M), Ref. 5978. Max. 54 cm TL. Museum: CSIRO CA801. From Bali Strait to Timor Sea.

References^a

- Allen, G.R. 1991. Damselfishes of the world. Mergus Publishers, Melle, Germany. 271 p.
- Eschmeyer, W.N. 1990. Catalog of the genera of recent fishes. California Academy of Sciences, San Francisco. 697 p.
- FAO. 1994. FAO yearbook, fishery statistics catches and landings 1992. Vol. 74. 677 p.
- Fischer, W. and P.J.P. Whitehead, Editors. 1974. FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Vols. 1-4. pag. var. FAO, Rome.
- Froese, R. and D. Pauly, Editors. 1995. FishBase: a biological database on fish (version 1.2). ICLARM, Manila, Philippines. 146 p.
- Froese, R. and D. Pauly, Editors. 1996. FishBase 96: Concepts, design and data sources. ICLARM, Manila, Philippines. 179 p.
- Gloerfelt-Tarp, T. and P.J. Kailola. 1984. Trawled fishes of Southern Indonesia and Northwestern Australia. Australian Development Assistance Bureau, Directorate General of Fisheries Indonesia, German Agency for Technical Cooperation. 406 p.
- Kottelat, M., A.J. Whitten, S.N. Kartikasari and S. Wirjoatmodjo. 1993. Freshwater fishes of Western Indonesia and Sulawesi. Periplus Editions, Hong Kong. 221 p. + 84 plates
- Kuiter, R.H. 1992. Tropical reef-fishes of the Western Pacific, Indonesia, and adjacent waters. PT Gramedia Pustaka Utama, Jakarta, Indonesia. 314 p.
- Myers, R.F. 1991. Micronesian reef fishes. A practical guide to the identification of the coral reef fishes of the tropical Central and Western Pacific. Coral Graphics, Guam. 298 p.
- Nelson, J.S. 1994. Fishes of the world. John Wiley and Sons, Inc., New York. 600 p.
- Randall, J.E., G.R. Allen and R.C. Steene. 1990. Fishes of the Great Barrier Reef and Coral Sea. University of Hawaii Press, Honolulu, Hawaii. 506 p.
- Randall, J.E. and P.C. Heemstra. 1991. Revision of Indo-Pacific groupers (Perciformes: Serranidae: Epinephelinae), with descriptions of five new species. Indo-Pac. 20, 332 p.
- Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea and W.B. Scott. 1991. World fishes important to North Americans. Exclusive of species from the continental waters of the United States and Canada. Am. Fish. Soc. Spec. Publ. (21): 243 p.
- Schuster, W.H. and R. Djajadiredja. 1952. Local common names of Indonesian fishes. W.W. Hoeve, Bandung, Indonesia. 276 p.
- Weber, M. and L.F. de Beaufort. 1911-1962. The fishes of the Indo-Australian Archipelago. Vols. 1-11. E.J. Brill Ltd., Leiden, Netherlands.

^aSee Appendix I, p. 284, for numbered (FishBase) references.

Using the NAN-SIS and FiSAT Software to Create a Trawl Survey Database for Western Indonesia^a

FRANCISCO TORRES, Jr., ANNADEL CABANBAN^b, SHERLYN BIENVENIDA
JOHN McMANUS, MARK PREIN and DANIEL PAULY^c

International Center for Living
Aquatic Resources Management
MCPO Box 2631, 0718 Makati City
Metro Manila, Philippines

TORRES, F., JR., A. CABANBAN, S. BIENVENIDA, J. McMANUS, M. PREIN and D. PAULY. 1996. Using the NAN-SIS and FiSAT software to create a trawl survey database for Western Indonesia [*Penggunaan perangkat lunak NAN-SIS dan FiSAT guna menciptakan suatu data-dasar survei trawl untuk Indonesia bagian barat*], p. 276-283. In D. Pauly and P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resources of Western Indonesia*. ICLARM Stud. Rev. 23, 312 p.

Abstract

This contribution presents the rationale for and some details on the creation of a database for the results of trawl surveys conducted from 1974 to 1981 by five research vessels (*R/Vs Mutiara 4, Jurong, Dr. Fridtjof Nansen, Bawal Putih 2 and Lemuru*), using the NAN-SIS software for the catch data, and the FiSAT software for the length-frequency data. These software — both distributed by the Food and Agriculture Organization (FAO) and maintained by their authors— allow further, detailed analysis of the data base made available herewith, and the readers are encouraged to perform such analyses using the contributions in this volume as examples and/or starting points.

Abstrak

Tulisan ini menyajikan alasan dan beberapa informasi rinci tentang pembentukan suatu data-dasar hasil-hasil survei yang dilaksanakan dari tahun 1974 hingga 1981 oleh lima kapal penelitian (kapal-kapal penelitian *Mutiara 4, Jurong, Dr. Fridtjof Nansen, Bawal Putih 2 dan Lemuru*) dengan menggunakan perangkat lunak NAN-SIS untuk data hasil tangkapan, dan FiSAT untuk data frekuensi-panjang. Perangkat-perangkat lunak ini - yang disebarluaskan oleh Food and Agriculture Organization (FAO) dan disimpan oleh para penulisnya-memungkinkan analisis yang lebih rinci terhadap data-dasar tersebut sebagaimana disajikan disini, dan selanjutnya mendorong para pembaca untuk melakukan analisis seperti ini dengan menggunakan tulisan kami ini sebagai contoh dan/atau langkah awal.

Introduction

Large-scale inferences on the structure of demersal fish communities drawn from more than one single survey or a series of surveys performed by the same ship appear to have been exceedingly rare (see Longhurst and Pauly 1987 and contributions in Bianchi 1992). One of the reasons for this, besides simple parochialism, is that an enormous effort is required to encode the results of such surveys (Pauly, this vol.). Indeed, this effort has often stumped the organizers of single surveys (see Box 1 for a historical snippet documenting the encoding problems of the JETINDOFISH survey).

This contribution documents the authors' experience in encoding data from numerous surveys, obtained by five different research vessels over a period of eight years. It also enables replication of the analyses in this volume (Pauly 1993).

Preliminary Data Encoding

The trawl data of *R/V Jurong*, covering areas 3-7 of the JETINDOFISH survey (Lohmeyer, this vol.) and of *R/V Bawal Putih 2*, covering areas 8 and 9 of the same survey (Martosubroto, this vol.), were made available to this project, through the last author, by Mr. Soewito of the Directorate General of Fisheries (DGF), Jakarta, in the form of a magnetic tape in EBCDIC format (density 1,600 bits/inch), produced on 2 November 1985 by the computer center ("Pusat Komputer") of the Department of Agriculture and in printout form (Fig. 1). The tape was read for a preliminary analysis by McManus (1989), and its contents transferred to diskettes. As both the tape and the diskettes subsequently became infected by a fungus, the data on the printout were re-entered by Ms. Luningning Malumay, from September 1992 to September 1994, into a database for trawl data designed by Vakily (1992), using the DataEase software (DataEase 1988).

The species names presented some difficulties, as expected in trawl surveys involving various taxonomists identifying hundreds of species from poorly known waters. More than 70% of the names, current at the time, have since been revised, and many synonymies resulted. The FishBase system

^aICLARM Contribution No. 1317.

^bPresent address: Borneo Marine Research Unit, Universiti Malaysia Sabah, 9th Floor Gaya Centre, Jalan Tun Fuad Stephens, Locked Bag 2073, 88999 Kota Kinabalu, Sabah, Malaysia.

^calso at: Fisheries Centre, the University of British Columbia, 2204 Main Mall, Vancouver, B.C. Canada V6T 1Z4; pauly@fisheries.com

Box 1. JETINDOFISH data processing problem.

[Boks 1. Masalah pemrosesan data JETINDOFISH.]

Modules one and two of the JETINDOFISH project have made several cruises and collected considerable data. It was originally intended that these data would be processed in Rome using a computer program package developed for the FAO Arabian Sea Project.^a This package, modified to meet JETINDOFISH needs, was done as part of the Indian Ocean Project (IOP). The capability no longer exists in Rome to process the data. The process requires expertise knowledgeable of the program and the type of data being processed.

The events have created a problem in getting the data already collected and recorded on forms intended for the Rome program, processed (400 data sheets have been completed). There are several possible solutions to this problem. The simplest solution of course would be to run the data in Rome using the FAO system and computer equipment. The next possible solution is to obtain the computer program from Rome, find a compatible computer in Indonesia and run the data in Indonesia. The advantages of this approach are the closeness to researchers and simplicity in handling data problems. If a compatible computer system is not available in Indonesia, then the next alternative is to "liberate" the FAO program on to another computer system. This is not possible with all computer systems. The last alternative is to create a new system in Indonesia.

The first alternative has the least cost in both time and money, although it is understood that funds are not available in the South China Sea Fisheries Development and Coordinating Programme (SCSP) for computer expenditure. The last is the most expensive in time and money.

There are very few IBM data systems in Jakarta. To place the FAO program in these systems would require computer expertise familiar with the system and computer time both of which would have to be purchased. The capability to do this was not located. The DGF is now using the ICL computer system at the Bureau of Statistics (BPS). This system will not in anyway accept the FAO programs. The BPS staff, however, indicated they could write the necessary program to process the data provided someone else described the functions. BPS would have to be reimbursed for the cost. The problem with this approach would be the necessary expertise to translate the needs for processing the data into a format needed by the BPS staff.

The United States Agency for International Development (USAID) has an ongoing project to computerize the Ministry of Agriculture data. This project will be installing in September a computer for this purpose. In the meantime the project is using a computer that can liberate the FAO program. The project also has computer expertise that can accomplish this task. This project can immediately adopt the FAO program to a local computer, have the survey data key entered, and begin to process the data. The project will also train DGF personnel in the use of the system and will later transfer the data to the Ministry of Agriculture computer system when it is installed.

The USAID project appears to be the least expensive solution for obtaining a computer system and offers early processing of the existing survey data. This will allow changes to be made in the existing program. It is, however, recommended that any form of telecommunications or other sophisticated computer operation not be used.

The USAID project which is under contract to the Iowa State University will need a request from the DGF to initiate this task. This can be with a telephone call to the individual who is in charge of the USAID project.

B.G. Thompson

Consultant

South China Sea Fisheries Development
and Coordinating Programme

February 1980

^aFlowers, J.M. 1978. A data processing and basic analysis system for demersal fisheries surveys. Regional Fishery Survey and Development Project, Bahrain, Iran, Iraq, Kuwait, Oman, Qatar, Saudi Arabia, United Arab Emirates. Field Doc. FI:DP/RAB/71/278/4, 79 p. FAO, Rome, Italy.

was used to update many names and to verify their concordance with known geographic ranges (Froese et al., this vol.). The final dataset was summarized by area-depth combinations and processed as described in McManus (this vol.).

A second dataset was created, based on *Mutiara 4* catch records for the years 1976-1979, by Messrs. M. Badrudin and Suhendro Budihardjo shortly before and during their visit to ICLARM, in May-June 1993. This dataset consisted of catch records from 998 trawl hauls, initially encoded using the Lotus

1-2-3 spreadsheet. The records were then turned into NAN-SIS files for Dr. G. Bianchi (see Bianchi et al., this vol.). This dataset was then expanded to include the survey period November 1974-1976 (522 stations) at IMR (Bergen).

A third dataset documented the *R/V Dr. Fridtjof Nansen* trawl survey conducted in Northwestern Sumatra on 6-30 August 1980 (Aglen et al. 1981 and Bianchi, this vol.). This had been encoded using NAN-SIS, a database system developed by staff of the *Dr. Fridtjof Nansen* Project of the Institute of Marine Research, Bergen, Norway, and distributed by FAO (Strømme 1992). Given our wish to make the result of the surveys documented in this book widely available inside and outside Indonesia (Pauly, this vol.), we decided to use NAN-SIS for storing all catch data.

Further, we opted to use FiSAT (FAO-ICLARM Stock Assessment Tools), a software recently released by FAO (Pauly and Garcia 1994; Gayanilo et al. 1996) to store all length-frequency (L/F) data emanating from the surveys.

Use of NAN-SIS for Catch Data Storage

While now disseminated for use by any group with trawl survey data, NAN-SIS was initially developed to meet the needs of the project which deployed the *R/V Dr. Fridtjof Nansen* (actually the predecessor of the ship currently so named) since 1975, in various parts of the intertropical belt (see Bianchi 1992). The data accumulated were stored in different formats, first as datasheets, then as electronic files for various brands of (micro)computers.

One of the key features of NAN-SIS is its routine for reading files created using various formats (Strømme 1992). We found this routine particularly helpful, given the various datasets created using other systems. This routine is, however, not well-documented by Strømme (see Box 2 for further details).

Because only up to 999 stations at the time can be transferred from another file, the database on Western Indonesian trawl data was split into three subsets:

1. the 958 stations of the surveys of *R/V Jurong* (746 stations), *Bawal Putih 2* (121), *Dr. Fridtjof Nansen* (79) and *Lemuru* (12), and coded JI.
2. the 522 stations of *R/V Mutiara 4* covering the period 1974-1976 entered directly in NAN-SIS at IMR, coded MU, and
3. the 998 stations of *R/V Mutiara 4* (covering the period 1976-1979) coded MM.

C A T C H D E T A I L S				

HAUL NUMBER : 0001				
TOTAL CATCH : 82.60 KG				
FAM / GEN / SPECIE	CATCH(KG)	NUMBER	CATCH RATE (KG/HK)	DENSITY (KG/SQ KM)
CLUPEIDAE				
SARDINELLA				
LONGICEPS	68.00	1989.00	45.33	526.90
NOMEIDAE	3.80	21.00	2.53	18.20
SPHYRNIDAE	1.90	1.00	1.26	9.10
CARANGIDAE				
DECAPTERUS				
MACROSOMA	8.90	42.00	5.93	42.70
TOTAL :			55.06	597.11
TOTAL NO OF SPECIES : 005				
EFFORT (HRS) : 1.50				
AREA TRAWLED (SQ/KM) : 0.208				
HAUL NUMBER : 0002				
TOTAL CATCH : 40.80 KG				
FAM / GEN / SPECIE	CATCH(KG)	NUMBER	CATCH RATE (KG/HK)	DENSITY (KG/SQ KM)
CRARS	5.20	17.00	5.46	17.50
MYCTOPHIDAE	10.60	76.00	7.20	56.40
MACROSOMA	0.80	47.00	0.53	2.70
SCOMBRIDAE				
SCOMBER				
JAPONICUS	1.60	2.00	1.06	5.40
TRICHIURIDAE				
TRICHIURUS				
LEPTURUS	16.70	52.00	11.13	56.40
APOGONIDAE	3.50	120.00	2.53	11.80
POMATOMIDAE	0.50	5.00	0.53	1.60
CARANGIDAE				
DECAPTERUS				

Fig. 1. Facsimile of a typical R/V Jurong catch data sheet.

[Gambar 1. Fax tentang suatu bentuk penyajian data hasil tangkapan kapal penelitian Jurong.]

Fig. 2 provides the rationale for this split.

Encoding of "fresh" data using NAN-SIS, as opposed to transferring already encoded data, was rather straightforward, the only problem, if any, being the need to update the local "species catalog" (i.e., the list of mnemonics; see Box 2) based on the global catalog of the *Dr. Fridtjof Nansen* Project staff (the global catalog required for global comparative analyses was kindly supplied by Dr. Bianchi).

Use of FiSAT for Encoding and Storing Length-Frequency Data

The FiSAT software, created through the merging of ICLARM's Compleat ELEFAN (Pauly 1987; Gayanilo et al. 1988) with FAO's Length-based Fish Stock Assessment (LFSA) (Sparre 1987), and the addition of complementary routines (Pauly and Sparre 1991) was released only in 1995 after thorough debugging.

However, a beta version of FiSAT was available to us, and was used to encode a large fraction of the length-frequency samples, including their weight and ancillary information, (Table 1) collected during the various surveys documented in this volume.^d

FiSAT was used for this rather than NAN-SIS, which also can store length-frequency data because the former:

1. has been taught at numerous stock assessment courses in Indonesia and elsewhere;
2. will be widely available in Indonesia and elsewhere, and most importantly;
3. contains numerous routines for analysis of length-frequency datafiles.

Pauly et al. (this vol.) provide numerous examples of application of FiSAT to the L/F database documented here.

^dThe catch and effort data for Bali Straits lemuru of Ghofar and Mathews (this vol.) were also encoded using FiSAT, as a two column file (on Diskette 2 of the database).

Table 1. Example of a length-frequency file with incorporated length-frequency data on *Decapterus macrosoma* collected during surveys by various research trawlers (here: *R/V Bawal Putih 2* and *R/V Dr. Fridtjof Nansen*), from 15 May 1980 to 13 November 1980.

[Tabel 1. Contoh suatu penyajian frekuensi-panjang *Decapterus macrosoma* yang dikumpulkan selama survei oleh kapal-kapal penelitian trawl (*Bawal Putih 2* dan *Dr. Fridtjof Nansen*) dari 15 Mei 1980 hingga 13 November 1980.]

Midlength ^{a)} (TL, cm)	Date					
	15/05/80 ^{b)}	15/08/80	17/08/80	19/08/80	19/08/80	13/11/80
7.5						1.00
8.5						2.00
9.5						4.00
10.5	18.00					27.00
11.5	36.00					24.00
12.5	36.00					28.00
13.5	43.00					29.00
14.5	51.00					11.00
15.5	50.00					3.00
16.5	27.00			2.00	1.00	2.00
17.5	3.00			7.00	9.00	5.00
18.5	3.00			5.00	4.00	
19.5	2.00		4.00	1.00	1.00	
20.5	1.00					
21.5	1.00					
22.5	1.00	1.00				
23.5		1.00				
24.5		2.00				
25.5		0.00				
26.5		0.00				
27.5		2.00				
28.5		4.00				
29.5		3.00				
30.5		1.00				
31.5		2.00				
Sum	272.00	16.00	4.00	15.00	15.00	136.00
Weight of sample (g)	7900	3400	NA	700	700	4000

^a i.e., the midpoint of the length class interval.

^b Details are provided, in this example, only for the first sample viz: location of station [08° 45.5° South Latitude, 116° 35.0° East Longitude]; Station number: 4; Research Vessel: *Bawal Putih 2*; cod end mesh size: 4 cm; Gear type: Demersal trawl; Bottom depth: 45 meters; Time started trawling: 12:32 hrs; other information. The actual FiSAT file contains similar details on each station with length-frequency data.

However, these applications used only part of the data, and similar contributions can still be extracted from the database through the application of FiSAT (or other software).

Incorporation of Occurrence Records into FishBase

In addition to storing survey catch data into NAN-SIS and L/F data into FiSAT, the project which led to this volume also included strong interaction, at ICLARM, with the FishBase Project. This project is involved in the creation and maintenance of an electronic encyclopedia of fish in the world, available on CD-ROM (Froese and Pauly 1996; Froese et al., this vol.).

The interactions with FishBase were two-way:

1. It was used to check the name and distribution of fish in the database documented here (e.g., McManus, this vol. and Pauly et al., this vol.).

2. It is used as repository of occurrence records of fish sampled during the surveys documented in this volume. This was achieved by selecting, from the NAN-SIS database documented above and from Widodo (1976), occurrence records for all species otherwise known to occur in Indonesia (Froese et al., this vol.), thus documenting the range covered by these surveys, and as far as possible, the range of the fishes in Western Indonesia.

These records were read from NAN-SIS (and from a spreadsheet with Widodo's records) into the "OCCURRENCES" table of FishBase (Froese and Capuli 1996); subsequent information, e.g., on sizes, was then added from the FiSAT files.

Thus, the users of the FishBase 97 will find that it thoroughly covers the fishes of Western Indonesia, for which a special map was created within FishBase.

- the Tropics, Denmark, 1986 and Philippines, 1987. FAO Fish. Rep. (389).
- 001403** Murty, S.V. 1983. Observations on some aspects of biology of silverbelly *Leiognathus bindus* (Valenciennes) from Kakinada. Indian J. Fish. 30(1):61-68.
- 001415** Bouhlel, M. 1985. Stock assessment of the king fish *Scomberomorus commerson* inhabiting the coastal waters of Djibouti Republic and state of fish stocks. Development of fisheries in the areas of the Red Sea and the Gulf of Aden, RAB/83/023/INT/18. Field Doc. 40 p. FAO/UNDP, Rome.
- 001416** Devaraj, M. 1983. Maturity, spawning and fecundity of the king seer, *Scomberomorus commerson*, in the seas around the Indian peninsular. Indian J. Fish. 30:203-230.
- 001418** Palsson, W. A. and T. W. Pietsch. 1989. Revision of the Acanthopterygian fish Family Pegasidae (Order Gasterosteiformes). Indo-Pac. Fishes (18):38 p.
- 001419** Woodland, D.J. 1990. Revision of the fish family Siganidae with descriptions of two new species and comments on distribution and biology. Indo-Pac. Fish. (19):136 p.
- 001435** Beck, U. and A. Sudradjat. 1978. Variations in size and composition of demersal trawl catches from the north coast of Java with estimated growth parameters for three important food fish. Contrib. Demersal Fish. Proj. (Indones.-Ger.) 4:1-80.
- 001439** Conand, F. 1991. Biology and phenology of *Amblygaster sirm* (Clupeidae) in New Caledonia, a sardine of the coral environment. Bull. Mar. Sci. 48:137-149.
- 001442** Sanders, M.J. and S.M. Kedidi. 1984. Stock assessment for the spotted sardinella (*Sardinella sirm*) caught by purse seine adjacent to the border between Egypt and Sudan. Project for Development of Fisheries in Areas of the Red Sea and Gulf of Aden, UNDP/FAO RAB/83/023/04. Cairo. 28 p.
- 001443** Gjøsæter, J., P. Dayaratne, O.A. Bergstad, V. Gjøsæter, M.I. Sousa and I.M. Beck. 1984. Ageing tropical fish by growth rings in the otoliths. FAO Fish. Circ. 176. 54 p.
- 001444** Dayaratne, P. and J. Gjøsæter. 1986. Age and growth of four *Sardinella* species from Sri Lanka. Fish. Res. 4:1-33.
- 001447** Sadhotomo, B. and S.B. Atmadja. 1985. On the growth of some pelagic fishes in the Java Sea. J. Pen. Perikanan Laut (33):53-60. (in Bahasa Indonesia).
- 001449** Corpuz, A., J. Saeger and V. Sambalay. 1985. Population parameters of commercially important fishes in Philippine waters. Tech. Rep. Univ. Philipp. Visayas, Dept. Mar. Fish. (6):99 p.
- 001451** Edwards, R.R.C. 1985. Growth rates of Lutjanidae (snappers) in tropical Australian waters. J. Fish Biol. 26:1-4.
- 001452** Sadhotomo, B. et al. 1983. The dynamics of trevally, *Pentapiron longimanus* at Java Sea. Mar. Fish. Res. Rep. (28):82 p. (In Bahasa Indonesian)
- 001454** Gjøsæter, J. and M.I. Sousa. 1983. Reproduction, age and growth of the Russell's scad *Decapterus russelli* (Rüppell 1829) (Carangidae) from Sofala Bank, Mozambique. Rev. Invest. Pesq. Maputo 8:83-108.
- 001455** Brinca, L., A. Jorde da Silva, L. Sousa, I.M. Sousa and R. Saetre. 1983. A survey of the fish resources at Sofala Bank, Mozambique, September 1982. Reports on surveys with the R/V DR. FRIDTJOF NANSEN. Institute of Marine Research, Bergen. Instituto de Investigacao Pesqueira, Maputo, Mozambique.
- 001462** Boonraksa, V. 1987. Preliminary resource analysis of chub mackerel (*Rastrelliger* spp.) and round scads (*Decapterus* sp.) in the west coast of Thailand. Paper presented at the 3rd Working Group Meeting of the Malacca Strait Project/BOBP, 18-26 August 1986, Phuket, Thailand.
- 001463** Luther, G. 1973. Observations on the biology and the fishery of the Indian mackerel, *Rastrelliger kanagurta* (Cuvier) from Andaman Islands. Indian J. Fish. 20(2):425-447.
- 001464** Sekharan, K.V., M.S. Muthu, G. Sudhakava and B.N. Rao. 1969. Spawning concentrations of the sardine, *Sardinella gibbosa* (Bleeker), off the North Andhra coast in March-April 1969. Indian J. Fish. 16(122):156-160.
- 001465** Sousa, M.I. and M. Gislason. 1985. Reproduction, age and growth of the Indian mackerel *Rastrelliger kanagurta* (Cuvier, 1816) from Sofala Bank, Mozambique. Rev. Invest. Pesq. Maputo (14):1-28.
- 001466** Udupa, K.S. and C.H. Krishna Bhat. 1984. Age and growth equation of the Indian mackerel from purse seine catches off Karnataka coast. Indian J. Fish. 31(1):61-67.
- 001467** Anon. 1985. Report of the Second Working Group Meeting on the Mackerels (*Decapterus* and *Rastrelliger* spp.) in the Malacca Strait, 4-9 October 1985, Colombo, Sri Lanka. Bay of Bengal Programme Document. 23 p.
- 001470** Chullasorn, S., K. Chotiyaputta and R. Chayakul. 1974. Preliminary study of king mackerel (*S. commerson* Lacepède 1802) in the Gulf of Thailand. Annu. Rep. Pelagic Fish. Unit, Mar. Fish. Div. Bangkok 1:331-377. (in Thai). (mimeo)
- 001474** Silvestre, G.T., and M.L. Soriano. 1988. Effect of incorporating sigmoid selection on optimum mesh size estimation for the Samar Sea multispecies trawl fishery. p. 482-492. In S.C. Venema, J.M. Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO/DANIDA Follow-up Training Course on Fish Stock Assessment in the Tropics, Denmark, 1986 and Philippines, 1987. FAO Fish. Rep. (389).
- 001485** Yohannan, T.M. 1979. The growth pattern of Indian mackerel. Indian J. Fish. 26(1/2):207-216.
- 001486** Silvestre, G. 1986. Preliminary analysis of the growth, mortality and yield-per-recruit of ten trawl-caught fish species from the Philippines. Univ. Phil., Coll. Fish., Dept. Mar. Fish. Tech. Rep. (7):1-41.
- 001488** Sanders, M.J. and G.R. Morgan. 1989. Review of the fisheries resources of the Red Sea and Gulf of Aden. FAO Fish. Tech. Rep. (304):138 p.
- 001498** Edwards, R.R.C., A. Bakhader and S. Shaher. 1985. Growth, mortality, age composition and fishery yields of fish from the Gulf of Aden. J. Fish Biol. 27:13-21.
- 001504** Bleeker, P. 1849. A contribution to the knowledge of the ichthyological fauna of Celebes. J. Indian Arch. & E. Asia 3:65-74.
- 001511** Bleeker, P. 1853. Diagnostische beschrijvingen van nieuwe of weinig bekende vischsoorten van Batavia. Tiental I-VI. Natuurkd. Tijdschr. Neder.-Indië 4:451-516.
- 001524** Sanders, M.J., S.M. Kedidi and M.R. Hegazy. 1984. Stock assessment for the horse mackerel (*Trachurus indicus*) caught by purse seine and trawl in the Gulf of Suez. Project for the Development of Fisheries in the areas of the Red Sea and Gulf of Aden, FAO/UNDP RAB/81/002/20. Cairo. 47 p. (mimeo).
- 001529** Sanders, M.J., S.M. Kedidi and M.R. Hegazy. 1984. Stock assessment for the goldstripe sardinella (*Sardinella gibbosa*) caught by small scale fishermen along the Egyptian Red Sea coast. Project for the Development of Fisheries in the areas of the Red Sea and Gulf of Aden, FAO/UNDP RAB/81/002/27. Cairo. 29 p. (mimeo).
- 001531** Sanders, M.J., S.M. Kedidi and M.R. Hegazy. 1984. Stock assessment for the Indian mackerel (*Rastrelliger kanagurta*) caught by purse seine from the Gulf of Suez and more southern Red Sea waters. Project for the Development of Fisheries in the areas of the Red Sea and Gulf of Aden, FAO/UNDP RAB/83/003/03. Cairo. 25 p. (mimeo).
- 001532** Sanders, M.J., S.M. Kedidi and M.R. Hegazy. 1984. Stock assessment for the brushtooth lizardfish (*Saurida undosquamis*) caught by trawl in the Gulf of Suez. Project for the Development of Fisheries in the areas of the Red Sea and Gulf of Aden, FAO/UNDP RAB/83/023/05. Cairo. 28 p. (mimeo).
- 001539** Cantor, T.E. 1849. Catalogue of Malayan fishes. J. R. Asiatic Soc. Bengal 18(2):981-1443.
- 001555** Evermann, B.W. and A. Seale. 1907. Fishes of the Philippine Islands. Bull. U.S. Bur. Fish. 26(1906):49-110.
- 001601** Munro, I.S.R. 1958. The fishes of the New Guinea region. Papua New Guinea Agric. J. (1956) 10(4):97-369.
- 001602** Myers, R.F. 1991. Micronesian reef fishes. Second Ed. Coral Graphics, Barrigada, Guam. 298 p.
- 001617** Scott, J. S. 1959. An introduction to the sea fishes of Malaya. Min. of Agriculture. Government Press, Kuala Lumpur, Malaysia. 180 p.
- 001632** Suvatti, C. 1981. Fishes of Thailand. Royal Institute of Thailand, Bangkok. 379 p.
- 001633** Taylor, W. R. 1964. Fishes of Arnhem Land. Rec. Amer.-Aust. Exped. Arnhem Land 4:454-307.
- 001652** Linnaeus, C. 1758. Systema Naturae per Regna Tria Naturae secundum Classes, Ordines, Genera, Species cum Characteribus, Differentiis Synonymis, Locis. 10th ed., Vol. 1. Holmiae Salvii. 824 p.

- 001687 Cuvier, G. L. 1816. Le Règne Animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Les reptiles, les poissons, les mollusques et les annélides. Édition 1. v. 2. 532 p.
- 001724 Carcasson, R.H. 1977. A field guide to the coral reef fishes of the Indian and West Pacific Oceans. William Collins Sons & Co. Ltd, Glasgow. 320 p.
- 001751 Banerji, S.K. and T.S. Krishnan. 1973. Acceleration of assessment of fish populations and comparative studies of similar taxonomic groups. p. 158-175. *In* Proceedings of the Symposium on Living Resources of the Seas Around India. Spec. Publ., Centr. Mar. Biol. Res. Inst., Cochin, India. 748 p.
- 001809 Anon. 1986. Surveys of the Fish Resources of Angola. January 1985 - June 1986. Data Files. Reports on Surveys with the R/V Dr. Fridtjof Nansen. NORAD - FAO/UNDP GL082/001. pag. var.
- 001830 Eschmeyer, W.N. 1990. Catalog of the genera of recent fishes. California Academy of Sciences, San Francisco, USA. 697 p.
- 001836 Pauly, D. 1979. Gill size and temperature as governing factors in fish growth: a generalization of von Bertalanffy's growth formula. *Berichte des Instituts für Meereskunde an der Christian-Albrechts Universität Kiel*, (63). 156 p.
- 001911 Pauly, D. 1980. A selection of simple methods for the assessment of tropical fish stocks. *FAO Fish. Circ.* (729):54 p.
- 001918 Cuvier, G. L. 1829. Le règne animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Nouvelle édition. Paris, 2:122-406.
- 001920 Randall, J. E. 1956. A revision of the surgeonfish genus *Acanthurus*. *Pac. Sci.* 10(2):159-235.
- 001921 Randall, J.E. 1987. Three nomenclatorial changes in Indo-Pacific surgeonfishes (Acanthurinae). *Pac. Sci.* 41(1-4):54-61.
- 001966 Chan, E.H. and H.C. Liew. 1986. A study on tropical demersal species (Malaysia). Report to the International Development Research Centre: IDRC/3-A-83-1905, Singapore, 64 p.
- 001975 Morgan, G.R. and D. Pauly. 1987. Analysis of length-frequency data: some available programs and user's instructions. p. 373-462. *In* D. Pauly and G.R. Morgan (eds.) Length-based methods in fisheries research. ICLARM Conference Proceedings 13, 468 p.
- 002021 Magnusson, J. 1970. Deep sea fishing development. The Philippine marine fisheries biology. FAO-UNDP report to the Philippine Government. FI: SF/Phil 11. 84 p.
- 002023 Tiews, K., I.A. Ronquillo and L.M. Santos. 1971. On the biology of anchovies (*Stolephorus Lacépède*) in Philippine waters. *Philipp. J. Fish.* 9(1/2):92-123.
- 002029 Armada, N. and G. Silvestre. 1981. Demersal fish resource survey in Samar Sea and Carigara Bay. UP-NSDB Project 7811.1c Ag. 56 p. College of Fisheries, University of the Philippines in the Visayas, Iloilo City, Philippines.
- 002030 Dawson, C.E. 1966. *Gunterichthys longipenis*, a new genus and species of ophidioid fish from the northern Gulf of Mexico. *Proc. Biol. Soc. Wash.* 79:205-214.
- 002044 Dwiponggo, A. and M. Badrudin. 1979. Data of trawl survey by R.V. Mutiara-IV. Contribution of the Demersal Fisheries. Project No. 6A. Marine Fisheries Research Institute (LPPL), Jakarta. 128 p.
- 002045 Martosubroto, P. and D. Pauly. 1976. R/V Mutiara-IV survey data. November 1974 to July 1976. Marine Fisheries Research Institute. Contributions of the Demersal Fisheries Project No. 3, Jakarta. 136 p.
- 002047 Dwiponggo, A. And M. Badrudin. 1980. Length-frequency measurements of demersal fish. Contribution of the Demersal Fisheries Project, RIMF Spec. Rep. (7b), 94 p.
- 002088 Pauly, D. and N. David. 1981. ELEFAN I, a BASIC program for the objective extraction of growth parameters from length-frequency data. *Meeresforsch.* 28(4):205-211.
- 002089 Pauly, D. 1977. The Leiognathidae (Teleostei): their species, stocks and fisheries in Indonesia, with notes on the biology of *Leiognathus splendens* (Cuvier). *Penelitian Laut Indones./Mar. Res. Indones.* 19:73-93.
- 002107 Menon, A.G.K. and S. Monkolprasit. 1974. Cynoglossidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO. Vol. II, pag. var.
- 002108 Kühlmorgen-Hille, G. 1974. Leiognathidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO. Vol. II, pag. var.
- 002109 Bathia, U. and T. Wongratana. 1974. Mugilidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (Fishing Area 71). Volume 3. [var. pag.]. FAO, Rome.
- 002110 Kühlmorgen-Hille, G. 1974. Mullidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO. Vol. III, pag. var.
- 002112 Chan, W., F. Talbot and S. Sontirat. 1974. Pomadasyidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). Volume 3. FAO, Rome.
- 002113 Chan, W., U. Bathia and D. Carlsson. 1974. Sciaenidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). Volume 3. FAO, Rome.
- 002115 Eggleston, D. 1974. Sparidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71), Volume 4. FAO, Rome.
- 002116 Abe, T. 1974. Sphyracidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71), volume 4. [var. pag.] FAO, Rome.
- 002117 Abe, T. and D. Pathansali. 1974. Synodontidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). volume 4. [var. pag.]. FAO, Rome.
- 002135 Roux, C. 1986. Pomadasyidae. p. 327-330. *In* J. Daget, J.-P. Gosse and D.F.E. Thys van den Audenaerde (eds.) Check-list of the freshwater fishes of Africa (CLOFFA). ISNB, Brussels; MRAC, Tervuren; and ORSTOM, Paris. Vol. 2
- 002136 Randall, J.E. 1980. Two new Indo-Pacific labrid fishes of the genus *Halichoeres*, with notes on other species of the genus. *Pac. Sci.* 34(4):415-432.
- 002137 Randall, J.E. 1981. Revision of the labrid fish genus *Labropsis* with description of five new species. *Micronesica* 17(1-2):125-155.
- 002138 Randall, J.E. 1983. Revision of the Indo-Pacific labrid fish genus *Wetmorella*. *Copeia* (4):875-883.
- 002139 Randall, J.E. and R. Lubbock. 1981. Labrid fishes of the genus *Paracheilinus*, with descriptions of three new species from the Philippines. *Japan. J. Ichthyol.* 28(1):19-30.
- 002142 Randall, J.E., T.H. Fraser and E.A. Lachner. 1990. On the validity of the Indo-Pacific cardinalfishes *Apogon aureus* (Lacepède) and *A. fleurieu* (Lacepède), with description of a related new species from the Red Sea. *Proc. Biol. Soc. Wash.* 103(1):39-62.
- 002178 Dalzell, P. and R.A. Ganaden. 1987. A review of the fisheries for small pelagic fishes in Philippine waters. *Tech. Pap. Ser. Bur. Fish. Aquat. Resour. (Philipp.)* 10(1):58 p. Bureau of Fisheries and Aquatic Resources, Quezon City, Philippines.
- 002221 Bianchi et al. 1989. Gabon - Angola Survey of the R/V Dr. Fridtjof Nansen. NAN-SIS database.
- 002290 Loubens, G. 1980. Biologie de quelques espèces de poissons du lagon Néo-Calédonien. III. Croissance. *Cah. Indo-Pac.* 2:101-153.
- 002295 Carpenter, K.E. and G.R. Allen. 1989. FAO Species Catalogue. Vol. 9. Emperor fishes and large-eye breams of the world (family Lethrinidae). An annotated and illustrated catalogue of lethinid species known to date. FAO Species Synop. No. 125(9):118 p.
- 002300 Ralston, S. and H.A. Williams. 1988. Depth distributions, growth, and mortality of deep slope fishes from the Mariana Archipelago. NOAA Technical Memorandum NMFS. NOAA-TM-NMFS-SWFC-113. 47 p.
- 002302 Hughes, G.M. and M. Morgan. 1973. The structure of fish gills in relation to their respiratory function. *Biol. Rev.* 48:419-475, and

- Supplementary Publication SUP 90005, British Library, Lending Division, Boston Spa, Wetherby, Yorkshire, LS23 7BQ.
- 002308** Gray, I.E. 1954. Comparative study of the gill area of marine fishes. *Biol. Bull. Mar. Biol. Lab. Woods Hole* 107:219-225.
- 002311** Hughes, G.M. 1966. The dimensions of fish gills in relation to their function. *J. Exp. Biol.* 45:177-195.
- 002325** Munz, F.W. and W.N. McFarland. 1973. The significance of spectral position in the rhodopsins of tropical marine fishes. *Vision Res.* 13:1829-1874.
- 002334** Randall, J.E., G.R. Allen and R.C. Steene. 1990. *Fishes of the Great Barrier Reef and Coral Sea*. University of Hawaii Press, Honolulu, Hawaii. 506 p.
- 002462** Day, F. 1889. *Fishes*. p. 1-509. In W.T. Blanford (ed.) *The fauna of British India, including Ceylon and Burma, Volume 2:i-xiv*. Taylor and Francis, London.
- 002504** Peñaflor, G.C. 1988. Growth of *Leiognathus splendens* based on daily otolith rings and length-frequency analysis. *Asian Fish. Sci.* 2:83-92.
- 002505** Donaldo, S.S. 1979. Contributions to the biology of common slipmouth, *Leiognathus splendens* (Cuvier, 1829) caught from Manila Bay. University of the Philippines, Diliman, Quezon City. 55 p. M.Sc. thesis.
- 002677** Randall, J.E. 1972. A revision of the labrid fish genus *Anampses*. *Micronesica* 8(1-2):151-190.
- 002682** Kuronuma, K. 1961. A check list of fishes of Vietnam. United States Consultants, Inc.; International Cooperation Administration Contract - IV-153. Division of Agriculture and Natural Resources, United States Operations Mission to Vietnam. 66 p.
- 002683** Schneider, W. 1990. FAO species identification sheets for fishery purposes. Field guide to the commercial marine resources of the Gulf of Guinea. Prepared and published with the support of the FAO Regional Office for Africa. FAO, Rome. 268 p.
- 002688** Hensley, D. A. and K. Amaoka. 1989. A redescription of *Pseudorhombus megalops*, with comments on *Cephalopsetta ventrocellata* (Osteichthyes: Pleuronectiformes: Paralichthyidae). *Proc. Biol. Soc. Wash.* 102(3):577-585.
- 002689** Randall, J.E. and J.H. Choat. 1980. Two new parrotfishes of the genus *Scarus* from the Central and South Pacific, with further examples of sexual dichromatism. *Zool. J. Linn. Soc.* 70:383-419.
- 002745** Randall, J. E. and S. Shen. 1978. A review of the labrid fishes of the genus *Cirrhitilabrus* from Taiwan, with description of a new species. *Bull. Inst. Zool., Acad. Sinica* 17(1):13-24.
- 002746** Randall, H.A. and G.R. Allen. 1977. A revision of the damselfish genus *Dascyllus* (Pomacentridae) with description of a new species. *Rec. Aust. Mus.* 31(9):349-385.
- 002747** Randall, J.E. and V.C. Springer. 1973. The monotypic Indo-Pacific labrid fish genera *Labrichthys* and *Diproctacanthus* with description of a new related genus, *Larabicus*. *Proc. Biol. Soc. Wash.* 86(23):279-298.
- 002748** Randall, J.E. and R.H. Kuitert. 1982. Three new labrid fishes of the genus *Coris* from the Western Pacific. *Pac. Sci.* 36(2):159-173.
- 002795** Hureau, J.C. 1979. Mullidae. p. 402-404. In J.C. Hureau and Th. Monod (eds.) *Check-list of the fishes of the north-eastern Atlantic and of the Mediterranean (CLOFNAM)*. UNESCO, Paris. Vol. 1.
- 002799** Smith, M.M. and R.J. McKay. 1986. Haemulidae. p. 564-571. In M.M. Smith and P.C. Heemstra (eds.) *Smiths' sea fishes*. Springer-Verlag, Berlin.
- 002830** Thomson, J.M. 1984. Mugilidae. In W. Fischer and G. Bianchi (eds.) *FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51)*. volume 3. [pag. var.]. FAO, Rome.
- 002857** Herre, A.W.C.T. and A.F. Umali. 1948. English and local common names of Philippine fishes. U. S. Dept. of Interior and Fish and Wildl. Serv. Circular No. 14, U. S. Gov't Printing Office, Washington. 128 p.
- 002858** Conlu, P.V. 1986. Guide to Philippine flora and fauna. Fishes. Volume IX. Natural Resources Management Center, Quezon City.
- 002860** Valenciennes, A. 1847. Poissons. Catalogue des principales espèces de poissons, rapportées de l'Amérique méridionale. In A. d'Orbigny. *Voyage dans l'Amérique méridionale*.
- 002871** Bianchi, G. 1985. FAO species identification sheets for fishery purposes. Field guide to the commercial marine and brackish-water species of Tanzania. Prepared and published with the support of TCP/URT/4406 and FAO (FIRM) Regular Programme. FAO, Rome. 199 p.
- 002872** Bianchi, G. 1985. FAO species identification sheets for fishery purposes. Field guide to the commercial marine and brackish-water species of Pakistan. Prepared with the support of PAK/77/033/ and FAO (FIRM) Regular Programme. FAO, Rome. 200 p.
- 002877** Boonwanich, T. 1991. Population dynamics of *Saurida elongata* and *S. undosquamis* (Synodontidae) in the southern Gulf of Thailand. *Fishbyte* 9(1): 23-27.
- 002926** Sambily, V.C. 1991. Depth-distribution patterns of demersal fishes of the Samar Sea, Philippines, and their use for estimation of mortality. M.Sc. thesis, University of the Philippines in the Visayas. 66 p.
- 002948** Whitehead, P.J.P. 1974. Clupeidae. In W. Fischer and P.J.P. Whitehead (eds.) *FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71)*. volume 1. [pag. var.]. FAO, Rome.
- 003079** Randall, J.E., D. Golani, and A. Diamant. 1988. *Sargocentron marisrubri*, a new squirrelfish (Bericyformes: Holocentridae) from the Red Sea. *Isr. J. Zool.* 35:187-197.
- 003084** Ralston, S. 1988. Length-weight regressions and condition indices of lutjanids and other deep slope fishes from the Mariana Archipelago. *Micronesica*, 21: 189-197.
- 003090** Manooch, C.S. 1987. Age and growth of snappers and groupers. p. 329-373. In J.J. Polovina and S. Ralston (eds.) *Tropical snappers and groupers: biology and fisheries management*. Ocean Resour. Mar. Policy Ser. Westview Press, Inc., Boulder and London.
- 003109** Hutchins, B. 1984. Balistidae. In W. Fischer and G. Bianchi (eds.) *FAO species identification sheets for fishery purposes. Western Indian Ocean Fishing area 51*. Prepared and printed with the support of the Danish International Development Agency (DANIDA). FAO, Rome. Vol. 1.
- 003111** Munro, J.L. 1983. Epilogue: Progress in coral reef fisheries research, 1973-1982. p. 249-265. In J.L. Munro (ed.) *Caribbean coral reef fishery resources. ICLARM Studies and Reviews* 7.
- 003128** Abe, T. 1974. Balistidae. In W. Fischer and P.J.P. Whitehead (eds.) *FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71)*. volume 1. FAO, Rome.
- 003131** Sainsbury, K.J., P.J. Kailola and G.G. Leyland. 1985. Continental shelf fishes of the northern and north-western Australia. CSIRO Division of Fisheries Research; Clouston & Hall and Peter Pownall Fisheries Information Service, Canberra, Australia. 375 p.
- 003132** Allen, G.R. and R. Swainston. 1988. *The marine fishes of north-western Australia: a field guide for anglers and divers*. Western Australian Museum, Perth.
- 003136** Manickchand-Heileman, S. and J. Flüs. 1990. Species composition and seasonality of coastal demersal fish stock in Trinidad, West Indies. *Caribb. Mar. Stud.* 1:1-21.
- 003145** Randall, J.E. 1986. Acanthuridae. p. 811-823. In M.M. Smith and P.C. Heemstra (eds.) *Smiths' sea fishes*. Springer-Verlag, Berlin.
- 003151** Rofen, R.R. 1963. *Handbook of the food fishes of the Gulf of Thailand*. Compiled and edited by the George Vanderbilt Foundation and the University of California Scripps Institution of Oceanography, La Jolla, California. 736 p.
- 003197** Smith-Vaniz, W.F. 1986. Carangidae. p. 638-661. In M.M. Smith and P.C. Heemstra (eds.) *Smiths' sea fishes*. Springer-Verlag, Berlin.
- 003207** Russell, B.C. 1991. Description of a new species of *Nemipterus* (Pisces: Perciformes; Nemipteridae) from the western Pacific, with re-descriptions of *Nemipterus marginatus* (Valenciennes), *N. mesoprion* (Bleeker) and *N. nematopus* (Bleeker). *J. Nat. History*, 25:1379-1389.
- 003225** Brinca, L., V. Mascarenhas, B. Palha de Sousa, L. Palha de Sousa, I.M. Sousa, R. Saetre and I. Timochin. 1984. A survey on the fish resources at Sofala Bank- Mozambique, May-June 1983. Reports on Surveys with the R/V Dr. Fridtjof Nansen. Instituto de Investigacao Pesqueira, Maputo, Mozambique.
- 003242** Wenner, C.A. 1978. Anguillidae. In W. Fischer (ed.) *FAO species identification sheets for fishery purposes. West Atlantic (Fishing Area 31)*. volume 1. [pag. var.]. FAO, Rome.
- 003243** Armitage, R.O., D.A. Payne, G.J. Lockley, H.M. Currie, R.L. Colban,

- B.G. Lamb and L.J. Paul, Editors. 1981. Guide book to New Zealand commercial fish species. New Zealand Fishing Industry Board, Wellington, New Zealand.
- 003257** Castle, P.H.J. and J.E. McCosker. 1986. Muraenidae. p. 165-176. *In* M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 003263** Compagno, L.J.V. 1986. Dasyatidae. p. 135-142. *In* M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 003268** Ritterbush, S. 1975. An assessment of the population biology of the Bali Strait lemuru fishery. *Mar. Fish. Res. Rep.* (1):1-38.
- 003277** Berry, F.H. and W.F. Smith-Vaniz. 1978. Carangidae. *In* W. Fischer (ed.) FAO species identification sheets for fishery purposes. West Atlantic (Fishing Area 31). volume 1. FAO, Rome. [var. pag.]
- 003279** Wongratana, T. and U. Bathia. 1974. Ariidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (Fishing Area 71). volume 1. [pag. var.]. FAO, Rome.
- 003280** Chan, W., F. Talbot, and P. Sukhavisidh. 1974. Carangidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). Volume 1. FAO, Rome.
- 003281** Sukhavisidh, P. and D. Eggleston. 1974. Centropomidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). volume 1. [pag. var.]. FAO, Rome.
- 003282** Langham, N. 1974. Coryphaenidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO, Vol. 2, pag. var.
- 003287** Smith-Vaniz, W.F. 1984. Carangidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 1. [pag. var.]. FAO, Rome.
- 003290** Jayaram, K.C. 1984. Ariidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 1. FAO, Rome. pag. var.
- 003383** Organisation for Economic Co-operation and Development. 1990. Multilingual dictionary of fish and fish products. Fishing News Books, Oxford.
- 003391** Menon, A.G.K. 1984. Cynoglossidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 2.
- 003392** Poss, S.G. 1984. Dactylopteridae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 2.
- 003397** Bauchot, M.-L. 1987. Poissons osseux. p. 891-1421. *In* W. Fischer, M.L. Bauchot and M. Schneider (eds.) Fiches FAO d'identification pour les besoins de la pêche. (rev. 1). Méditerranée et mer Noire. Zone de pêche 37. Vol. II. Commission des Communautés Européennes and FAO, Rome.
- 003399** Kühlmorgen-Hille, G. 1974. Gerreidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO, Volume 2, pag. var.
- 003404** Wongratana, T. 1974. Lactariidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (Fishing Area 71). volume 2. [pag. var.]. FAO, Rome.
- 003408** Menon, A.G.K. 1974. Polynemidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71). Rome, FAO, Volume 3, pag. var.
- 003409** Woodland, D.J. 1984. Gerreidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 2. [pag. var.]. FAO, Rome.
- 003412** McKay, R.J. 1984. Haemulidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 2. FAO, Rome. pag. var.
- 003414** Eggleston, D. 1974. Priacanthidae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71), Volume 3. FAO, Rome.
- 003416** Carlsson, D. 1974. Rachycentridae. *In* W. Fischer and P.J.P. Whitehead (eds.) FAO species identification sheets for fishery purposes. Eastern Indian Ocean (Fishing Area 57) and Western Central Pacific (Fishing Area 71). volume 3. [var. pag.] FAO, Rome.
- 003424** James, P.S.B.R. 1984. Leiognathidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 2. FAO, Rome. pag. var.
- 003430** Munro, I.S.R. 1960. Handbook of Australian Fishes No. 33. Reprinted (Leiognathidae) from *Fish. News Lett. Aust.* 19(6):17-20 by Publicity Press Pty. Ltd., 71-75 Regent St., Sydney.
- 003436** Canagaratnam, P. and J.C. Medcof. 1956. Ceylon's beach seine fishery. *Bull. Fish. Res. Strn., Ceylon* 4(4):1-32.
- 003437** Central Fisheries Dept. Pakistan. 1955. Marine fishes of Karachi and the coasts of Sind and Makran. Govt. of Pakistan Press, Karachi.
- 003438** Chua, T.E. and H.C. Lai. 1978. Fishes. p. 49-87. *In* T.E. Chua and J.A. Mathias (eds.) Coastal resources of West Sabah. Penerbit Universiti Sains Malaysia, Pulau Pinang.
- 003439** Fowler, H.W. 1936. A synopsis of the fishes of China. Part 6. The mackerels and related fishes. Family Carangidae, continued. *Hong Kong Nat.* 7(1):61-80. Reprint edition (1972) Vol. 1. Antiquariat Junk, Netherlands.
- 003440** Günther, A. 1860. Catalogue of the Acanthopterygian fishes in the collection of the British Museum. British Museum, London, Vol. II.
- 003441** Macleay, W. 1884. Supplement to descriptive catalogue of the fishes of Australia. *Proc. Linn. Soc. N.S.W.* 9:2-64.
- 003442** Misra, K.S. 1959. An aid to commercial fishes. *Rec. Indian Mus.* 57:255-259.
- 003443** Jayaram, K.C. 1981. The freshwater fishes of India, Pakistan, Bangladesh, Burma and Sri Lanka - a handbook. Zoological Survey of India, Calcutta. 475 p. + plates.
- 003444** Hardenberg, J.D.F. 1936. On a collection of fishes from the estuary and lower and middle course of the river Kapuas (W. Borneo). *Treubia* 15(3):225-254.
- 003470** Kumaran, M. and J.E. Randall. 1984. Mullidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 3.
- 003476** Knapp, L.W. 1984. Platycephalidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 3. FAO, Rome. pag. var.
- 003479** Menon, A.G.K. and M. Babun Rao. 1984. Polynemidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 3. FAO, Rome. pag. var.
- 003490** Lal Mohan, R.S. 1984. Sciaenidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 4. FAO, Rome. pag. var.
- 003503** Poss, G.S. and K.V. Rama Rao. 1984. Scorpaenidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 4. FAO, Rome. pag. var.
- 003505** Menon, A.G.K. 1984. Soleidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 4. FAO, Rome. pag. var.
- 003507** Bauchot, M.-L. and M.M. Smith. 1984. Sparidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). volume 4. [var. pag.] FAO, Rome.
- 003509** Poll, M., G.G. Teugels and P.J.P. Whitehead. 1986. Clupeidae. p. 41-55. *In* J. Daget, J.-P. Gosse and D.F.E. Thys van den Audenaerde (eds.) Check-list of the freshwater fishes of Africa (CLOFFA). ORSTOM, Paris and MRAC, Tervuren. Vol. 1.
- 003517** Haedrich, R.L. 1984. Stromateidae. *In* W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 4. FAO, Rome. pag. var.

- 003520 Cressey, R. and R.S. Waples. 1984. Synodontidae. In W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Volume 4. FAO, Rome.
- 003539 Vari, R. 1984. Teraponidae. In W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 4. [pag. var.]. FAO, Rome.
- 003555 Ronquillo, I.A. 1975. A review of the roundscad fishery in the Philippines. Philipp. J. Fish. 1(1 & 2):86-126.
- 003556 Magnusson, J. 1970. Report on assignment as Marine Fisheries Biologist with the UNDP (SF)/FAO Deep Sea Fishing Project in the Philippines. (January, 1966- June, 1969). FAO Report. 86 p. (mimeo).
- 003557 Aprieto, V.L. and E.P. Viloso. 1979. Catch composition and relative abundance of trawl-caught fishes in the Visayan Sea. Fish. Res. J. Philipp. 4(1):9-18.
- 003560 Sekharan, K.V. 1968. Growth rate of the sardines, *Sardinella albelli* (Val.) and *S. gibbosa* (Bleek.), in the Mandapam area. Indian J. Fish. 15(1 & 2):68-80.
- 003579 Menon, M. and N. Radhakrishnan. 1974. Present status of knowledge regarding the biology of Indian mackerel *Rastrelliger kanagurta* (Cuvier). 15th IPFC Proc., Wellington, New Zealand, 18-27 October 1972. Sect. III:343-350.
- 003590 Merrett, N.R. 1990. Chlorophthalmidae. p. 351-360. In J.C. Quero, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.) Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. Vol. 1.
- 003605 Schroeder, R.E. 1982. Length-weight relationships of fishes from Honda Bay, Palawan, Philippines. Fish. Res. J. Philipp. 7(2):50-53.
- 003607 Jones, G. 1985. Revision of the Australian species of the fish family Leiognathidae. Aust. J. Mar. Freshwat. Res. 36:559-613.
- 003614 Jayabalan, N. 1988. Age and growth of the ponyfish, *Leiognathus splendens* (Cuvier) caught off Porto Novo coast. p. 259-261. In M. Mohan Joseph (ed.) The First Indian Fisheries Forum, Proceedings. Asian Fisheries Society, Indian Branch, Mangalore.
- 003621 Guanco, M.R. 1991. Growth and mortality of Indian mackerel *Rastrelliger kanagurta* (Scombridae) in the Visayas Sea, Central Philippines. Fishbyte 9(2):13-15.
- 003624 Majid, A. and A. Imad. 1991. Growth of *Pomadasys kaakan* (Haemulidae) off the coast of Pakistan. Fishbyte 9(2):19-20.
- 003626 Edwards, R.R.C. and S. Shafer. 1991. The biometrics of marine fishes from the Gulf of Aden. Fishbyte 9(2):27-29.
- 003627 Mathews, C.P. and M. Samuel. 1991. Growth, mortality and length-weight parameters for some Kuwaiti fish and shrimp. Fishbyte 9(2):30-33.
- 003641 Bawazeer, A.S. 1987. The fishery management of the stock of chim, the giant sea catfish *Arius thalassinus* in Kuwait waters. Kuwait Bull. Mar. Sci. 9:87-100.
- 003642 Brothers, E.B. and C.P. Mathews. 1987. Application of otolith microstructural studies to age determination of some commercially valuable fish of the Arabian Gulf. Kuwait Bull. Mar. Sci. 9:127-157.
- 003649 Arora, H.L. 1952. A contribution to the biology of the silver belly, *Leiognathus splendens* (Cuv.). Indo-Pac. Fish. Counc. Tech. Pap. 4:75-80.
- 003653 James, P.S.B.R. and M. Badrudeen. 1986. Studies on the maturation and spawning of the fishes of the family Leiognathidae from the seas around India. Indian J. Fish. 33(1):1-26.
- 003655 Rao, K.S. 1967. Reproductive cycles and lipid levels in *Leiognathus splendens* (Cuvier). J. Mar. Biol. Ass. India 9(2):303-322.
- 003667 Mangalik, A. 1965. Makanan dan tabiat makan dari dua jenis 'ikan peperek', *Leiognathus splendens* (Cuvier), dan *Gazza minuta* (Bleeker). Thesis, Institut Pertanian Bogor, Indonesia. 30 p. (in Bahasa Indonesia).
- 003669 Torres, F. Jr. 1991. Tabular data on marine fishes from Southern Africa, Part II: Growth parameters. Fishbyte 9(2):37-38.
- 003670 van der Elst, R. 1981. A guide to the common sea fishes of Southern Africa. C. Struik, Cape Town. 367 p.
- 003674 Boonwanich, T. and S. Amornchairojkul. 1982. Length composition of *Saurida undosquamis* and *S. elongata* in the western coast of the Gulf of Thailand. Marine Fisheries Division, Department of Fisheries Publ. No. 13, 8 p. Bangkok, Thailand.
- 003675 Sommani, P. 1989. Estimation of the Bertalanffy growth parameters of fish in the Gulf of Thailand by unweighted and weighted least square methods. SEAFDEC/TD/RES/21 Thailand. 37 p.
- 003676 Siripakhavanich, S. 1990. A study on population dynamics of the brushtooth lizard fish, *Saurida undosquamis* (Richardson) off the upper western coast in the Gulf of Thailand. Faculty of Fisheries, Kasetsart University, Thailand. 77 p.
- 003678 Torres, F. Jr. 1991. Tabular data on marine fishes from Southern Africa, Part I. Length-weight relationships. Fishbyte 9(1):50-53.
- 003686 Nielsen, J.G. 1990. Ophidiidae. p. 564-573. In J.C. Quero, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.) Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. Vol. 2.
- 003784 Morales-Nin, B. 1988. Caution in the use of daily increments for ageing tropical fishes. Fishbyte 6(2):5-6.
- 003785 Dalzell, P. 1988. Small pelagic fisheries investigations in the Philippines. Part II. The current status. Fishbyte 6(3):2-4.
- 003786 Philbrick, C. 1988. Length frequency analysis of pelagic fish species. Fishbyte 6(3):5-6.
- 003804 Dalzell, P., S.R. Lindsay, and H. Patiale. 1991. Fisheries resources survey of the Island of Niue. Tech. Doc. Inshore Fish. Res. Proj. S. Pac. Comm 3. A report prepared in conjunction with the South Pacific Commission Inshore Fisheries Research Project, and the FAO South Pacific Aquaculture Development Project for the Government of Niue, July 1990.
- 003807 Smith, A. and P. Dalzell. 1993. Fisheries resources and management investigations in Woleai Atoll, Yap State, Federated States of Micronesia. Inshore Fish. Res. Proj., Tech. Doc., South Pacific Commission. Noumea, New Caledonia. 64 p.
- 003810 Russell, B.C. 1990. Nemipterid fishes of the world. (Threadfin breams, whiptail breams, monocle breams, dwarf monocle breams, and coral breams). Family Nemipteridae. An annotated and illustrated catalogue of nemipterid species known to date. FAO Fisheries Synops. 12(125):149 p.
- 003972 McCosker, J.E. and P.H.J. Castle. 1986. Ophichthidae. p. 176-186. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004055 Cressey, R.F. 1986. Synodontidae. p. 270-273. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004066 Hulley, P.A. 1986. Myctophidae. p. 282-321. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004104 Olney, J.E. and D.F. Markle. 1986. Carapidae. p. 350-354. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004181 Heemstra, P.C. 1986. Trachichthyidae. p. 410-413. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004201 Randall, J.E. and P.C. Heemstra. 1986. Holocentridae. p. 415-427. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004233 Smith-Vaniz, W.F. 1986. Carangidae. p. 815-844. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen and E. Tortonese (eds.) Fishes of the north-eastern Atlantic and the Mediterranean. UNESCO, Paris. vol. 2.
- 004241 Ebeling, A.W. 1986. Melamphidae. p. 427-432. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004313 Eschmeyer, W.N. 1986. Scorpaenidae. p. 463-478. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004318 Heemstra, P.C. and T.J. Martin. 1986. Ambassidae. p. 507-508. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004319 Heemstra, P.C. and J.E. Randall. 1986. Serranidae. p. 509-537. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004327 Heemstra, P.C. 1986. Teraponidae. p. 543-544. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 004330 Makwaia, E.D.S. and L.B. Nhwani. 1992. Population parameters of *Sardinella* species in the coastal waters of Dar es Salaam, Tanzania. NAGA, The ICLARM Q. 15(1):25-28.
- 004331 Nhwani, L.B. and G. Bianchi. 1987. Preliminary assessment of

- 006323** Crawford, R. 1993. World record game fishes 1993. The International Game Fish Association, Pompano Beach, Florida.
- 006328** Walford, L. and R. Wicklund. 1973. Contribution to a world-wide inventory of exotic marine and anadromous organisms. FAO Fish. Tech. Pap. No. 121. 49 p.
- 006365** Fouda, M.M. and G.V. Hermosa, Jr. 1993. A checklist of Oman fishes. Sultan Qaboos University Press, Sultanate of Oman. 42 p.
- 006425** Lijam, N. 1999. On the accuracy of natural mortality of fish stocks estimated from water temperature and growth parameters. Dissertation, Univ. North Wales, Bangor, U.K. 81 p.
- 006490** Shiokawa, T. 1988. Managements of ribbon fish resources in the Central Japan Sea. Japan Fisheries Resource Conservation Association, Tokyo. 102 p.
- 006543** Karrer, C. 1990. Zeniidae. p. 629-630. In J.C. Quero, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.) Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. Vol. 2.
- 006545** Karrer, C. 1990. Oreosomatidae. p. 637-640. In J.C. Quero, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.) Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. Vol. 2.
- 006567** Aglen, A., L. Føyn, O.R. Godø S. Myklevoll and O.J. Østvedt. 1981. A survey of the marine fish resources of the north and west coast of Sumatra, August 1980. Reports on Surveys with the R/V 'Dr. Fridtjof Nansen', Institute of Marine Research. Bergen. 55 p.
- 006660** Tyler, J.C. 1986. Triacanthodidae. p. 887-890. In M.M. Smith and P.C. Heemstra (eds.) Smiths' sea fishes. Springer-Verlag, Berlin.
- 006771** Kailola, P.J. 1991. The fishes of Papua New Guinea: a revised and annotated checklist. Vol. III. Gobiidae to Molidae. Research Bulletin No. 41, Research Section, Dept. of Fisheries and Marine Resources, Papua New Guinea. 153 p.
- 006773** Pietsch, T.W. and D.B. Grobecker. 1987. Frogfishes of the world. Systematics, zoogeography, and behavioral ecology. Stanford University Press, Stanford, California. 420 p.
- 006783** Tongyai, M.L. 1970. Plah in-see, *Scomberomorus* spp., of Thailand, 1967. p. 557-564. In J.C. Marr (ed.) Kuroshio - a Symposium on the Japan Current. Honolulu, East-West Center Press, Honolulu, Hawaii.
- 006810** Ofori-adu, D.W. 1988. List of fishes, shellfishes and other marine food resources in the Ghanaian coastal waters. Mar. Fish. Res. Tech. Pap. No. 1. 43 p.
- 006822** Lewis, A.D., B.R. Smith and C.P. Ellway. 1983. A guide to the common tuna baitfishes of the South Pacific Commission area. South Pacific Commission, Handbook No. 23, Noumea, New Caledonia
- 006871** Last, P.R. and J.D. Stevens. 1994. Sharks and rays of Australia. CSIRO, Australia. 513 p. 84 plates.
- 006934** Chen, H.M., K.T. Shao and C.T. Chen. 1994. A review of the muraenid eels (Family Muraenidae) from Taiwan with descriptions of twelve new records. Zool. Stud. 33(1):44-64.
- 006956** De la Paz, R., N. Aragonés, and D. Agulto. 1988. Coral-reef fishes off western Calatagan, Batangas (Luzon Island, Philippines) with notes on new and rare captures and controversial taxa. Philipp. J. Sci. 117:237-318.
- 006992** Sudradjat, A., U. Rempe and S. Ehrlich. 1982. Biometric comparison of the splendid ponyfish, *Leiognathus splendens* (Cuvier) from the Sunda shelf. Bull. Pen. Perikanan 2(1):17-36.
- 007050** Kottelat, M., A.J. Whitten, S. N. Kartikasari and S. Wirjoatmodjo. 1993. Freshwater fishes of Western Indonesia and Sulawesi. Periplus Editions (Hong Kong) Ltd. 221 p. + plates.
- 007100** Khin, U. 1948. Fisheries in Burma. Supdt. Govt. Printing and Stationery, Rangoon, Burma. 186 p.
- 007238** Castle, P.H.J. 1984. Muraenesocidae. In W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Volume 1. FAO, Rome.
- 007247** Allen, G.R. 1991. Damselfishes of the world. Mergus Publishers, Melle, Germany. 271 p.
- 007293** Heemstra, P.C. 1984. Monodactylidae. In W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). Vol. 3. FAO, Rome. [pag. var.].
- 007294** Allen, G.R. 1984. Toxotidae. In W. Fischer and G. Bianchi (eds.) FAO species identification sheets for fishery purposes. Western Indian Ocean (Fishing Area 51). volume 4. [var. pag.]. FAO, Rome.
- 007300** Paxton, J.R., D.F. Hoese, G.R. Allen and J.E. Hanley. 1989. Pisces. Petromyzontidae to Carangidae. Zoological Catalogue of Australia, Vol. 7. Australian Government Publishing Service, Canberra, 665 p.
- 007401** Smith-Vaniz, W.F. 1987. The saber-toothed blennies, Tribe Nemophini (Pisces: Blenniidae): an update. Proc. Acad. Nat. Sci. Philadelphia 139:1-52.
- 008524** Randall, J.E. and J.E. McCosker. 1992. Revision of the fish genus *Luzonichthys* (Perciformes: Serranidae: Anthiinae), with descriptions of two new species. Indo-Pac. Fishes 21:1-21.
- 008525** Gon, O. 1993. Revision of the cardinalfish genus *Cheilodipterus* (Perciformes: Apogonidae), with description of five new species. Indo-Pac. Fishes 22:1-59.
- 008527** Hoese, D.F. and H.K. Larson. 1994. Revision of the Indo-pacific gobiid fish genus *Valenciennea*, with descriptions of seven new species. Indo-Pac. Fishes 23:1-71.
- 008630** Mould, B. 1994. A world list of rays. The scientific nomenclature and distribution of the recent Batoidea (Batoidea, Elasmobranchii, Chondrichthyes). University of Nottingham.
- 008631** Kuitert, R. H. 1992. Tropical reef-fishes of the western Pacific Indonesia and adjacent waters. Gramedia Pustaka Utama, Jakarta. 314 p.
- 008912** Böhlke, J.E. and J.E. Randall. 1981. Four new garden eels (Congridae, Heterocongrinae) from the Pacific and Indian oceans. Bull. Mar. Sci. 31(2):366-382.
- 008927** Randall, J.E. and E. Clark. 1993. *Helcogramma vulcana*, a new triplefin fish (Blennioidae: Tripterygiidae) from the Banda Sea, Indonesia. Revue fr. Aquariol. 20(1):27-32.
- 008935** Springer, V.G. and J.E. Randall. 1992. *Platygiobopsis akihito*, new genus and species of gobiid fish from Flores, Indonesia. Japan. J. Ichthyol. 38(4):349-355.
- 008937** Poss, S.G. 1982. A new apoactinid fish of the genus *Kanekonia* from Indonesia and redescription of *K. florida*. Japan. J. Ichthyol. 28(4):375-380.
- 008940** Randall, J.E. 1993. *Acanthurus tristis*, a valid Indian ocean surgeonfish (Perciformes: Acanthuridae). Spec. Publ. J.L.B. Smith Inst. Ichth. (54):1-8.
- 008989** Allen, G.R. and R.H. Kuitert. 1989. *Hoplolatilus luteus*, a new species of malacanthid fish from Indonesia. Rev. Fr. Aquariol. 16(2):39-41.
- 008991** Dooley, J.K. 1978. Systematics and biology of the tilefishes (Perciformes: Branchiostegidae and Malacanthidae) with descriptions of two new species. NOAA Tech. Rep. NMFS Circ. No. 411:1-78.
- 009018** Winterbottom, R. and M. Burrige. 1993. Revision of the species of *Priolepis* possessing a reduced transverse pattern of cheek papillae and no predorsal scales (Teleostei; Gobiidae). Can. J. Zool. 71:494-514.
- 009069** Dooley, J.K. and P.J. Kailola. 1988. Four new tilefishes from the northeastern Indian Ocean, with a review of the genus *Branchiostegus*. Japan. J. Ichthyol. 35(3):247-260.
- 009070** Wantiez, L. 1993. Les poissons des fonds meubles du lagon Nord et de la Baie de Saint-Vincent de Nouvelle-Caledonie: Description des peuplements structure et fonctionnement des communautes. Ph.D. Thesis, Universite d' Aix-Marseille II, France.
- 009137** Masuda, H. and G.R. Allen. 1993. Meeresfische der welt - Groß-Indopazifische region. Tetra Verlag, Herrenteich, Melle. 528 p.
- 009184** Matsuura, K. 1994. *Arothron caeruleopunctatus*, a new puffer from the Indo-western Pacific. Japan. J. Ichthyol. 41(1):29-33.
- 009232** Satapoomin, U., R.H. Kuitert, and J.E. Randall. 1994. First record of the parrotfish *Scarus viridifucatus* from Thailand (the Andaman Sea) and Indonesia. Phuket mar. biol. Cent. Res. Bull. (59):5-9.
- 009360** Randall, J.E. and M. Goren. 1993. A review of the gobioid fishes of the Maldives. Ichthyol. Bull. J.L.B. Smith Inst. Ichthyol. (58):1-37, 5 pls.
- 009407** Kuitert, R.H. and H. Debelius. 1994. Southeast Asia tropical fish guide. IKAN-Unterwasserarchiv, Frankfurt, Germany, 321 p.
- 009682** Collette, B. B. 1996. Belonidae. Needlefishes. In K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes.

- The Western Central Pacific.
- 009685** Feltes, R.M. 1996. Polynemidae. Threadfins. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009710** Lieske, E. and R. Myers. 1994. Collins Pocket Guide. Coral reef fishes. Indo-Pacific & Caribbean including the Red Sea. Haper Collins Publishers, 400 p.
- 009760** Ivantsoff, W. 1996. Atherinidae. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009771** Richards, W. 1996. Triglidae. Gurnards, sea robins, armored gurnards, and armored sea robins. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009772** Sasaki, K. 1996. Sciaenidae. Croakers, drums. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009774** Amaoka A. and D.A. Hensley. 1996. Paralichthyidae. Sand flounders. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009786** Ivantsoff, W. 1996. Pseudomugilidae. Blue eyes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009790** Knapp, L. 1996. Platycephalidae. Flatheads. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009792** Hensley, D.A. 1996. Pleuronectidae. Righteye flounders. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009799** Larson, H. 1996. Centropomidae. *In* K.E. Carpenter and V. Niem (eds.) FAO identification guide for Fishery Purposes. The Western Central Pacific.
- 009800** Heemstra, P. C. 1996. Drepanidae. Sicklefishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009804** Matsuura, K. 1996. Triacanthidae. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009812** Harrison, I.J. and H. Senou. 1996. Mugilidae. Mulllets. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009822** Munroe, T. and M. Nizinski. 1996. Engraulidae. Anchovies. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009823** Westneat, M.W. 1996. Labridae. Wrasses, hogfishes, razorfishes, corises, tuskfishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009824** Amaoka, K. and D.A. Hensley. 1996. Bothidae. Lefteye flounders. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009826** Fritzsche, R. A. 1976. A review of the cornetfishes, genus *Fistularia* (Fistulariidae), with a discussion of intrageneric relationships and zoogeography. *Bull. Mar. Sci.* 26(2):196-204.
- 009835** Paxton, J.R. 1996. Myctophidae. Lanternfishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009836** Ivantsoff, W. 1996. Telmatherinidae. Sailfin silversides. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009839** Parin, N. V. 1996. Exocoetidae. Flyingfishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009840** Compagno, L.J.V. 1996. Dasyatidae. Stingrays. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009843** Collette, B.B. 1996. Hemiramphidae. Halfbeaks. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009844** Paxton, J.R. 1996. Megalomycteridae. Bignose fishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009862** Compagno, L.J.V. 1996. Myliobatidae. Eagle rays. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009893** Munroe, T. 1996. Soleidae. Soles. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009895** Munroe, T. 1996. Cynoglossidae. Tongue Soles. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009909** Compagno, L.J.V. 1996. Rhinobatidae. Guitarfishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009912** Compagno, L.J.V. 1996. Narcinidae. Longnose electric rays. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009913** Compagno L.J.V. 1996. Narkidae. Shortnose electric rays. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009915** Compagno, L.J.V. 1996. Rhinidae. Sharkfin guitarfishes or wedgefishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009918** Compagno, L.J.V. 1996. Gymnuridae. Butterfly rays. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 009936** Allen, G. R. and R. C. Steene. 1995. Notes on the ecology and behaviour of the Indonesian cardinalfish (Apogonidae) *Pterapogon kauderni* Koumans. *Revue fr. Aquariol.* 22(1-2):7-10.
- 009947** Randall, J. E. 1996. Mullidae, goatfishes. *In* K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific.
- 010591** Randall, J.E. 1988. Three new damselfishes of the genus *Chromis* (Perciformes: Pomacentridae) from the Indian Ocean. *Revue. fr. Aquariol.* 15(2):49-56.
- 010988** Collette, B.B. 1974. The garfishes (Hemiramphidae) of Australia and New Zealand. *Records of the Australian Museum* 29(2):11-105.
- 011015** Eschmeyer, W.N. and B.B. Collette. 1966. The scorpionfish subfamily Setarchinae, including the genus *Ectreposebastes*. *Bull. Mar. Sci.* 16(2):349-375.
- 011081** Herre, A.W.C.T. 1940. Notes on fishes in the Zoological Museum of Stanford University, VII, New and rare Philippine gobies from the Herre 1936-1937 oriental expedition, and in the collections of the Bureau of Science. *Philipp. J. Sci.* 72(4):357-367.
- 011441** Randall, J.E. 1995. Coastal fishes of Oman. University of Hawaii Press, Honolulu, Hawaii. 439 p.
- 011893** Allen, G.R., R.H. Kuitert, and J.E. Randall. 1994. Descriptions of five new species of cardinalfishes (Apogonidae: *Apogon*) from Maumere Bay, Flores, Indonesia and surrounding regions. *Revue fr. Aquariol.* 21(1-2):27-38.
- 011894** Allen, G.R. and R.H. Kuitert. 1994. Descriptions of two new species of cardinalfishes (Apogonidae) from Malaysia and Indonesia. *Revue fr. Aquariol* 21(1-2):19-23.

Appendix II. Indexes

[Lampiran II. Indeks.]

Author*/Name Index

- Abdul Aziz bin I. 37-38
Abdullah, M. 26, 38
Abdullah, M.A.S. 291
Abe, T. 287, 288
Abe, Y. 294
Abu Khair Mohammad Mohsin 294
Abushusha, T.L. 294
Adkin, F. 291
Adli, A. 54
Aglan, A. 123, 130, 277, 281-282, 295
Agulto, D. 295
Aken, H.M. van 7, 14
Akimichi, T. 294
Al-Ogaily, S.M. 294
Aldan, D.T. 294
Alexander, L.M. 6, 14
Allan, R.J. 10, 13
Allen, G.R. 218, 275, 284, 287-288, 292, 294-296
Alverson, D.L. 2, 6
Amaoka, A. 296
Amaoka, K. 284, 288, 296
Amornchairojkul, S. 290
Anderson, J. 26, 38
Anderson, W.D., Jr. 294
Annigeri, G.G. 291
Anon. 25-28, 30-38, 40-41, 45, 50, 54, 134, 281-282, 286-287, 293
Aoyama, T. 49, 54
Apparao, T. 285
Aprieto, V.L. 290
Araga, C. 82, 90, 284
Aragones, N. 295
Armada, N. 287
Armitage, R.O. 288
Arora, H.L. 290
Atmadja, S.B. 285, 286
Auerbach, P.S. 291
Azmi Ambak, Mohd. 294
Babenerd, B. 85, 90
Babun Rao, M. 289
Badrudeen, M. 290
Badrudin, M. 48, 50, 54-55, 61, 63, 65-66, 76, 92, 277, 281-282, 287
Bagnis, R. 292
Bailey, C. xii, 32, 37-38, 105, 122
Bailey, R.M. 218, 275, 284, 291
Baillon, N. 294
Bakhader, A. 286
Bakun, A. 15, 17, 23
Balan, V. 285
Banerji, S.K. 285, 287
Banon, S. xii, 285
Barnett, M.A. 291
Basyori, A. 48, 54
Bathia, U. 287, 289
Bauchot, M.L. 284, 289-292, 294
Bauer, R. 9, 13
Baum, G.A. 68, 76
Bawazeer, A.S. 290
Bay of Bengal Programme 291
Beardsley, G.L. 284
Beaufort, L.F. de 82, 90, 217-218, 275
Beck, I.M. 286
Beck, U. 286
Beddington, J.R. 2, 6, 67, 76, 90-91, 101
Bekker, V.E. 292
Ben Yami, M. 285
Ben-Tuvia, A. 293
Bennett, J. 292
Bergstad, O.A. 286
Berkeley, H. 28, 39
Bertage, H.P. 64, 76
Berry, F.H. 289
Bertalanffy, L. von 136, 199, 216
Bianchi, G. vii, 1, 5-6, 13, 50, 54-55, 57, 63, 65-66, 92-94, 99-101, 123-126, 130, 135, 276-278, 281-282, 284, 287-291, 293-295
Bienvenida, S.L. 5, 41, 50-51, 54, 61, 85, 90, 109, 124, 127, 130-131, 134-135, 146, 218, 276
Bilio, M. vii, x, xi
Birowo, S. 64, 76
Birtwistle, W. 41, 46
Bishop, B.P. 218
Bjarnason, B. 105, 122
Blaber, S.J.M. 292
Blanford, W.T. 288
Bleeker, P. 284, 286
Bliss, C.I. 65, 76
Bloch, M.E. 284
Boëly, T. 105-106, 119, 122
Bogdanov, A.S. 46
Böhike, J.E. 295
Bond, C.E. 218, 275, 284, 291
Boonraksa, V. 286
Boonwanich, T. 288, 290
Boraey, F.A. 294
Borges, T.C. 292
Borgese, E. M. 39
Bottomanne, C.J. 35, 38
Bouhleb, M. 286, 293
Boyer, T. 6
Brandhorst, W. 78
Brinca, L. 286, 288
Brodie, J.W. 117
Brook, D.R. 5, 6
Brooker, J.R. 218, 275, 284, 291
Brothers, E.B. 290
Brouard, F. 284
Bruce, R. W. 284
Buchary, E.A. vii
Budihardjo, S. 45, 50, 54-55, 63, 65-66, 68, 76, 92, 277, 281, 285, 291
Burgess, W.E. 292
Burhannudin 285, 293
Burrige, M. 294, 295
Butcher, J. 24, 31, 38
Bykov, V.P. 292
Cabanban, A. 5, 11, 13, 41, 50-51, 54, 61, 85, 90, 105, 107, 109, 116-117, 120, 124, 127, 130-131, 134-135, 146, 218, 276, 279
Caces-Borja, P. 284, 285, 293
Cada, R. 135
Campbell, D.R. 291
Canagaratnam, P. 289
Cantor, T.E. 286
Capuli, E.C. 5, 51, 92, 135, 137, 217, 277, 279, 282
Carcasson, R.H. 287
Carlsson, D. 287, 289
Carpenter, K.E. 218, 284, 287, 295-296
Carvalho, J.L. 41, 46
Castañeda, P. 34, 38
Castle, P.H.J. 289, 290, 293, 295
Central Fisheries Dept. Pakistan 289
Cervigón, F. 292
Chabanne, J. 285
Chan, E.H. 287
Chan, W. 287, 289
Chavez, F. 16
Chayakul, R. 286
Chen, C.F. 293
Chen, C.T. 292, 295

*Excludes authors of species names, e.g., "(Bleeker 1867)".

Chen, H.M. 295
Cheunpan, A. 285
Chirichigno, N.F. 293
Chiu, T.-S. 292
Choat, J.H. 288, 294
Chotiyaputta, K. 286
Christensen, J. M. 46, 285, 286
Christensen, V. 1, 5-6, 13-14, 17, 23, 75-76
Christian, E. 292
Christy, F.T., Jr. 2, 6
Chua, T.E. 289
Chullasorn, S. 286
Cinco, E. 293
Cipriani, R. 292
Clark, C.W. 67, 76
Clark, E. 295
Clifford, H. 26-27, 30, 38
Cohen, D.M. 285, 291
Colban, R.L. 288
Collijn, H. 29, 38
Collette, B.B. 284, 294, 295, 296
Compagno, L.J.V. 284, 289, 296
Conand, F. 286, 293
Conlu, P.V. 288, 291
Contemprate, A. 130
Cooke, J.G. 2, 6, 90
Coppola, S.R. 284
Coronado, G. 136, 216
Corpuz, A. 286
Corpuz, V.T. 34, 38
Crawford, R. 295
Crawfurd, J. 25, 38
Cressey, R.F. 290
Crossman, E.J. 294
Csirke, J. 23
Currie, H.M. 288
Curtis, C. 17, 23
Cury, P. 17, 23
Cushing, D.H. 33, 38, 77, 85, 90
Cuvier, G. L. 287
Dadang Karyana 105
Daget, J. 287, 289, 291
Dahlström, P. 3, 6
Dalzell, P. 13, 287, 290, 292
Danil'chenko, P.G. 292
DataEase 276, 282
David, N. 287
Dawson, C.E. 287, 292, 293
Day, F. 284, 288
Dayaratne, P. 286
De Beaufort, L.F. 292
de Freitas, A. 292
de la Hoz, M.M. 291
De la Paz, R. M. 293, 295
de Leon, F.P. 293
Debelius, H. 295
Delgado de Molina, A. 293
Delsman, H.C. 35, 38
Department of Fisheries., Malaysia. 292
Deshmukh, V.M. 285
Devaraj, M. 286, 291
DGF (Directorate General of Fisheries) 76, 276, 282
Dhulked, M.H. 291
Diamant, A. 288
Dias, M. 293
Dizon, A.E. 13-14
Dizon, L.B. 101, 130
Djajadiredja, R. 219, 275
Djamali, A. 285, 293
Donaldo, S.S. 288
Dooley, J.K. 295
Doty, M.S. 64, 76
Doubleday, W.G. 1, 6, 83, 90
Dunker, J.C. 7, 13
Durand, J. 292
Durand, M.H. 17, 23
Dwiponggo, A. xii, 37-38, 55, 61, 63, 66, 76, 83, 90, 105-106, 122, 281-283, 285, 287
Ebeling, A.W. 290
Eccles, D.H. 292
Edi Amin 105
Edwards, A.J. 292, 293
Edwards, R.R.C. 286, 290
Eggleston, D. 287, 289
Ehrich, S. 295
Ekman, S. 57, 61
Ellway, C.P. 295
Emery, A. R. 284
Emery, K.D. 40-41, 46, 63-64, 76, 100-101
Erzini, K. 294
Eschmeyer, W.N. 137, 216, 218-219, 275, 287, 290, 296
Evermann, B.W. 286
Everson, A.R. 294
Fager, E.W. 91, 101
FAO 67, 76, 219, 275, 284, 292
FAO/IOP 120, 122
Farquhar, G. Brooke 122
Federizon, R.R. 92, 101, 124, 130, 293
Feltes, R.M. 296
Feral, J.P. 292
Ferrell, D.J. 294
Føyn, L. 123, 130, 277, 281-282, 295
Firth, R. 32, 36, 38
Fischer, W. 50, 54, 82, 90, 132, 134, 218, 275, 284, 287-292, 294-295
Flowers, J.M. 277
Flüs, J. 288
Formacion, S.P. 136, 216
Fouda, M.M. 295
Fowler, H.W. 284, 289
Fox, P. 285
Fraser, T.H. 284, 287
Fritzsche, R.A. 296
Froese, R. vii, xii, 3, 5-6, 51, 92, 101, 135-137, 216-219, 275, 277, 279, 282
Gabriel, W.L. 91, 101
Ganaden, R.A. 287
Ganaden, S.R. 285
Garcia, S.M. 2, 6, 277, 282
Garibaldi, L. 284, 292
Gauch, H.G. 124, 130
Gayanilo, F.C., Jr. 5-6, 136, 216, 277-278, 281-282
Gelfeld, D. 6
George, K. 285
Ghofar, A. 5, 11, 79, 107, 116, 120, 145-146
Gibbs, R.H. 291
Ginsburg, N. 39
Gislason, M. 286
Gjøsaeter, J. 286, 293
Gjøsaeter, V. 286
Glaser, T. 285
Glass, C.W. 292
Gloerfelt-Tarp, T. vii, x, xii, 83, 90, 217-220, 275, 297
Gobée, E. 30, 38
Godø, O.R. 123, 130, 277, 281-282, 295
Golani, D. 288
Gold, B.D. 6, 14
Gomon, J.R. 294
Gon, O. 295
Goodyear, R.H. 291
Goren, M. 295
Gosse, J.P. 287, 289, 291
Gould, H.R. 100-101
Grabda, E. 294
Gramberg, J.S.G. 39
Grandperrin, R. 284
Gray, I.E. 288
Greenfield, D. W. 293
Grobeck, D.B. 295
Guano, M.B. 290

Guéze, P. 293
 Gulland, J.A. 1-2, 6, 44-45, 54, 62-63, 68, 76, 90, 134
 Gunderson, D.R. 56, 61
 Günther, A. 289
 Hackert, E.C. 10, 13
 Hadisubroto, I. 48, 54
 Haedrich, R.L. 289, 291
 Haga, B.J. 29, 39
 Halstead, B.W. 291
 Hamada, R. 285
 Hanley, J.E. 295
 Hansen, P.E.H. 294
 Hanson, A. 76
 Hanyu, I. 292
 Hardenberg, J.D.F. 105-106, 113, 122, 289
 Hariati, T. xii, 285
 Harrison, I.J. 296
 Hasselman 29, 31-32, 39
 Hastenrath, S. 10, 13, 15, 23
 Hatta 36
 Haudricourt, A.-G. 294
 Heemstra, P.C. 218, 275, 288-293, 295-296
 Heese, T. 294
 Hegazy, M.R. 286
 Hendrickx, M. 292
 Hensley, D.A. 288, 291, 296
 Hermes, R. 133
 Hermosa, G.V., Jr. 295
 Herre, A.W.C.T. 35, 39, 284, 288, 291, 296
 Høisæter, T. 125, 130
 Hida, T. 2, 6
 Hill, M.O. 93, 101, 124, 130
 Hirano, R. 292
 Hoese, D.F. 284, 293, 295
 Holt, S.J. 67, 76
 Honebring, R. 292
 Honig, P. 39
 Hosillos, L.V. 101, 130
 Hughes, G.M. 287, 288
 Hulley, P.A. 290
 Hureau, J.C. 288, 290-292, 294-295
 Husby, D.M. 15, 17, 23
 Hussain, A. 294
 Hussain, N.A. 291
 Hutchins, B. 288
 Hutchins, J. B. 284
 Hutomo, M. 285
 Ianelli, J. 291
 Ikehara, I.I. 293
 Ilahude, A.G. 64, 76, 85, 90
 Imad, A. 290
 Inada, T. 285
 Ingles, J. 285
 Interior, R. 293
 International Game Fish Association. 291
 Iqbal, M. 294
 Isa, M.M. 291
 Isom Hadisubroto 105
 Ito, B.M. 294
 Ivantsoff, W. 292, 296
 Iwamoto, T. 285
 Jabat, M. 292
 Jajuri 120
 James, P.S.B.R. 289, 290
 Janzen, D. 6
 Jarchau, P. 48, 51, 54
 Jayabalan, N. 290
 Jayaram, K.C. 289
 Jenne, R.L. 17, 23
 Jensen, E.A. 85, 90
 Jhingran, A.G. 292
 Johnson, D. 6
 Jones, A. 102, 122
 Jones, G.P. 294
 Jones, S. 294
 Jorde da Silva, A. 286
 Jothy, A.A. 285
 Joung, S.-J. 292
 Kailola, P.J. vii, x, 83, 90, 217-220, 275, 288, 294, 295
 Kalk, M. 293
 Kamara, A.B. 291
 Kami, H.T. 293
 Kampen, P.H. van 33, 35, 39
 Karnella, S. J. 293
 Karrer, C. 290, 291, 294, 295
 Kartikasari, S.N. 217, 275, 295
 Karunasinghe, W.P.N. 294
 Katili, J.A. 7, 14
 Kedidi, S.M. 286, 294
 Keller, G.H. 41, 46
 Khin, U. 295
 Klausewitz, W. 177
 Knapp, L.W. 289, 296
 Kosaka, M. 285
 Kottelat, M. 217, 275, 295
 Krey, J. 85, 90
 Krishna Bhat. C.H. 286
 Krishnan, T.S. 285, 287
 Krisnandhi, S. 36-37, 39
 Kühlmorgen-Hille, G. 50, 54, 285, 287, 289
 Kuitert, R.H. 217-218, 288, 295-296
 Kulbicki, M., 293-294
 Kumaran, M. 289, 294
 Kuronuma, K. 288, 294
 Lachner, E.A. 218, 275, 284, 287, 291, 293
 Lai, H.C. 289
 Lal Mohan, R.S. 289
 Lam, W.C. 37, 39
 Lamb, B.G. 288
 Lamb, P.J. 10, 13, 15, 23
 Langham, N. 289
 Larson, H. 295, 296
 Last, P.R. 295
 Latiff, S.S. 41, 46
 Law, R.M. 67, 76
 Le Poulain, F. 292
 Lea, R.N. 218, 275, 284, 291
 Leblic, I. 294
 Lee, Y.-C. 294
 Lemus, A.J. 292
 Lester, R.J.G. 291
 Levitus, S.R. 6
 Lewis, A.D. 295
 Leyland, G.G. 288
 Leyndekkers, J.V. 85, 90
 Lieske, E. 296
 Liew, H.C. 287
 Lijam, N. 295
 Lindsay, S.R. 290
 Linnaeus, C. 286
 Liu, H.-C. 294
 Lockley, G.J. 288
 Lohmeyer, U.P. 13, 77, 90, 92, 99, 103, 111, 115, 117, 132, 135, 276, 281-282
 Longhurst, A.R. 5-6, 12, 14, 91-92, 101, 123, 130, 276, 282
 Loubens, G. 287, 294
 Low, H. 27, 39
 Lubbock, R. 287
 Luna, S. vii, 5, 51, 92, 135, 137, 217, 277, 279
 Luther, G. 285, 286
 Machida, Y. 294
 Maclean, J.L. 101, 130
 Macleay, W. 289
 Macnae, W. 293
 Magnusson, J. 287, 290
 Mahyam Binti Mohd. Isa 40, 46

Makwaia, E.D.S. 290
 Malumay, L. 101, 276, 282
 Mangalik, A. 290
 Mangundjojo 54
 Manickchand-Heileman, S. 288
 Manooch, C.S. 288
 Mansor Mat Isa 40, 46
 Marahudin, F. xii, 37-38, 105, 122
 Markle, D.F. 290
 Marquardt, D.W. 136, 216
 Márquez, R. 292
 Marr, J. 46, 285, 295
 Martin, T.J. 290
 Martosewojo, S. 285, 293
 Martosubroto, P. vii-ix, 1, 4-5, 7, 9, 15-17, 24, 40 -41, 43-46, 47-49, 51-56, 61-63, 65-66, 76-77, 79, 91, 102, 104, 108-109, 113, 118, 123, 131, 135, 197, 199, 217, 276, 281-282, 285, 287
 Mascarenhas, V. 288
 Massinga, A. 292
 Masuda, H. 82, 90, 284, 295
 Mathews, C.P. 5-6, 11, 30, 39, 79, 107, 116, 120, 145-146, 290
 Mathias, J.A. 289
 Matsuura, K. 295, 296
 Maugé, L. A. 291
 Maul, G.E. 291
 Maxwell, C.N. 33, 38-39
 May, R.M. 2, 6, 67, 76, 101
 Mazellier, P. 292
 McCosker, J.E. 289, 290, 292, 295
 McFarland, W.N. 288
 McKay, R.J. 288, 289, 294
 McLennan, D.A. 5-6
 McManus, J.W. vii, 5-6, 41, 50-51, 54, 57, 61, 76, 85, 90-92, 98, 100-101, 109, 124, 127, 130-131, 134-135, 146, 218, 276-277, 279-281, 283
 McPherson, G.R. 294
 Meade, R.H. 7, 13-14
 Medcof, J.C. 289
 Menasveta, D. 40-41, 43, 46
 Mendelssohn, R. 17, 23
 Mendo, J. 5-6, 17, 23
 Menon, A.G.K. 287, 289, 293
 Menon, M. 290
 Menon, N.G. 291
 Mercer, M.C. 101
 Merrett, N.R. 290
 Merta, I.G.S. 293
 Milliman, J.D. 7, 13-14
 Misra, K.S. 289
 Miyasaka, A. 293
 Moeljanto, R. 293
 Mohan Joseph, M. 290
 Monkolprasit, S. 287, 293
 Monod, Th. 288, 292
 Mook, W.G. 7, 9, 14
 Moor, J.H. 26, 39
 Moosa, M.K. 293
 Morales-Nin, B. 290
 Morgan, G.R. 5-6, 283, 286-287
 Morgan, M. 287
 Mou Tham, G. 293
 Mould, B. 295
 Mubarak, H. 293
 Muck, P. 5-6
 Muhamad Nasir Abdul Salam 294
 Mulyadi, E. 285
 Munro, I.S.R. 50, 54, 82, 90, 132, 134, 284, 286, 289
 Munro, J.L. 288
 Munroe, T. 296
 Munz, F.W. 288
 Murdjijo, F.X. ix
 Murdy, E.O. 284, 292
 Murphy, G. 285
 Muthu, M.S. 286
 Muus, B.J. 3, 6
 Myers, R.F. 218, 275, 286, 296
 Myklevoll, S. 123, 130, 277, 281-282, 295
 Naamin, N. 69, 76
 Nakamura, I. 284, 294
 Nammalwar, P. 285
 Narasimham, K.A. 291
 Nauen, C.E. 284
 Neiman, A.A. 41, 46
 Nelson, C.S. 15, 23
 Nelson, G.J. 284
 Nelson, J.S. 61, 218, 275, 284
 Newton, C. 2, 6
 Nhwani, L.B. 290
 Nichols, N. 10, 14
 Nielsen, J. 290-292, 294
 Niem, V. 295, 296
 Nizinski, M. 296
 Njock, J.C. 291
 Nontji, A. 64, 76, 85, 90
 Norkis, M.J. 94, 101
 Nouguier, J. 294
 Obika, Y. 293
 Ofori-adu, D.W. 295
 Okamura, O. 294
 Okera, W. 291
 Olney, J.E. 290
 Ommanney, F.D. 37, 39, 41, 46
 Ong, K.S. 41, 46
 Organisation for Economic Co-operation and Development 289
 Østvedt, O.J. 123, 130, 277, 281-282, 295
 Overholtz, W.J. 91, 101
 Ozanne-Rivierre, F. 294
 Padilla, J.E. 294
 Palha de Sousa, B. 288
 Palha de Sousa, L. 288
 Palko, B.J. 284
 Palomares, M.L. xii, 285, 292
 Palsson, W. A. 286
 Panggabean, T. 48, 54
 Parin, N.V. 292, 294, 296
 Pariwono, J.I. 10, 13
 Parrish, J.D. 294
 Parrish, R.H. 15, 17, 23
 Pathansali, D. 37, 39, 41, 46, 287
 Patiale, H. 290
 Patten, J. M. 292
 Paul, L.J. 288
 Pauly, D. vii-x, xii, 1-7, 11-15, 17, 23-24, 30, 37, 39-41, 44-56, 61-63, 65-66, 68, 75-79, 83, 85, 90-93, 100-102, 104-105, 107-109, 113, 116-117, 120, 122-124, 127, 130-131, 134-135, 146, 177, 197, 199, 216-217, 219, 275-279, 281-287, 290-292
 Paxton, J.R. 295, 296
 Payne, D.A. 288
 Pel, H. van 33-34, 39
 Pender, P.J. 291
 Peng, C.-Y. 292
 Peñafior, G.C. 288, 292
 Pereyra, W.T. 2, 6
 Philbrick, C. 290
 Pielou, E.C. 67, 76
 Pietsch, T.W. 286, 295
 Plante, R. 285
 Plas, van der 32
 Platt, T. 23
 Poll, M. 289
 Polovina, J.J. 91, 101, 288, 294
 Pope, J.G. 2, 6, 67, 72, 76
 Por, F.D. 293
 Poss, S.G. 289, 295
 Post, A. 290, 291, 294, 295
 Postma, H. 7, 9, 14

Poutiers, J.M. 292
Prabhu, M. 285
Prein, M. 41, 50-51, 54, 61, 85, 90, 109, 124, 127, 130-131, 134-135, 146, 218, 276
Punjanan, J. 7, 14
Quero, J.C. 290-291, 294-295
Radhakrishnan, N. 290
Radtke, R.L. 294
Rafail, S.Z. 284
Ralston, S. 91, 101, 287, 288, 294
Rama Rao, K.V. 289
Randall, H.A. 288
Randall, J.E. 218, 275, 284, 287-296
Rao, B.N. 286
Rao, K.S. 290
Rao, K.V.S. 291
Rathborne, A.B. 26, 39
Rau, A. 284
Rau, N. 284
Razak, H. 293
Refait, D. 294
Rempe, U. 295
Reuter, J. 33, 39
Richards, A.F. 41, 46
Richards, W. 284, 296
Ricker, W.E. 45-46, 67, 76, 90
Riehl, P.M. van 64, 76
Ritragasa, S. 2, 6, 50, 54
Ritterbush, S. 289
Rivard, D. 1, 6, 83, 90
Robaina, G. 292
Robins, C.R. 218, 275, 284, 291
Robinson, H.C. 28, 39
Robinson, M. 9, 13
Rodriguez, B. 292
Rodriguez, M.I. 285
Roedel, P. 54, 90
Rofen, R.R. 288
Rohlf, F.J. 67, 76
Rongo, J.M. 136, 216
Ronquillo, I.A. 284, 285, 287, 290, 293
Rosa, H., Jr. 285
Rose, J.H. 291
Rosenblatt, R. H. 292
Roux, C. 287
Roy, C. 5, 15, 17, 57, 124, 130, 134
Russell, B.C. 288, 290
Sadhotomo, B. 5, 120, 286
Saeger, J. 1, 4-5, 41, 44, 46-48, 54-56, 61, 63, 65-66, 76, 90, 104, 130, 135, 197, 199, 277, 286
Saetre, R. 286, 288
Saila, S. 54, 90
Saimima, S. 7, 14
Sainsbury, K.J. 288
Saldanha, L. 290, 291, 294, 295
Sale, P.F. 294
Sambalay, V.C., Jr. 136, 216, 286, 288
Samuel, M. 290
Sanders, M.J. 286
Sansaloni, J.A. 122
Santana, J.C. 293
Santos, L.M. 287
Santosa, W. 48
Sardjono, I. 45-46, 54, 63, 76
Sasaki, K. 296
Satapoomin, U. 295
Sauchomal, S. 294
Saunders, M.J. 290
Schaefer, M.B. 67-68, 76
Schippers, A.W. Jzn. 28, 39, 32
Schneider, J.G. 284
Schneider, M. 284, 289
Schneider, W. 288, 292
Schroeder, R.E. 290
Schuster, W.H. 219, 275, 284
Scialabba, N. 284, 285
Scott, J. S. 286
Scott, W.B. 218, 275, 284, 291, 294
SCSP (South China Sea Fisheries Development and Coordinating Programme) 40, 46, 63, 76
Seale, A. 286
Sedana Merta, I. Gede 40, 46, 105
Sekharan, K.V. 285, 286, 290
Senou, H. 296
Seshappa, G. 285
Shackleton, L.Y. 11, 14
Shaharil, T. 30, 39
Shaher, S. 286, 290
Shao, K.T. 292, 295
Sharp, G.D. 5, 7, 12-14, 15, 17, 23, 57, 107, 124, 130, 134, 146
Shen, S. 288
Shen, S.-C. 292
Shepard, F.P. 100-101
Sherman, K. 6, 14
Shindo, S. 50, 54
Shiokawa, T. 295
Shirai, S. 284
Shomura, R.S. 294
Silva, C. 292
Silvestre, G. 2, 5-6, 286, 287
Siripakhavanich, S. 290
Sivasubramaniam, K. 40, 46, 293
Sjarif, S. 64, 76
Slutz, R.J. 17, 23
Small, M. 292
Smith, A. 290
Smith, B.R. 295
Smith, D.G. 291
Smith, I.R. 2, 5-6
Smith, J.L.B. 293
Smith, M.M. 288-293, 295
Smith, P.E. 116, 122
Smith, S.E. 292
Smith-Vaniz, W.F. 288-290, 295
Sneath, P.H.A. 67, 76
Soegiarto, A. 64, 76
Soemarto 105, 122
Soeriatmadja, R.E. 64, 76
Soerjodinoto, R. 106, 122
Soewito 7, 54, 276, 282
Sokal, R.R. 67, 76
Soliman, F.M. 294
Sommani, P. 290
Somvanshi, V.S. 285
Sontirat, S. 287
Soriano, M.L. 40, 46, 282, 286
Sousa, F.E. 285
Sousa, I. 292
Sousa, L. 286
Sousa, M.I. 286, 288, 293
Sparre, P. 5-6, 136, 216, 277-278, 281, 283, 290
Spiering, W. 48, 54
Springer, V.C. 288
Springer, V.G. 291, 293, 295
Sreenivasan, P.V. 291, 293
Starnes, W.C. 293
Stead, D.G. 32, 39
Steel, R.G.D. 65, 76
Steene, R.C. 218, 275, 288, 292, 296
Stehmann-Nielsen, E. 85, 90
Steurer, P.M. 17, 23
Stevens, J.D. 295
Strømme, T. 2, 6, 124, 127, 130, 277, 280-283
Sua, T. 294
Subhat Nurhakim 105, 122
Sudekum, A.E. 294
Sudhakava, G. 286
Sudjastani, T. 285

Sudjianto 105
 Sudradjat, A. 48, 54, 286, 295
 Sujastani, T. 5, 40-41, 45-46, 48-49, 51-53, 63, 65, 76, 104, 108-109, 113
 Sukarno 36
 Sukhavisidh, P. 289
 Sulak, K.J. 292
 Susanto, V. 47
 Suvapepun, S. 130
 Suvatti, C. 286
 Suwaso. 291
 Swainston, R. 288, 294
 Sy, J.C. 34, 38
 Talbot, F. 287, 289
 Talwar, P.K. 292
 Tampubolon, G.H. 40, 46, 285, 293
 Tandog-Edralin, D. 285
 Taylor, L. 294
 Taylor, W. R. 286
 Tenison-Woods, J.E. 39
 Ter Braak, C.J.F. 127, 130
 Teugels, G.G. 289
 Teulières, M.-H. 294
 Therriault, J.C. 23
 Thiagarajan, R. 285
 Thollot, P. 293
 Thompson, B.G. 277
 Thomson, J.M. 288
 Thys van den Audenaerde, D.F.E. 287, 289, 291
 Tiews, K. 6, 37, 39, 47, 54, 62, 76, 284, 285, 287, 293
 Timochin, I. 288
 Tinker, S.W. 284
 Tirant, G. 291
 Tongyai, M.L. 295
 Torres, F.S.B., Jr. 5, 11, 13, 41, 50-51, 54, 61, 85, 90, 105, 107, 109, 116,
 120, 124, 130-131, 134-135, 146, 218, 276, 279, 290
 Torrie, J.H. 65, 76
 Tortonese, E. 290, 291, 292, 294
 Toshino, T. 82
 Tranter, D.J. 85, 90
 Tsukayama, I. 5-6, 15, 23
 Tyler, A.V. 91, 101
 Tyler, J.C. 295
 Tzeng, C.-S. 292
 U Hla Win. 294
 Uchida, R.N. 291
 Uchiyama, J.H. 291
 Udupa, K.S. 286
 Uktolseya, J. 69, 76
 Umali A.F. 288
 Unar, M. 38-39, 45, 48, 54
 Ursin, E. 199
 Uyeno, T. 284
 Vakily, J.M. 3, 6, 276, 283
 Valencia, M.J. 40, 46
 Valenciennes, A. 288
 van der Elst, R.P. 290, 291
 Zalinge, N.P. 285
 Vari, R. 290
 Venema, S.C. 1, 5, 13, 16, 41, 46, 77, 79, 90, 102, 104-105, 108, 120, 122,
 132, 134, 285, 286, 290
 Verdoorn, F. 39
 Vergara, R. 292
 Veth, P.J. 27, 39
 Viloso, E.P. 290
 Vore, R. 292
 Wade-Pauly, S. 292
 Wahyuono, H. 291
 Walczak, P.S. 291
 Walford, L. 295
 Wantiez, L. 293, 295
 Waples, R.S. 290
 Warwick, R.M. 5-6
 Wasilun 48
 Watson, R.A. 291
 Weber, M. 82, 90, 217-218, 275, 292
 Weber, W. 285
 Wenner, C.A. 288
 Westneat, M.W. 296
 Wheeler, A. 292
 White, T. 78
 Whitehead, P.J.P. 50, 54, 82, 90, 132, 134, 218, 275, 284, 287, 288, 289,
 290, 291, 292, 294
 Whitten, A.J. 217, 275, 295
 Wicklund, R. 295
 Widodo, J. 5, 40, 46, 48, 51, 54, 120, 279, 283, 285
 Wijeyaratne, M.J.S. 294
 Wilkinson, R.J. 28, 39
 Williams, F. 2, 6
 Williams, H.A. 287, 294
 Williams, J. T. 284
 Williams, M. 6
 Willing, R.S. 291
 Winterbottom, R. 284, 292, 294-295
 Wirjoatmodjo, S. 217, 275
 Wirjoatmodjo, S. 295
 Wojciechowski, J. 285
 Wongratana, T. 284, 287, 289
 Woodland, D.J. 286, 289
 Woodruff, S.D. 17, 23
 Wright, L.D. 7, 13-14
 Wu, M.C. 10, 13
 Wyrski, K. 7, 9, 10, 14-15, 17, 23, 64, 76, 85, 90, 99-101, 107, 122, 124, 130
 Yamakawa, T. 293
 Yap, C.L. 34, 37, 39
 Yeh, S.-Y. 294
 Yohannan, T.M. 286
 Yoshino, T. 82, 90, 284
 Zachman, N. 62-63, 76
 Zannou, L.H. 294
 Zeitzschel, B. 130
 Zijlstra, J.J. 14

Geographic Index*

- Aceh 25, 28, 40-42, 44-45
Aden, Gulf of 186
Africa 12, 137, 154, 165, 173, 193
 East 143, 162-163, 169, 186, 188, 190, 195, 206, 214
 South 11-12, 141, 151-152, 182, 201, 203
 West 2-3, 6, 8, 283
Alas Strait 118-119, 133
Ambon 169
America, South 8, 10-12
Americas 10-11
Anak Krakatau 117
Anambas Islands 25, 34
Andaman 150
 Islands 7, 201
 Sea 8, 141
Angola 130
Arabia 30-31
Arafura
 Sea 8-9, 12-13, 79, 277
 Shelf 7
Arends 108
Asahan 160, 202
Asia 8, 13-14, 16, 62
 East 101
 Southeast 6, 7-8, 10-13, 15-17, 24, 38, 45-47, 54, 64, 98, 100-101, 123, 130, 137, 162, 173, 175, 182, 184, 186, 188, 190, 193, 195, 201, 204, 208, 210, 214, 282
Atlantic 12, 214
 Ocean 204
 Western 165
Ator Island 7
Australasia 13
Australia ix-x, xii, 7-9, 10, 14, 16, 77-79, 83, 90, 122, 131, 137, 148, 154, 158, 162, 163, 167, 171, 175, 178, 180, 182, 184, 186, 188, 190, 193, 195, 199, 201,
 Northern 141, 143, 214
 Southeast 204
 Western vii, 141, 145, 151-152
Babar Island 7
Badjo 139, 143, 148, 175, 182, 201
Bagan Si Api Api 24-26, 29-31, 37-39, 120
Bali 77-78, 82, 85, 88-90, 91, 102, 107, 110-111, 114-116, 118, 120, 131-133, 143, 145
 Shelf 115-116
 Straits 77, 83, 85, 87, 89, 102-103, 105, 107, 114-116, 120, 122, 145, 146, 278
Banda
 Aceh 160, 202
 Banda Sea 7-9, 13, 17
Bandung 143, 178
Bangka 25, 34, 52, 66, 106, 113, 115
 Island 63
 Strait 113
Bangkaru 124, 128
Bangladesh 102, 180
Banjarmasin 110-111
Bantam 25, 32
Banyuwangi 114-115, 118
Barat Daya 7
Barito River 110
Barro 111
Batang 119-120
Batavia 24, 27-28, 32-34
Bawean 34, 108, 119, 160
 Island 105, 120
Belau 210
Belawan 103
Belitung 25, 34, 52, 66
 Island 113
Bengal, Bay of 7, 12-13, 46, 124
Benguela upwelling region 11-12
Benoa 102-103, 115-116, 118
Bergen 6, 277, 280
Berlin 47
Biabangan Peninsula 114
Biliton Island 63
Bintan 178
Blambangan 87, 115
Borneo 17, 25-27, 46, 141
 South 143, 175, 182, 205
 West 148, 171
Brazil 8
Bremen 77
Breueh 124, 125, 128
Brouwerstraat 25-28
Bugis 139, 143, 145, 163, 180, 186, 201, 208, 210
Burma 6, 31
Calang 124, 128
California, Gulf of 160
Cambodia 29
Cape of Good Hope 11-12
Cape Town Agulhas 12
Caribbean 8
Celebes Sea 7, 9
Cempai Bay 119
Ceram 169
China 27, 34, 173, 195, 201, 204
 Sea, South 6-8, 14, 17-18, 23, 25, 46-47, 52-54, 56-58, 64, 90, 103-106, 108, 120-121, 150, 197-198
Cilacap 78, 103, 115, 117-118
Cirebon 113
Comores 182
Congo 12
Costa Rica 130
Cupel 115
Denmark 132
Denpasar 132
Dindings 25, 33
Durban 141, 158
Engano 85
Equator 12
Ethiopia 92
Europe 3, 27, 35
Fiji 137, 158, 178, 182
Flores 118-119, 131
 Sea 7-9, 17, 64, 105, 111, 118-119
FMS *see* Malay States, Federated
Formosa, Strait of 7, 9
Gabon 12
Galapagos Islands 160, 165
Gaspar Strait (of) 106, 113
Germany, Federal Republic of ix, xii, 37, 47, 77-78, 131
Geser 169
Grajagan Bay 83, 88, 114-116, 118
Great Barrier Reef 151-152
Gresik 108
Guinea 6
 Gulf of 101
Haria 169
Hawaii 101, 156, 165, 190
Hong Kong 31, 145
Idi 162
India 31, 139, 141, 143, 167, 178, 180, 182, 194, 186
Indian
 Archipelago 38-39
 Ocean vii, 2, 6-14, 25, 45-46, 54, 65, 77-78, 82, 90, 91-92, 99-100, 102-103, 106-107, 115-119, 124, 130-131, 141, 145, 148, 156, 158, 167, 175, 178, 180, 188, 191, 210, 277, 282
Indies, West 165
Indo-Australian Archipelago 39, 82, 90

*Excludes occurrence records in species list, p. 222 to 274.

Indo-Pacific 5, 10-13, 82, 139, 165, 177
 Ocean 190, 206, 275
 Indonesia
 Southern vii, x, xii
 Western 1, 5, 15, 18-22, 24, 45, 47, 51, 55, 62, 69, 77, 90, 91, 102-103, 123, 131, 135-145, 147-164, 166-170, 172-174, 178, 182, 188, 190, 192-193, 195, 199-200, 202-203, 207, 209-210, 212, 214, 276, 279, 281-282
 Indonesian Archipelago 7, 10, 12-13, 15, 23, 156, 160, 162, 163, 173, 175, 178, 193
 Irian Jaya 13, 191
 Jakarta vii, 10, 25, 31, 36, 38, 48, 65, 78, 102, 103, 105, 113, 115, 117, 118, 120, 122, 133, 139, 141, 143, 145, 148, 160, 162-163, 165, 167, 169, 171, 173, 175, 178, 180, 182, 186, 195, 201, 203, 205-206, 108, 210, 212, 277
 Bay 162
 Jambi Bay 46
 Jambuairst 124
 Japan 38, 46, 82, 90, 137, 151, 152, 154, 158, 160, 162-163, 165, 169, 175, 186, 188, 190, 197, 199, 204, 208, 210, 214
 Java 9, 13-14, 16-17, 24-32, 35-36, 38-39, 47, 51, 53, 56-58, 63-66, 68-78, 83-90, 100-104, 106-110, 112-120, 122, 133, 138, 141, 143, 145, 148, 160, 162-163, 167, 169, 171, 175, 178, 180, 182, 186, 195, 205-206, 208, 210, 212
 Sea vii, 4, 7-9, 17-18, 23, 33-35, 37, 45-46, 50, 52-58, 60-66, 68, 70, 74-75, 90, 102-106, 108, 110-111, 113, 116-117, 119-122, 138, 145, 148, 150, 152, 160, 162-163, 165, 167, 171, 178, 180, 182, 184, 195, 199, 201-202, 208, 282-283
 Central 138, 150, 152, 163, 178, 180, 182, 184, 195, 199, 201, 210
 East 145
 North 138, 195
 West vii, 139, 141, 143, 145, 148, 160, 162-163, 165, 167, 169, 171, 173, 175, 178, 180, 182, 186, 195, 201, 203, 205, 206, 208, 210, 212
 Jawa 90, 109 *see* Java
 Jepara 108
 Juwana 119-121
 Kalimantan 16, 52-54, 56-58, 60-66, 68-69, 71, 73-74, 78, 82, 92, 103, 106, 110-111, 113, 119-120, 138, 141
 Strait 108
 Central 148
 Northern 197, 199
 South 148, 171, 178
 Southern 178
 Western 197-199
 Kangean 111, 118
 Islands 25, 32
 Kapingamarangi 206
 Karimata 106
 Strait 18, 56-57, 113, 122
 Karimun Java 108, 120
 Kedah State 46
 Kelantan 25, 32, 38
 Kembung Island 108
 Keramian 119
 Kokos 124-125, 128
 Komodo Island 119
 Korea 143
 Kota Agung 116-117
 Kotabaru 69, 111 *see also* Laut Island
 Krakatau 84-85, 116-117
 Krawang 25, 32
 Kuala Kurau 25, 28
 Kuala Lumpur 25
 Kupang 119
 Kuta Beach 115
 Kutanihong 124-125, 128
 Kuwait 39
 Labuhan 116-117
 Lombok 133
 Lampung 63, 65
 Bay 116-117
 Langkawi 25, 34
 Langsa 160
 Laurot Islands 110-111
 Laut Island 111 *see also* Kotabaru
 Lombren Island 119
 Lombok 7, 119, 131, 133-134
 Basin 8
 Strait 118
 London 182
 Lord Howe Islands 206, 210
 Luhu 169
 Madagascar 8, 11-12, 137, 141, 143, 158, 173, 178, 182, 184, 186
 Madura 25, 27, 63, 66, 108-111, 118, 139, 143, 145, 160, 162-163, 165, 171, 175, 182, 186, 201, 203, 205-206, 208, 210, 212
 Strait 27, 37, 65, 108, 111
 Makassar 7, 32, 139, 145, 148, 163, 171, 186, 201, 208, 210, 212
 Strait 7-9, 17, 64, 108, 111, 116, 119-122
 Malacca 17, 25-26, 29
 Strait (of) 17-18, 23-24, 26, 36-38, 40-42, 44-46, 48-49, 51-53, 56, 103-104, 108, 162, 197, 201-202
 Malay(an) Peninsula 25-26, 28, 30-31, 33-34, 38-39, 57
 States, Federated (FMS) 34, 38-39
 Malaya 24, 30, 32, 34-39, 46
 British 33, 39
 Federation of 36-38
 Malaysia 16, 24-25, 32, 37-38, 41, 52, 78, 103, 124, 139, 148
 Peninsular 39, 45-46
 Mandar 111
 Mangkok 32
 Manila 105, 120
 Mariana Trench 8
 Marianas 206
 Marquesas Island 156
 Masalembo 108, 119-120
 Mata Siri 119, 120
 Mataram 133
 Mauritius 178, 182, 186
 Medan 25, 28
 Mediterranean 151-152, 193, 201, 204, 206
 Mega 85
 Melanesia 201
 Mentawai Islands 85
 Merak 113, 116-117
 Mergui Archipelago 25, 34
 Mexico 130, 160, 165
 Micronesia 201, 206, 210
 Mindanao 7
 Mindanao Strait 9
 Moçambique Channel 11-12
 Molucca Strait 17 *see* Malacca Strait
 Mombasa 77
 Muncar 102, 107, 114-116
 Muntjar 122, 145
 Namibia 11-12
 Negara 116
 Netherlands Indies 24, 28-29, 33-35, 38-39
 New Caledonia 162, 165, 169, 171, 180, 184, 193, 195
 New Guinea 57, 82, 132, 137
 New Guinea 54, 90
 Nias 82, 85-86, 124, 128
 Nusa Barung 117-118
 Nusa Tenggara 102, 104, 118
 Barat 133
 Oceania 188
 Okinawa 137, 154, 158, 182, 186
 Oman 92
 Gulf of 141
 Ombai Strait 7
 P.P. Karimunjawa 119
 Lima 119
 Pacific 10, 191
 Basin 10
 Ocean 6-13, 17, 65, 118, 160
 Central 156
 Eastern 160
 Indo-West 137, 143, 150, 151-152, 169, 173, 182, 184, 186, 193, 199, 201, 203, 208, 214

North 2
 West 7, 10
 Western 145, 160, 167, 197
 Central 54, 82, 90, 178, 180
 Pacitan 118
 Padang 78
 Pakistan 92, 139
 Pang Pang Bay 114-115
 Pangkor 25, 28, 34, 37
 Panjang 116
 Papagayo, Gulf of 130
 Papua New Guinea 12-13, 141
 Pare Pare 111
 Pejantan 119
 Pekalongan 25, 31, 119-121, 138, 160, 165, 202
 Pelabuhan Ratu 117-118
 Pemalang 120
 Penaitan 116
 Penang 25, 28-29, 31, 33-34, 37, 39, 46
 Pengambangan 114, 116
 Perak 34
 Persian Gulf 78, 104, 139, 143, 171, 180, 182, 208
 Peru 6, 23, 160, 165
 Central 5
 Northern 5
 Perupok 25, 32, 36
 Philippine Trench 8
 Philippines 7, 16-17, 34, 57, 92, 101, 103, 130, 139, 141, 145, 167, 171, 206
 Phuket 145
 Pini 124, 128
 Plantation District 31
 Pontianak 25, 27
 Princess of Orange Bank 115
 Probolinggo 25, 31, 108
 Pulau Ketam 25, 28
 Pulau Laut Kecil 110, 120
 Pulau Pini 85
 Pulau-Pulau Seribu 113 *see also* Thousand Islands
 Puput 141
 Queensland 175
 Raas Strait 108
 Ragay Gulf 92, 101, 130
 Rapa 165
 Red Sea 137, 143, 148, 151-152, 154, 169, 171, 178, 180, 186, 193, 201, 204, 206, 210, 214
 Rembang 119-120
 Réunion 182
 Islands 186
 Riau 40-45, 120, 143, 178
 Lingga Archipelago 25, 34
 Riouw 148, 169
 Rokan 28-29, 37-38
 Rome 77, 105, 277
 Ryukyu Islands 33, 167, 182, 191, 201
 Ryukyus 206
 Sabang 124, 128
 Samar Sea 92, 101, 130
 Samoa 201, 206, 210
 Sampit Bay 111-110, 148
 San Bernardino Strait 7
 Saparua 169
 Sapekan 25, 32
 Sarawak 25, 27, 39, 197
 Savu Basin 8
 Sawu Sea 7, 118
 Selat Panjang *see* Brouwerstraat
 Semangka Bay 116-117
 Semarang 25, 28, 31, 49, 103-104, 108, 110, 167
 Seribu Island 162
 Seychelles 182, 184, 186, 201
 Siam 24, 29-30, 32 *see also* Thailand
 Gulf of 26, 34, 54 *see also* Thailand, Gulf of
 Siberut 124-125, 128
 Sibolga 124, 128
 Sierra Leone 6, 283
 Simeulue 124, 128
 Singapore 17, 24-26, 28-31, 33-34, 36, 39, 52, 103, 113
 Singkibaru 124, 128
 Somalia 11-12, 92, 124, 139
 South Seas 33
 Southern Ocean 12
 Spratly Islands 101
 Sri Lanka 154, 158, 167, 178, 180, 182, 184, 186
 St. Helena 204
 Straits Settlements 25, 38-39
 Sudan 137
 Port 180
 Suez Canal 151-152, 193, 201, 204, 206
 Sulawesi 103, 111
 South 139, 143, 145, 148, 163, 171, 175, 180, 182, 186, 201, 208, 210, 212
 Sulu Sea 17
 Sumatra 13, 16-17, 24-25, 27-31, 34, 38, 40, 52-53, 56-58, 62-66, 68-70, 72, 74, 77-78, 82, 85-92, 99-100, 103, 106, 112-113, 115-120, 122, 123-124, 127-128, 130-131, 160, 191, 202, 282
 Shelf 87
 Central 165
 East 143, 163, 210
 North 41-45
 Northwestern 277
 South 148
 Sumba 118, 131
 Sumbawa 107, 119, 131, 133
 Sunda
 Islands 100, 197
 Lesser 16, 102
 Shelf 7-8, 17, 46, 48-49, 62, 108, 111
 Strait 17-18, 23, 63, 85, 102-103, 106, 115-117
 Surabaya 25, 27-28, 31, 108, 118
 Surigao Straits 7
 Tafelhoek 115
 Tahiti 186
 Taiwan 137, 143, 145
 Tambelan Islands 25
 Tampanuli 27
 Tanah Bala 85, 124, 128
 Tanah Masa 124, 128
 Tanimbar Islands 7
 Tanjung Pura 25, 27
 Tanjung
 Pinang Harbour 50
 Priok 103
 Selatan 56, 60, 171
 Tanzania 12, 177
 Tegal 25, 32, 102, 103, 113, 119-121, 162
 Tehuantepec, Gulf of 130
 Teluk Betung 116-117
 Tg. Antab 114
 Thailand 24, 31, 37, 39, 45, 49, 65, 145
 Gulf of 2, 6-7, 14, 23, 30, 37, 39, 47, 54, 62, 67, 72, 76, 90, 130, 154, 197
 Thousand Islands 25, 34, 113,
 Timor 7, 118-119, 131
 Sea 7-9
 Trench 7
 Torres Strait 7
 Trengganu 24-25, 27-28, 30-32, 38-39
 Tuangku 124, 128
 Tuban 25, 31
 Ujung Kulon 116
 Ujung Pandang 111
 Veth 27-28
 Vietnam 31
 Wahai 169
 Walvis Bay 11
 Western Archipelago 24-28, 30, 33, 35, 37-38
 Yemen 92
 Zanzibar 158, 182

Species Index***

- Abalistes* sp. 99
Abalistes stellaris 4, 57
Abalistes stellatus 59, 95, 126, 129, 137, 214-215 *see also* Balistidae
- Acanthocybium* sp. 116 *see also* Wahoo
Acanthocybium solandri 107
- Acanthurids(-ae) 85, 95
Acanthurus bleekeri 220
Acanthurus mata 95, 220
- Acropomatidae 96
- Aetobatus* sp. 95
Aetobatus narinari 95
- Ajapu 139 *see* *Dussumieria acuta*
- Ajong-ajong 205 *see* *Scomberomorus guttatus*
- Alepes* spp. 57, 108
Alepes djedaba 4, 126
Alepes kalla 4
- Alopias* sp. 96, 98-99
- Alopiidae 96
- Alu-alu 206-207 *see* *Sphyræna obtusata*
- Amblygaster clupeioides* 137 *see also* Clupeiidae
Amblygaster leiogaster 137 *see also* Clupeiidae
Amblygaster sirm 108, 113, 121, 137-139 *see also* Sardinella, spotted
- Anadontostoma chacunda* 4, 43, 57, 109
- Anchisomus multistriatus* 220
- Anchoveta 5-6, 23 *see* *Engraulis ringens*
- Anchovies 120, 134, 204 *see* Engraulidae
- Anthias fasciatus* 220
Anthias squamipinnis 220
- Apogon 49
ceramensis 49
darnleyensis 220
fasciatus 220
quadrifasciatus 220
timorensis 220
- Apogonidae 109
- Aprion virescens* 95, 137, 190-191
- Argyrosomus* sp. 220
- Ariidae 4, 43, 51, 57, 59, 69-73, 95, 108-109, 134, 137
- Ariomma brevimanum* 220
Ariomma brevimanus 220
Ariomma indica 220
Ariomma indicum 220
- Arius* spp. 95, 111 *see also* Catfish
Arius coelatus 57
Arius maculatus 57
Arius thalassinus 57, 59, 95, 126, 220
Arius venosus 57
- Arothron* sp. 220
Arothron caeruleopunctatus 220
Arothron leopardus 49
Arothron manillensis 220
Arothron manillensis 220
- Ariids 59
- Ateleomycterus marmoratum* 220
Ateleomycterus marmoratus 220
- Atrobucca brevis* 220
Atrobucca nibe 99
- Atropus atropus* 4, 57
- Atule* spp. 57
Atule mate 107
- Aulopiformes 137
- Auxis* spp. 13, 116, 118 *see also* Tuna; Tuna, frigate; Mackerels, frigate
Auxis thazard spp. 107-108, 133 *see also* Tuna, frigate
- Axinurus thynnoides* 220
- Bali sardinella 102, 105-106, 115-116 *see* *Sardinella lemuru*
- Balistidae 4, 57-59, 69-73, 95, 137
- Banjara 201 *see* *Rastrelliger kanagurta*
- Banjara 201 *see* *Rastrelliger kanagurta*
- Barenose, rippled 188-189 *see* *Gymnocranius grandoculis*
- Barracudas 117-118
Barracuda, obtuse 206-207 *see* *Sphyræna obtusata*
Barracuda, small 117 *see* *Sphyræna flavicauda*
- Barukang 148 *see* *Netuma thalassina*
- Bathycongrus guttulatus* 221
- Bawal 163, 208 *see* *Parastromateus niger*; *Pampus argenteus*
hitam 163-164 *see* *Parastromateus niger*
putih 208-209 *see* *Pampus argenteus*
- Bebete 182 *see* *Leiognathus equulus*
lubangang 175 *see* *Lactarius lactarius*
lumuh 186 *see* *Gazza minuta*
- Beloso 152-153 *see* *Saurida undosquamis*
sirip pendek 150 *see* *Saurida micropectoralis*
- Bengol 162 *see* *Decapterus russelli*
deles 160 *see* *Decapterus macrosoma*
- Bentong 165 *see* *Selar crumenophthalmus*
- Besiko 169 *see* *Diagramma pictum*
- Bete kalo 139 *see* *Dussumieria acuta*
lalaki 145 *see* *Sardinella lemuru*
-bete sulamang 186 *see* *Gazza minuta*
- Bidji angka 195 *see* *Upeneus sulphureus*
- Bigeye(s), red 94, 199-201 *see* *Priacanthus macracanthus*
- Bijinangka 193-194 *see* *Upeneus moluccensis*
- Bondol 178-179 *see* *Leiognathus splendens*
- Bonito 116, 120 *see* *Sarda orientalis*
- Bothidae 134
- Botto-botto 201 *see* *Rastrelliger kanagurta*
- Brachiurans (crabs) 50
- Branchiostegus australiensis* 220
Branchiostegus gloerfelti 220
Branchiostegus sp. 220
Branchiostegus sp. cf. *argentatus* 220
Branchiostegus sp. cf. *doliatus* 220
- Bream(s), palefin threadfin 113, 197-198 *see* *Nemipterus thosaporni*
- Bregmaceros* sp. 117
- Bulus 160 *see* *Decapterus macrosoma*
blangseng 160 *see* *Decapterus macrosoma*
lake 201 *see* *Rastrelliger kanagurta*
- Bun bun 165 *see* *Selar crumenophthalmus*
- Caelorinchus maculatus* 220
- Caesio* spp. 43 *see also* Lutjanidae
Caesio cuning 108
Caesio erythrogaster 4
- Caesionidae 26, 33, 95 *see also* Fusilier
- Carangidae 4, 43-44, 57, 59, 69-72, 95-96, 134, 137
- Carangids 59, 116, 204 *see* *Megalaspis cordyla*; Carangidae
- Carangoides* spp. 4, 57, 95
Carangoides armatus 126
Carangoides malabaricus 95, 126-127, 129, 137, 154-155 *see also* Trevally, Malabar
- Caranx* spp. 57, 107-108
Caranx heberi 220
Caranx ignobilis 137, 156-157 *see also* Trevally, giant
Caranx kleinii 220
Caranx para 220
Caranx radiatus 220
Caranx sem 220
Caranx semifasciatus 95

***Excludes the species in the list from p. 222 to 274

- Caranx sexfasciatus* 95
Caranx tille 95, 137, 157, 159 *see also* Trevally, tille
Carcharhinus sp. 95
Carcharhinus sealei 57, 59, 126
Carcharias taurus 220
Catfish 111 *see Arius* spp.
 giant 148-149 *see Netuma thalassina*
Centrolophidae 96
Centrophorus sp. 126-127, 129 *see also* Shark, deepwater
Cephalopholis boenack 220
 Cephalopholis boenak 220
Cephalopods 93
Ceratium 137
Chauliodus 85
 Chauliodus sloanei 220
 Chauliodus sloani 220
Cheilinus bimaculatus 220
Chiloscyllium griseum 220
 Chiloscyllium hasseltii 220
Chirocentridae 4, 57, 59, 69-72
Chirocentrus spp. 70
 Chirocentrus dorab 4, 43, 51, 107-108
Chlorophthalmus agassizi 126-127, 129 *see also* Greeneyes
Cladocerans 201
Clupeids 59, 111, 145, 204
Clupeidae 4, 43, 51, 57-59, 69-73, 95, 137
Clupeiformes 137
Coelorinchus maculatus 220
Coilia dussumieri 57
Congrogadoides spinifer 220
Congrogadus spinifer 220
Cookeolus boops 220
 Cookeolus japonicus 220
Coryphaena hippurus 107
Crabs 50, 57, 59, 69-72, 148, 190 *see also* Brachiurans
Croakers 116, 118
Crocodiles 28
Crustaceans 43, 50, 93
 decapod 50
Ctenophores 208
Cubiceps whiteleggi 220
Cuttlefish 57, 59, 69-73, 109
Cynoglossus sp. 220
Cypselurus spp. 111
Da-tanda 171 *see Pomadasys argenteus*
Dactyloptena orientalis 126
Damselshes 275
Dasyatidae 4, 57, 59, 95
Dasyatis sp. 94-95, 98-99
 Dasyatis kuhlii 95
Da-tanda 171-172 *see Pomadasys argenteus*
Dawah 208 *see Pampus argenteus*
Dawahan 208 *see Pampus argenteus*
Decapterus spp. 43, 46, 57, 105-106, 108-109, 116-117, 119, 122 *see also*
 Scads, layang; Mackerels, chub
 Decapterus kurroides 96, 98-99
 Decapterus macrosoma 96, 99, 120-121, 279, 137, 160-161, 279 *see also*
 Scad, shortfin
 Decapterus maruadsi 95
 Decapterus russelli 46, 99, 121, 137, 162-163 *see also* Scad; Scad,
 Indian
 Decapterus tabl 99
Deles 160 *see Decapterus macrosoma*
Dentex tumifrons 220
Dero 141 *see Pellona ditchela*
Diagramma pictum 95, 126, 137, 169, 170-171 *see also* Sweetlips, painted
Diatoms 201
Dibas 163 *see Parastromateus niger*
Diodon sp. 113 *see also* Porcupine fish
 Diodon hystrix 96, 99
Diodontidae 96, 108-109
Djambon 210 *see Terapon jarbua*
Djambon 210 *see Terapon jarbua*
Djambung 210 *see Terapon jarbua*
Djandjan 210 *see Terapon jarbua*
Djangdjan 210 *see Terapon jarbua*
Djapuh 139 *see Dussumieria acuta*
Djogor 212 *see Trichiurus lepturus*
Djurung 143 *see Sardinella gibbosa*
Djuwi djuwi 143 *see Sardinella gibbosa*
Doderleinia berycoides 220
Dodok 178, 182 *see Leiognathus splendens*
Doederleinia berycoides 220
Dolphinfish 117-118
Domul 169 *see Diagramma pictum*
Dorang 163 *see Parastromateus niger*
 hitam 163 *see Parastromateus niger*
Drepane longimana 4
 Drepane punctata 95
Drepanidae 4, 57-59, 69-73, 95
Duri padi 148 *see Netuma thalassina*
Duri utek 148 *see Netuma thalassina*
Dussumieria sp. 95
 Dussumieria acuta 4, 57-59, 95, 108-109, 137, 139-141 *see also*
 Sardine, rainbow
 Dussumieria elopsoides 95, 139
 Dussumieria hasselti 49
Echinoderms 50
Echinoideans 50
Elagatis bipinnulata 107-108, 117 *see also* Runner, blue
Encrasicholina devisi 221
 Encrasicholina punctifer 221
Engraulidae 4, 43, 51, 57-59, 69-72, 95, 109, 134 *see also* Anchovies
Engraulids 59, 111 *see also* Engraulidae
Engraulis ringens 5 *see also* Anchoveta
Epinephelinae 218
Epinephelus sp. 218, 220
 Epinephelus diacanthus 218, 220
 Epinephelus epistictus 220
 Epinephelus faveatus 220
 Epinephelus magniscuttis 218, 220
 Epinephelus microdon 218, 220
 Epinephelus morrhua 220
 Epinephelus poecilnotus 218, 220
 Epinephelus polyphkadion 220
 Epinephelus quoyanus 218, 220
 Epinephelus stictus 220
 Epinephelus tauvina 95, 98-99
Erong-erong 210 *see Terapon jarbua*
Etelis sp. 96, 98-99
Eugomphodus taurus 220
Euthynnus sp. 116 *see also* Tuna
 Euthynnus affinis 107-108, 133 *see also* Kawakawa
Feroxodon multistriatus 220
Fistularia spp. 109, 126
Fistulariidae 4
Flatfishes 43
Flyingfish 111, 116, 118-119
Formio niger 4, 43, 57, 59, 69-72
Formionidae 4, 57, 109
Fusiliers 26, 33-34 *see* Caesionidae
Gadji 169 *see Diagramma pictum*
Gadji-gadji 169 *see Diagramma pictum*
Gadjih 169 *see Diagramma pictum*

Galeocerdo cuvier 220
 Garut 171 see *Pomadasys argenteus*
Gazza minuta 57, 95, 98-99, 126, 129, 137, 186-187
 Gebel 163 see *Parastromateus niger*
 Gempar 178 see *Leiognathus splendens*
 Gemper 178 see *Leiognathus splendens*
 Gempylids 85
 Gerot-gerot 171, 173-174 see *Pomadasys argenteus*; *Pomadasys maculatus*
 Gerreidae 4, 43, 57, 59, 69-72, 95, 109, 134, 137
Gerres spp. 49
 Gintong 165 see *Selar crumenophthalmus*
Glaucosoma sp. 96, 99
 Glaucosomatidae 96
Gnathanodon speciosus 96, 99
 Goatfish 113
 sulphur 195-196 see *Upeneus sulphureus*
 goldfish 193-194 see *Upeneus moluccensis*
 Gombong 201 see *Rastrelliger kanagurta*
 Gonostomatidae 85
 Greeneyes 127 see *Chlorophthalmus agassizi* spp.
 Groupers 275
 Grunts 116
 Grunt, saddle 173-174 see *Pomadasys maculatus*
 Grunt, silver 171-173 see *Pomadasys argenteus*
 Gugup 148 see *Netuma thalassina*
 Gungut 148 see *Netuma thalassina*
Gymnocranius grandoculis 96, 99, 126, 129, 137, 188-189, 220
 Gymnocranius robinsoni 220
Gymnothorax thyrsoideus 220
 Haemulidae 95, 137
 Hairtail 127 see *Trichiurus lepturus* spp.
 largehead 212-213 see *Trichiurus lepturus*
 Hajam 167 see *Pentaprion longimanus*
 Halfbeaks 116
Harpadon nehereus 126
 Harpadontidae 109
 Herring 3
Heteropriacanthus cruentatus 221
 Heterosomata 57, 69-73
Hexanchus griseus nakamurai 220
 Hexanchus nakamurai 220
Himantura sp. 220
 Himantura uarnak 95, 98-99, 220
Hologymnosus sp. 220
 Hologymnosus rhodonotus 220
Hyperoglyphe sp. 96
Ilisha spp. 4, 57-59, 95
 Ilisha elongata 95
Inegocia bosschei 220
 Inegocia japonica 220
 Inegocia japonicas 220
 Japuh 139, 141 see *Dussumieria acuta*
 Jarbua terapon 210-211 see *Terapon jarbua*
 Jellyfish 126
 Jobfish, green 190-191 see *Aprion virescens*
 Jobfish, sharptooth 191-193 see *Pristipomoides typus*
Johnius coitor 220
 Johnius laevis 220
 Kaban bulus 160 see *Decapterus macrosoma*
 laes 160 see *Decapterus macrosoma*
 padara 160, 162 see *Decapterus macrosoma*; *Decapterus russelli*
 patek 160, 162 see *Decapterus macrosoma*; *Decapterus russelli*
 Kadji 169 see *Diagramma pictum*
 Kaji 169-171 see *Diagramma pictum*
 Kakunir 195 see *Upeneus sulphureus*
 Kambing-kambing 214-215 see *Abalistes stellatus*
 Kandibas 163 see *Parastromateus niger*
 Kapet 163 see *Parastromateus niger*
 Karang besar 156-157 see *Caranx ignobilis*
 Karang tile 158-159 see *Caranx tile*
 Karang trevali 154-155 see *Carangoides malabaricus*
 Karong-karong 210 see *Terapon jarbua*
 Katji 169 see *Diagramma pictum*
 Katji-katji 169 see *Diagramma pictum*
Katsuwonus pelamis 11, 13, 107 see also Skipjack
 Kawakawa 133 see *Euthynnus affinis* spp.
Kembong 29, 34 see *Rastrelliger* spp.
 Kembung lelaki 201-203 see *Rastrelliger kanagurta*
 Kempar 186 see *Gazza minuta*
 Kempor labu 186 see *Gazza minuta*
 Keretang 210 see *Terapon jarbua*
 Kerong-kerong tambu 210-211 see *Terapon jarbua*
 Kerongan 210 see *Terapon jarbua*
 Kerot-kerot 171, 173 see *Pomadasys argenteus*; *Pomadasys maculatus*
 Kerung-kerung 210 see *Terapon jarbua*
 Kibas 163 see *Parastromateus niger*
 Klemes 175 see *Lactarius lactarius*
 Korok 162 see *Decapterus russelli*
 Kotjol 206 see *Sphyræna obtusata*
 Krokot 171 see *Pomadasys argenteus*
 Krot 171-173 see *Pomadasys argenteus*; *Pomadasys maculatus*
 -krot 171, 173 see *Pomadasys argenteus*; *Pomadasys maculatus*
 Kunir 195-196 see *Upeneus sulphureus*
 Kuniran 195-196 see *Upeneus sulphureus*
 Kurisi 197-198 see *Nemipterus thosaporni*
 Kutjul 206 see *Sphyræna obtusata*
 Kuwe 154 see Trevally, Malabar; *Carangoides malabaricus*
 Lactariidae 57, 59, 95, 137
Lactarius lactarius 59, 69-73, 95, 99, 126-127, 129-130 137, 175-176 see
 also Trevally, false
 Ladjang 162 see *Decapterus russelli*
 Ladjeng 162 see *Decapterus russelli*
 lakek 160 see *Decapterus macrosoma*
 Ladjur 212 see *Trichiurus lepturus*
 Ladjuru 212 see *Trichiurus lepturus*
 Lagocephalidae 4
 Lajang 160, 162 see *Decapterus macrosoma*; *Decapterus russelli*
 deles 160 see *Decapterus macrosoma*
 lidi 160 see *Decapterus macrosoma*
 Lajur 212 see *Trichiurus lepturus*
 Lampa 148 see *Netuma thalassina*
 Langsar 206 see *Sphyræna obtusata*
 Langung 203, 205 see *Scomberomorus commerson*; *Scomberomorus guttatus*
 Lantern fish 116 see *Myctophidae*
 Lawang 208 see *Pampus argenteus*
 Layang 26, 31, 162-163 see *Decapterus russelli*
 deles 160-161 see *Decapterus macrosoma*
 Layur 212-213 see *Trichiurus lepturus*
 Leatherjackets 113, 117
 Leiognathidae 4, 43-44, 50-51, 54, 57, 59, 61, 69-73, 95, 109, 117, 119,
 123, 127, 134, 137, 177
 Leiognathids 59, 111, 113, 116, 118, 128-130, 177 see *Leiognathidae*
Leiognathus sp. 95, 98-99, 208
 bindus 4, 57, 94-95, 98-100, 126, 129, 137, 179, 181
 brevirostris 57
 daura 57
 elongatus 57, 95, 98-99
 equulus 57, 94-95, 98-100, 126, 129, 137, 182-183, 220
 fasciatus 95, 126, 129
 leuciscus 4, 126, 137, 184-185
 smithursi 57
 splendens 4, 51, 57-59, 95, 98-99, 126, 129, 137, 177-179

- Lelemah 175 *see* *Lactarius lactarius*
 Lemah 175 *see* *Lactarius lactarius*
 Lemahan 175 *see* *Lactarius lactarius*
 Lemuru 83, 102, 122, 145, 147, 278 *see* Sardine, oil; *Sardinella lemuru*
 Leptocephali 85
Lepturacanthus savala 4, 95, 98-99
Lestidiops mirabilis 220
Lestrolepis intermedia 220
 Lestrolepis japonica 220
 Lethrinidae 95, 134, 137
Lethrinus atkinsoni 220
 Lethrinus choerorhynchus 220
 Lethrinus elongata 220
 Lethrinus erythropterus 220
 Lethrinus fraenatus 220
 Lethrinus genivittatus 220
 Lethrinus hypselopterus 220
 Lethrinus laticaudis 220
 Lethrinus mahsena 220
 Lethrinus miniatus 95
 Lethrinus nebulosus 95, 220
 Lethrinus nematacanthus 220
 Lethrinus olivaceus 220
 Lethrinus variegatus 95, 99
 Limat 175 *see* *Lactarius lactarius*
 Lizardfish 94, 113, 127 *see* *Saurida undosquamis* spp.
 brushtooth 151-153 *see* *Saurida undosquamis*
 shortfin 150-151 *see* *Saurida micropectoralis*
 Lobster 57, 59, 89 *see* *Thenus* sp. 57
 palinurid 89
 shovelnose 89 *see* *Thenus* sp.
 spiny 89
 Lokmolok 182 *see* *Leiognathus equulus*
 Loliginidae 59, 96
Loligo sp. 96, 98-99, 126-127, 129-130
 Loligoidae 43
 Loncong 167-169 *see* *Pentaprion longimanus*
 Longkerong 210 *see* *Terapon jarbua*
 Longlong mata 141 *see* *Pellona ditchela*
 Lontong 167 *see* *Pentaprion longimanus*
 Lontjong 167 *see* *Pentaprion longimanus*
Lophioides gracilimanus 220
 Lowang 208 *see* *Pampus argenteus*
 Luntju 160 *see* *Decapterus macrosoma*
 Lutjanidae 4, 43-44, 57, 59, 69-72, 95-96, 134, 108-109, 137 *see also* *Caesio*
 spp.
Lutjanidae spp. 123 *see also* Snappers
 Lutjanids 59 *see also* Lutjanidae
Lutjanus sp. 95, 98-99
 Lutjanus argentimaculatus 95, 119
 Lutjanus erythropterus 126
 Lutjanus fulviflamma 220
 Lutjanus fulviflammus 220
 Lutjanus johni 220
 Lutjanus johnii 4, 57, 95, 220
 Lutjanus lineolatus 4
 Lutjanus malabaricus 61, 95, 98-99, 126, 129
 Lutjanus monostigma 220
 Lutjanus monostigmus 220
 Lutjanus rivulatus 95
 Lutjanus sanguineus 4, 57, 59, 95, 129
 Lutjanus timorensis 220
 Lutjanus timoriensis 220
 Lutjanus vitta 220
 Lutjanus vittus 220
 Mackerel 35, 46, 120
 Indian 46, 108, 116-117, 201-203 *see* *Rastrelliger* sp.; *Rastrelliger*
 kanagurta
 Indo-Pacific king 205-206 *see* *Scomberomorus guttatus*
 narrow-barred Spanish 113, 203-204 *see* *Scomberomorus*
 commerson
 Spanish 117-118
 chub 46 *see* *Decapterus* spp.
 frigate 13 *see* *Auxis* spp.
Macroparalepis mirabilis 220
Malakichthys sp. 96, 98-99
Malthopsis lutea 220
 Malthopsis luteus 220
 Mangahua 210 *see also* *Terapon jarbua*
 Mangmung 148 *see* *Netuma thalassina*
 Manjong 148 *see* *Netuma thalassina*
 Manjung 148 *see* *Netuma thalassina*
 Manjung kerbo 148 *see* *Netuma thalassina*
 Manriwasa lelung 163 *see* *Parastromateus niger*
 Manriwasakebo 208 *see* *Pampus argenteus*
 Manyung 148-149 *see* *Netuma thalassina*
 Maos 143 *see* *Sardinella gibbosa*
 Medusae 208
Megalaspis cordyla 4, 57, 107-109, 116, 119-120 *see also* Carangids; Scad,
 hairtail
Mene maculata 95
 Mengantih 171 *see* *Pomadasys argenteus*
 Mengida 143 *see* *Sardinella gibbosa*
 Menidae 95
Metapenaeus monoceros 50
Mobula diabolus 220
 Mobula diabolus 220
 Mojarra, longfin 167-169 *see* *Pentaprion longimanus*
 Molluscs 148
 Molok-molok 182 *see* *Leiognathus equulus*
 Mullidae 4, 57, 59, 69-72, 95, 109, 134, 137
 Mullids 59
Mulloides flavolineatus 220
Mulloidichthys flavolineatus 220
 Muraenesocidae 57, 69-73
 Mursiah 143 *see* *Sardinella gibbosa*
Mustelus sp. 96
 Mustelus griseus 95, 99
 Myctophidae 116-117, 126 *see also* lantern fish
 Myctophids 85, 111 *see* Myctophidae
 Myliobatidae 95
Myliobatis tobije 220
 Myliobatis tobije 220
Myripristis melanosticta 220
 Myripristis melanostictus 220
Naso sp. 95, 98-99, 220
 Naso lopezi 220
 Naso thynnoides 220
 Naso tuberosus 95
Nautilus sp. 99
 Nemipteridae 4, 43, 57, 59, 69-72, 92, 109, 134, 137
Nemipterus sp. 208, 221
 balinensoides 221
 bathybius 4, 57, 197 *see also* Bream(s), palefin threadfin
 gracilis 221
 hexodon 57
 japonicus 4, 46, 57
 marginatus 54, 57, 197, 199-220
 mesoprion 57
 metopias 221
 peronii 57
 tambuloides 129

- thosaporni* 137, 197-200 *see also* Bream(s), palefin threadfin
tolu 57
zysron 221
Netuma thalassina 137, 148-149, 220 *see also* Catfish, giant
 Njiuran 208 *see Pampus argenteus*
 Njor njoran 208 *see Pampus argenteus*
 Ophiuroids 50
Opisthopterus tardoore 126
 Opisthopterus valenciennensis 4
Osteogeneiosus militaris 4, 57
 Ostracods 201
Oxycheilinus bimaculatus 220
 Oysters, pearl 24
Pagrus major 99
 Palinurids 50
Pampus argenteus 4, 43, 57, 95, 137, 208-209 *see also* Pomfret, silver
 Pampus chinensis 4, 57
Pantolabus radiatus 220
Paracaesio sp. 95, 99
 Paracaesio kusakarii 99
Parapenaeopsis spp. 50
Paraplagusia longirostris 220
Parapriacanthus sp. 95
Parascolopsis sp. 221
 Parascolopsis qantasi 221
 Parascolopsis tanyactis 221
Parastromat 59
Parastromateus niger 126, 137, 163-164 *see also* Pomfret, black
Parupeneus sp. 221
 Parupeneus heptacanthus 221
 Parupeneus multifasciatus 221
 Parupeneus pleurospilus 221
 Peda-peda lotong 163 *see Parastromateus niger*
 Peda-peda puti 208 *see Pampus argenteus*
Pellona ditchela 126, 129, 137, 141-142 *see also* Pellona, Indian
 Pellona, Indian 141-142 *see Pellona ditchela*
 Penaeidae 43
Penaeus merguensis 50
 Penaeus semisulcatus 50
 Pentapodidae 4, 57, 59, 69-72
Pentapodus setosus (?) 4
Pentaprion longimanus 4, 49, 57, 59, 61, 95, 98-99, 126, 129, 137, 167-169
 see also Mojarra, longfin
 Penteng 165 *see Selar crumenophthalmus*
 Pentong 165 *see Selar crumenophthalmus*
 Peperek 180-181, 184, 186 *see Leiognathus bindus; Leiognathus leuciscus;*
 Gazza minuta
 bondolan 186-187 *see Gazza minuta*
 Tijina 178, 182 *see Leiognathus splendens; Leiognathus equulus*
 topang 182-183 *see Leiognathus equulus*
 Perciformes 137, 218
 Perek topang 182 *see Leiognathus equulus*
 Peridinium 137
Peristedion sp. 126, 129
 Petah 180, 186 *see Leiognathus bindus; Gazza minuta*
 Petek kuning 182 *see Leiognathus equulus*
Photonectes albipennis 221
 Photonectes albipennis 221
 Phyllosoma 89
 Phytoplankton 201
Pinjalo sp. 221
 Pinjalo lewisi 221
 Pinjalo pinjalo 221
 Platycephalidae 4, 43-44, 109
Plectorhinchus celebecus 221
 Plectorhinchus celebicus 221
 Plectorhinchus chaetodontoides 221
 Plectorhinchus chaetodontoides 221
 Plectorhynchidae 4
Plectorhynchus spp. 43
 Plectorhynchus pictus 4
Plectropomus pessuliferum 221
 Plectropomus pessuliferus 221
 Polychaetes 201
Polydactylus sextarius 221
 Polynemidae 57, 59, 69-73, 109, 134
Polynemus sextarius 126, 221
 Pomadasyidae 4, 43-44
 Pomadasyidae 57, 59, 69-73, 109
Pomadasys sp. 4, 95
 Pomadasys argenteus 95, 98-99, 137, 171-173 *see also* Grunt, silver
 Pomadasys argyreus 57, 59, 126, 129
 Pomadasys hasta 57
 Pomadasys maculatus 4, 95, 98, 126, 129, 137, 173-174 *see also*
 Grunt, saddle
 Pomfret, black 111, 163, 164 *see Parastromateus niger*
 silver 208-209 *see Pampus argenteus*
 Ponyfish(es) 123, 117, 127 *see* Leiognathidae
 common 182-183 *see Leiognathus equulus*
 orange-fin 180-181 *see Leiognathus bindus*
 splendid 178-179 *see Leiognathus splendens*
 whipfin 184-185 *see Leiognathus leuciscus*
 Porcupine fish 113 *see Diodon* sp.
 Porifera 50
 Porpoises 111, 118
 Potean 208 *see Pampus argenteus*
Poterion nautilus 50
 Potian 208 *see Pampus argenteus*
 Prawns 148
 Priacanthidae 4, 43, 57, 59, 70, 95-96, 109, 137
 Priacanthids 85
Priacanthus spp. 69-72, 96, 98-99, 221
 Priacanthus cruentatus 221
 Priacanthus fitchi 199-200, 221
 Priacanthus macracanthus 4, 57, 94, 96, 99, 137, 199, 201 *see also*
 Bigeye(s), red
 Priacanthus sagittarius 221
 Priacanthus tayenus 57, 126, 129
Pristipomoides sp. 95, 99
 Pristipomoides flavipinnis 99
 Pristipomoides typus 95, 126, 129, 137, 191-193
Psenes whiteleggii 220
Psettodes 59
 erumei 43, 57, 59
Pseudanthias fasciata 220
 Pseudanthias squamipinnis 220
Pteragogus flagellifer 221
 Pteragogus flagelliferum 221
Pterocaesio sp. 221
 Pterocaesio tessellata 221
 Pufferfish 111, 117
 Puput 142 *see Pellona ditchela*
 Putih 148 *see Netuma thalassina*
Pyrosoma 85
 Rachycentridae 57, 69-72
Rachycentron canadus 43
 Radja bau 169 *see Diagramma pictum*
Rastrelliger spp. 13, 29, 34, 37, 43-46, 57, 59, 108, 116-117 *see also*
 Kembong; Mackerel, Indian; Mackerels, chub
 Rastrelliger brachysoma 4, 108, 109, 121, 126
 Rastrelliger kanagurta 46, 49, 108-109, 121, 137, 201-203 *see also*
 Mackerel, Indian
 Rays 28, 43, 51, 57, 59, 69-73, 95, 134
 Rentjek bulus 160, 162 *see Decapterus macrosoma; Decapterus russelli*

- kaban 160, 162 *see* *Decapterus macrosoma*; *Decapterus russelli*
 padara 162 *see* *Decapterus russelli*
 patek 162 *see* *Decapterus russelli*
Rhechias guttulatus 221
Rhincodon typus 118 *see also* Sharks, whale
Rhiniodon typus 221
 Rhinobatidae 96
Rhinobatus sp. 96, 98-99
Rhynchobatus djiddensis 221
 Ronga 171 *see* *Pomadasys argenteus*
 Runner, blue 117-118 *see* *Elagatis bipinnulata*
Ryncobatus djiddensis 221
 Saängsa 201 *see* *Rastrelliger kanagurta*
 Salps 85, 208
Sarda sp. 120 *see also* Bonito
 Sarda orientalis 107, 116 *see also* Bonito
 Sardin 137-138 *see* *Amblygaster sirm*
 Sardine 119, 134
 oil 107, 122 *see* *Sardinella longiceps*; Lemuru
 rainbow 139-141 *see* *Dussumieria acuta*
Sardinella spp. 57, 59, 98, 115-116, 120, 122, 132 *see also* Sardines
 Sardinella brachysoma 108
 Sardinella fimbriata 121
 Sardinella fimbriata 143
 Sardinella gibbosa 4, 108-109, 113, 121, 137, 143-144 *see also*
 Sardinella, goldstriped
 Sardinella lemuru 83, 94, 98-99, 102, 106, 108-109, 116, 120, 137,
 145-147 *see also* *Sardinella*, Bali
 Sardinella longiceps 95, 107, 122, 145 *see* Sardine, oil
Sardinella 137
 Sardinella, Bali 145, 147 *see* *Sardinella lemuru*
 Sardinella, goldstriped 143-144 *see* *Sardinella gibbosa*
 Sardinella, spotted 137-139 *see* *Amblygaster sirm*
 Sardines 11-12, 35, 132 *see* *Sardinella* spp.
Saurida sp. 96
 Saurida elongata 4
 Saurida longimanus 4, 57
 Saurida micropectoralis 57, 95, 137, 150-151 *see also* Lizardfish, shortfin
 Saurida tumbil 4, 126-127, 129
 Saurida undosquamis 49, 57, 94, 96, 98-99, 126, 129-130, 137, 151-
 153 *see also* Snapper-lizardfish; Lizardfish, brushtooth
 Sawfish 28
 Scad 43, 46, 116, 120 *see* *Decapterus russelli*; *Selar* spp.
 bigeye 113, 116, 165-167 *see* *Selar crumenophthalmus*
 hardtail 119, 120 *see* *Megalaspis cordyla*
 Indian 162-163 *see* *Decapterus russelli*
 layang 105, 116-117, 119, 122 *see* *Decapterus* spp.
 shortfin 160-161 *see* *Decapterus macrosoma*
 Sciaenidae 4, 43-44, 57-59, 69-73, 109, 118
Scolopsis spp. 43, 57
Scomberoides sp. 4
 Scomberoides commersonianus 107-108
 Scomberomorus commerson 137, 203-204
 Scomberomorus commerson 4, 107
 Scomberomorus guttatus 4, 126, 129, 107-108, 137, 205-206
Scomberomorus spp. 13, 43, 57, 109
 Scombridae 4, 57, 59, 69-73, 137
 Scombrids species 13, 120 *see* *Rastrelliger* spp.
Scorpaena bynoensis 221
Scorpaena picta 221
 Scyllariidae 59
 Sea-snakes 28
 Sea-stars 50
 Seaweed 24
Sebastes tertius 221
Sebasticus tertius 221
Secutor insidiator 4, 57, 126
 Secutor ruconius 4, 95, 99
Selar spp. 57, 116, 108-109, 121 *see also* Scad, bigeye
 Selar bentong 165-167 *see* *Selar crumenophthalmus*
 Selar crumenophthalmus 95, 98, 121, 126, 137, 165-167 *see also*
 Scad, bigeye
Selaroides leptolepis 4, 57, 59
 Sepiidae 59
 Sepiolidae 43
Seriola rivoliana 99
Seriolina nigrofasciata 4, 57, 108
 Serranidae 43-44, 57, 59, 69-72, 95, 218
Setipinna spp. 57
 Shark 27-28, 43, 51, 57, 59, 69-73, 95, 108, 111, 132, 134
 deepwater 127 *see* *Centrophorus* sp.
 whale 118 *see* *Rhincodon typus*
 Shrimp 25-26, 30, 38-39, 57, 59, 69-73, 111, 117, 119, 129, 190, 201
 mantis 50, 148 *see* Stomatopods 50
 deep sea (*Solenocera*?) 89
 penaeid 50, 109, 204
 red 111, 116-117
Siderea thyrsoides 220
 Siganidae 44
Siganus spp. 43
Sillago sp. 221
 Sillago burrus 221
 Sillago ingenuua 221
 Sillago maculata burrus 221
 Sillago vittata 221
 Siluriformes 137
 Sintring 143 *see* *Sardinella gibbosa*
 Skipjack 11, 13, 117-118, 120, 133 *see* Tuna, skipjack; *Katsuwonus*
 pelamis
 Snapper-lizardfish 129 *see* *Saurida undosquamis*
 Snappers 119, 123, 130 *see* Lutjanidae
Solenocera spp. 50
 Soroi 145 *see* *Sardinella lemuru*
 Sparidae 134
Sphyræna spp. 43
 Sphyræna barracuda 57, 126
 Sphyræna flavicauda 117 *see also* Barracuda, small
 Sphyræna jello 57
 Sphyræna obtusata 4, 95, 98, 126, 137, 206-207 *see also* Barracuda,
 obtuse
 Sphyrænidae 4, 57, 59, 69-72, 95, 107-109, 137
 Sphyrænids 59
 Sponges 3, 50, 57, 59, 61
 Squalidae 96
Squalus sp. 96, 99
 Squids 3, 57, 59, 69-72, 85, 87, 130, 109, 111, 116-117, 119, 204
Squilla species 148
Stegostoma fasciatum 221
Stegostoma varium 221
Stolephorus spp. 4, 57, 59, 95, 108
 Stolephorus devisi 221
 Stolephorus indicus 95
 Stolephorus insularis 95
 Stolephorus punctifer 221
 Stomatopods 50 *see also* Shrimp, mantis
 Stromateidae 4, 57, 59, 69-72, 95, 109, 137
Strophidon sathete 221
Suggrundus bosschei 220
 Susu 175-176 *see* *Lactarius lactarius*
 Swaggi 199-201 *see* *Priacanthus macracanthus*
 Sweetlips, painted 169-171 *see* *Diagramma pictum*
 Synodontidae 4, 43, 57, 59, 69-72, 95-96, 109, 134, 137
Synodus englemani 221
 Synodus variegatus 221

- Taius tumifrons* 220
 Tamban 143 see *Sardinella gibbosa*
 sisik 143 see *Sardinella gibbosa*
 Tambi-tambi 175 see *Lactarius lactarius*
 Tana 175 see *Lactarius lactarius*
 Tandjan 143 see *Sardinella gibbosa*
 Tangkolok 208 see *Pampus argenteus*
 Tantjak 206 see *Sphyræna obtusata*
 Temban montjo 145 see *Sardinella lemuru*
 Tembang 143-144 see *Sardinella gibbosa*
 djawa 139 see *Dussumiera acuta*
 djuwi 143 see *Sardinella gibbosa*
 lakara 143 see *Sardinella gibbosa*
 mata kutjing 145 see *Sardinella lemuru*
 montjong 145 see *Sardinella lemuru*
 rakapeng 139 see *Dussumieria acuta*
 Tengere 203, 205 see *Scomberomorus commerson*; *Scomberomorus guttatus*
 Tenggara 205-206 see *Scomberomorus guttatus*
 papan 203-204 see *Scomberomorus commerson*
 Tengiri 203, 205 see *Scomberomorus commerson*; *Scomberomorus guttatus*
Terapon jarbua 95, 137, 210-211
Terapon theraps 126, 129
 Terapontidae 4, 43, 57, 59, 69-72, 95, 109, 137
 Tetraodontiformes 137
 Tetraodontidae 109
Thenus sp. 57, 89 see also Lobsters, shovelnose
 Thenus orientalis 43, 50, 69-73, 126
Therapon sp. 4
Thyrsitoidea marleyi 221
Thryssa spp. 57
Thunnus alalunga 107
 Thunnus albacares 11, 107 see also Tuna, yellowfin
 Thunnus obesus 13, 107 see also Tuna, bigeye
 Thunnus tonggol 13, 107 see also Tuna, longtail
Thyrsitoidea sp. 126, 129
Thyrsoidea macrura 221
 Tjalong 203, 205 see *Scomberomorus commerson*; *Scomberomorus guttatus*
 Tjangetjang 203, 205 see *Scomberomorus commerson*; *Scomberomorus guttatus*
 Tjapo 139 see *Dussumieria acuta*
 Tjaria 180 see *Leiognathus bindus*
 Tjeplak 208 see *Pampus argenteus*
 Tjeplek 163 see *Parastromateus niger*
 Tjiro 143 see *Sardinella gibbosa*
 Tong gentong 165 see *Selar crumenophthalmus*
 Toothpony 186-187 see *Gazza minuta*
 Torpedo-fish 28
 Totata 143 see *Sardinella gibbosa*
 Towoito 171 see *Pomadasys argenteus*
 Trash sp. 96
 Trevally, false 127, 130, 175-176 see *Lactarius lactarius*
 giant 156-157 see *Caranx ignobilis*
 Malabar 154-155 see *Carangoides malabaricus*
 tille 157, 159 see *Caranx tille*
 Triacanthidae 4
 Triakidae 95-96
 Trichiuridae 4, 43, 57, 59, 69-72, 95, 109, 134, 137
Trichiurus sp. 99
 Trichiurus lepturus 4, 95, 98-99, 107, 126-127, 129, 137, 212-213 see also Hairtail
 Triggerfish, starry 214-215 see *Abalistes stellatus*
Tripang 24, 27
 Tuna 3, 11, 13-14, 111, 116, 119, 120 see *Auxis* sp.; *Euthynnus* sp.
 bigeye 13, 107 see *Thunnus obesus*
 frigate 118, 120, 133 see *Auxis* sp.; *Auxis thazard*
 jumping 118
 little 118, 120
 longtail 13, 111, 113 see *Thunnus tonggol*
 skipjack 117-118, 120
 yellowfin 11, 111, 118, 120 see *Thunnus albacares*
 Turtles 57
Upeneus spp. 49, 99
 Upeneus bensasi 57
 Upeneus moluccensis 95, 126, 137, 193-194
 Upeneus sulphureus 4, 57, 59, 126, 129, 137, 195-196
 Upeneus vittatus 95, 98-99
 Usek-usek 205 see *Scomberomorus guttatus*
 Utik 148 see *Netuma thalassina*
 Wahoo 116 see *Acanthocybium* sp.
 Whales 24, 119
Xanthichthys caeruleolineatus 221
 Zooplankton 201, 208

Baseline studies of biodiversity: the fish resources of western Indonesia.
D. Pauly and P. Martosubroto, Editors. 1996. ICLARM Stud. Rev. 23, 312 p.
US\$16.50 surface, \$31.50 airmail, P345.

TITLES OF RELATED INTEREST

- An annotated bibliography of slipmouths (Pisces: Leiognathidae). D. Pauly and S. Wade-Pauly. 1981. ICLARM Bibliogr. 2, 62 p. US\$3 surface, \$7 airmail, P50.
- An atlas of the growth, mortality and recruitment of Philippine fishes. J. Ingles and D. Pauly. 1984. ICLARM Tech. Rep. 13, 127 p. US\$6.50 surface, \$11 airmail, P90.
- Growth, mortality and recruitment of commercially important fishes and penaeid shrimps in Indonesian waters. A. Dwiponggo, T. Hariati, S. Banon, M.L. Palomares and D. Pauly. 1986. ICLARM Tech. Rep. 17, 91 p. US\$4.50 surface, \$8.50 airmail, P70.
- Indonesian marine capture fisheries. C. Bailey, A. Dwiponggo and F. Marahudin. 1987. ICLARM Stud. Rev. 10, 196 p. US\$9.50 surface, \$18 airmail, P170.
- Trophic models of aquatic ecosystems. V. Christensen and D. Pauly, Editors. 1993. ICLARM Conf. Proc. 26, 390 p. US\$18 surface, \$36 airmail, P400.
- FishBase 96: concepts, design and data sources. R. Froese and D. Pauly, Editors. 1996. ICLARM. 179 p. Distributed with one CD-ROM disk for US\$95 (airmail). Available free of cost for collaborators of the FishBase project. Available only from the FishBase Project, ICLARM, MCPO Box 2631, 0718 Makati City, Philippines.
- The database mentioned in the Torres et al. paper, this volume, on p. 276-283, is available on diskettes. Diskette No. 1 (catch data) and Diskette No. 2 (length-frequency data) are available for US\$20 each, including handling and mailing. If both are ordered, their price is US\$30. Order directly from ICLARM, Manila. See address below.

How to Order

Enquirers from the USA, Canada and Central and South America should order directly from our exclusive distributor, **International Specialized Book Services (ISBS)**, 5804 N.E. Hassalo St., Portland, Oregon 97213-3644, to whom payment should be directed using the **airmail** price. ISBS telephone: 800-944-6190; e-mail: orders@isbs.com.

Enquirers from the Philippines may use the peso price and order directly from ICLARM. Enquirers elsewhere should address orders and payment (payable to ICLARM) in US\$ by bankdraft or UNESCO coupons to: The Editor, ICLARM, MCPO Box 2631, 0718 Makati City, Philippines. (ICLARM e-mail address: iclarm@cnet.com; fax: (63-2) 816-3183). We can accept US\$ checks only if from a US-based bank due to high clearance fees of other banks.

Airmail is strongly recommended.

ICLARM ORGANIZATIONAL STATEMENT

"For those who use and depend on fish and aquatic life in the developing world"

ICLARM's VISION

Our Goal : To enhance the well-being of present and future generations of poor people in the developing world through production, management and conservation of living aquatic resources.

Our Objectives : Through international research and related activities, and in partnership with national research and other institutions, to:

1. Improve the biological, socioeconomic and institutional management mechanisms for sustainable use of aquatic resource systems.
2. Devise and improve production systems that will provide increasing yet sustainable yields.
3. Help develop the capacity of national programs to ensure sustainable development of aquatic resources.

The Functions of ICLARM are to:

- Conduct and catalyze multidisciplinary strategic research and policy analysis of an international public goods nature on all aspects of aquatic resource management, conservation and use;
- Undertake research, training and information activities in partnership with others in national organizations in the developing and developed world;
- Develop global knowledge bases for living aquatic resources;
- Undertake global reviews and assessments of the status of aquatic resource and those who depend on them;
- Publish and disseminate widely research findings;
- Hold conferences, meetings and workshops to discuss current and future issues related to aquatic resources and to formulate advice for users and decisionmakers;
- Participate fully as a Center in the Consultative Group on International Agricultural Research (CGIAR) and in appropriate international intergovernmental activities.

The Guiding Principles for our Work Program are:

- Sustainability;
- Equity;
- Gender role in development;
- Participation;
- Systems approach; and
- Anticipatory research.

Our Values:

In our work, we are committed to:

- Excellence in achievement;
- Relevance to our beneficiaries' needs;
- Partnerships;
- Centerwide teamwork;
- Communication;
- Efficiency and flexibility in program delivery;
- Continual growth in our knowledge and understanding.

