

[Open Peer Review on Qeios](#)

The Spatial Politics of the Tonle Sap: A Multi-Scale Analysis of Conservation and Development Challenges

Mak Sithirith

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Key resources of Tonle Sap Lake are fish, biodiversity and natural resources. Its governance has been casted as a central pillar in the development of Cambodia. Although there are considerable literatures and documents, supporting the protection and conservation of the lake, there are also policies and development frameworks promoting hydropower investments and infrastructure developments, affecting the TSL. Accordingly, this study questions how the water, fishery, and biodiversity and infrastructure developments are embedded in the conservation and developments of TSL. In answering this question, the study conducts the literature reviews and case studies in the Tonle Sap Biosphere Reserves, employing the political ecology approach. It concludes that TSL is food producing engines of Cambodia, the heartbeats of the Mekong River Basin, and a global biodiversity hotspot. The protection and conservation of the lake bring benefits to all at national, regional and global levels, but the complex interplay of actors, arranged at various scales and levels, has collectively created the challenges facing the lake, giving rise to confrontation between the conservation and hydropower, questions about the long-term sustainability of its resources and the livelihoods that depend upon it.

Mak Sithirith

WorldFish Center, Cambodia;

E-mail: maksithirith@gmail.com / m.sithirith@cgiar.org

Keywords: biodiversity, biosphere reserves, conservation, fishery, Mekong river, zoning,

1. Introduction

Key resources of Tonle Sap Lake are fish, biodiversity and natural resources. Its governance has been casted as a central pillar in the development of Cambodia. Although there are considerable literatures and documents, supporting the protection and conservation of the lake, there are also policies and development frameworks promoting hydropower

investments and infrastructure developments, affecting the TSL. Accordingly, this study questions how the water, fishery, and biodiversity and infrastructure developments are embedded in the conservation and developments of TSL. In answering this question, the study conducts the literature reviews and case studies in the Tonle Sap Biosphere Reserves, employing the political ecology approach.

TSL is the largest freshwater lake in Southeast Asia and the MRB, covering 2500 km² in the dry season, but it increases to 10,000-13,000 million km² in the wet season. The lake is connected to the Mekong River via the Tonle Sap River (Kummu and Sarkkula, 2008; Kummu et al., 2014) and has an exceptional water regime. In the wet season, estimated 83 km³ water flows into TSL, of which the Mekong River contributes 54%, the lake's tributaries 34%, and the precipitation 13%. The lake absorbs volumes of floodwaters and reduces flooding along the Mekong River (Kummu et al., 2014). The water flows from the Mekong River, bringing sediments to TSL, providing rich nutrients for fish and biodiversity, and fertilizing the agricultural lands along the Mekong River and Delta. The dry season, from November to March, when the water level in the Mekong River is lower than that of the Tonle Sap Lake, the lake outflows its water to the Mekong mainstream through the Tonle Sap River, estimated at 82km³, of which 84% via Tonle Sap River and 13% via evaporation (Kummu et al., 2014).

The Mekong River, tributaries, hydrological processes, and the flows of TSL produce the Tonle Sap's flood pulse (timing, modality, speed, height, duration), which is of tremendous importance to fisheries productivity and biodiversity in the Lower Mekong Basin and Tonle Sap system (Poulsen et al., 2002). It is also a fact that the 'flood pulse' transforms both the physical and the human landscapes of the Lower MRB and TSL, submerging vast areas during the wet season, exposing those areas in the dry season, creating inundated forests with unique species adapted to the rhythms and cycles of the pulsing ecosystem, and providing a natural habitat for many species of flora and fauna. TSL is highly productive, and it is rich in fisheries. The lake is home to global species that need protection under international conventions (Davidson, 2006; IUCN, 2016).

The 1995 MRC (write out abbreviation at first mention) Agreement protects the TSL via three important protocols; first, the Mekong can be dammed and diverted; second, the average flows can be maintained; and third, maintaining average flows in the Mekong mainstream during the rainy season is a healthy optimum, which are: a) Of not less than the acceptable minimum monthly natural flow during each month of the dry season; b) To enable the acceptable natural reverse flow of the Tonle Sap to take place during the wet season (MRC,1995). Despite the assurance to protect TSL, it is very vague on protecting the dry season flow and the flow levels in general. Thus one can ask how can we ensure the 'acceptable' dry and wet season flow under the scenarios of hydropower dam development? Consequently, this paper examines the governance of TSL from a political ecology perspective (Sithirith, 2021).

2. Conceptual Framework of Political Ecology

This paper examines the conservation of TSL on the one hand, and the developments of infrastructures and its impacts on the other hands, from the political ecology approach. Middleton (2022) reviewed the concept of political ecology on large hydropower dams. He pinpoints that political ecology is an interdisciplinary field that has grown rapidly since the late 1980s. It has evolved from the merging of human ecology / cultural ecology, development geography, and political

economy, to include anthropology, human geography, political science, and environmental science, as well as fields such as political ecology on narratives and political ecology on scales. He finally highlights key concepts of political ecology on the large hydropower dams in the Mekong, focusing on hydro political ecology, rational hydrological approaches, the ontological politics of water, political economy and governance of large dams, knowledge production, and power relations, livelihoods, the commons, and water justice.

As this study looked into the Mekong River and Tonle Sap Lake; therefore, the researcher draws on the concepts of political ecology, particularly the political ecology on scales and narratives are drawn to analyze the political ecology of the Tonle Sap. In the Mekong, political ecology on Narratives are often employed by politicians, industry, or NGOs to frame problems and orient actions. In the Mekong Basin hydropower debate, narratives have been employed by various actors as a disguise for their agendas and to legitimise their activities (Watts and Peet 2004). Geheb and Sahardiman (2019) discuss the political ecology of Hydropower development in the Mekong River Basin, highlighting that hydropower development is occurring at a rapid, though controversial pace, pitting a variety of stakeholder groups against each other at both international scale and international scale, and affecting state relations across scales. It is the processes of socio-political construction of nature, viewing water as a medium that conveys power, and thus sources of both collaboration and conflict. While the Mekong hydropower narratives do, indeed, attempt to conflate the massive regulation of hydrological systems with large-scale social and economic ambitions, they are also intended to obscure a widespread and systemic effort to control and alienate the region's waters via engineering at multiple scales.

Political ecology is concerned with how geographic scale is constructed and how actors use it to legitimize or delegitimize environmental change (Zimmerman and Bassett, 2003). Environmental and social changes in one locations happens as the result of action, the processes and mechanisms takes place at local, national and international levels. Political ecology can help to illuminate how actors at one scale can construct and disseminate a version of scale that advances their agendas. Political ecology can help to highlight how powerful actors define the scale of environmental change and use these definitions to legitimize their actions (Marston and Smith, 2001; Rangan and Kull, 2009). In the Mekong Basin hydropower debate, hydropower developers and states often claim that the scalar benefits of hydropower, which are often measured at meso or national scales, are larger than the costs, which are often measured at local scales. In TSL, geographical space has been territorialized and reterritorialized into global, regional, and national levels, scale and spaces (Sithirith, 2011 & 2022). As Jan Penrose (2002: 280) put it: "Through territoriality, specific places (including territories) are constructed and it is this process that allows people to harness the material and emotional potential of space." In Penrose's terminology, territoriality helps harness 'the latent powers of space'. The production of space relates to its valorisation, commercialization, and the vital role that space plays in the processes of capital accumulation. Thus, space becomes at once a product and part of a process, serving various productive roles that help to generate revenue and profits for different agents (Lefebvre, 1991). Space holds two sources of latent power. First, a latent material power, and second, a latent emotional power (Penrose, 2002). Space comprises a substance fundamental to human life. Through its constitution of land, water, and atmosphere, space encompasses the basic prerequisites of human survival: the food that we eat, the water that we drink, the air that we breathe, and the resources for protecting ourselves. Second, when the substantive qualities of space are filtered through human experiences of time and process, strong attachments to space

have the capacity to invoke or release an emotional response. This is the latent emotional power of space (Penrose, 2002).

The political ecology study analyzes structures, prevailing institutions, actors, interests, and their powers that influence the management of the exploitation of natural resources (Andreas, Fernie, and Dainty, 2022). *Structures* are slow-moving entrenched patterns that are hard to impact in the short run, typically historical circumstances, cultural traits, and natural phenomena. They may, or are likely to, be restrictive for policy change. *Institutions* are patterns in the society described as "...humanly devised constraints that structure political, economic, and social interaction." (North, 1991:97). Institutions are rules and norms, both formal and informal, that provide structure for behavior and relationships in society. *Actors/agents*, finally, are those taking initiatives and pursuing interests, acting individually, in groups, or cooperating spontaneously through shared norms and or strategically to achieve desired future outcomes. All actors have varieties of interests, and different interests manifest themselves within and between different categories of actors. While actors, interests, and powers are open-ended categories with no definitive single content, a simplified inventory – identifying the most significant actors and their respective powers and interests – can still serve as a basic framework for a political economy analysis.

Central to numerous discussions about political ecology is the concept of power, politics, and policies (Jones, Jones, and Woods, 2004). Power is the commodity that sustains politics and policy, and 'politics' is the whole set of processes that are involved in achieving, exercising, and resisting power' while policy relates to the 'intended outcome'—the things that power allows one to achieve and that politics is about being in a position to do (Jones, Jones and Woods, 2004: 3). Resource politics is a politics of 'access' (Ribot and Peluso, 2003) and about issues of state centralization and decentralization and within these essential political strategies and policies relating to allocating and distributing rights to ordinary people, there are both territorial and non-territorial aspects (Wittayapak and Vandergeest, 2010).

3. Materials and Methods

The conceptual framework above is utilized to analyse the political ecology of the Tonle Sap. To do this, the study selected three sites in three Core Areas in the Tonle Sap Biosphere Reserve—Prek Toal, Boeung Tonle Chmar and Stung Sen, to undertake the data collection. Prek Toal is the first TSBR Core Area, located on the northwest side of TSL in Battambang Province, which is rich in fishery, birds and biodiversity. Boeung Tonle Chmar is a second TSBR Core Areas located in Kampong Thom Province. The third Core Area is the Stung Sen, located in the southern end of TSL, in Kampong Svay District, Kampong Thom Province (see Figure 1).




Figure 1. The studied areas in Tonle Sap Lake (Adopted from MRC, 2017).

The investigation gathered data from structured interviews with stakeholders at national and community levels. The research was conducted in three phases. First, between 2007 and 2011 information was collected on the geographical classification of TSL—the commercial fishing lot system, the zoning of TSL for environmental conservation, resource governance, and the politics of resources (Sithirith, 2011).

Second, between 2013 and 2013, a study was conducted that, focused on the aquatic and agricultural system in Tonle Sap, the linking between fisheries, agriculture, livelihoods, governance, and policy frameworks (Johnston et al., 2013). Additional data and information that were collected, focused on fisheries, agriculture, water, and livelihood activities of fishing communities in TSL. The participants involved in the study included: (i) villagers selected from thirteen villages including water-based, water-land-based and land-based communities around TSL; (ii) officials from government agencies at the national, provincial and local levels concerning the fisheries and agriculture in TSL; and (iii) NGOs and researchers worked in TSL. The study asked the participants around fish catch, agricultural activities, alternative livelihoods, water management, agricultural market and other related livelihoods activities.

Third, between August 2022 and March 2023, in collaboration with the Mekong River Commission (MRC), a gap analysis, using the IUCN Green List Framework, was undertaken to assess whether the current management of the Tonle Sap Multiple Use Area (TS-MUA) meets the IUCN Criteria or not; focused on good governance of biodiversity conservation in three Core Zones of the TSBR, the sound design and planning, effective management and successful conservation outcomes. During the study period; first, a stakeholder meeting was held with representatives from MAFF, FiA, MOWRAM, TSA, and IUCN to discuss the management of three Core Zones of TSL prior to the fieldwork. Second, the fieldwork conducted during November 2022 involved consultations with site managers, park rangers, NGOs, and meeting local communities. Key informant interviews (KIIs) with site managers, park rangers, NGOs and village chiefs were conducted in three Ramsar sites—Prek Toal, Boeung Tonle Chhmar, and Stung Sen, focusing on the governance of the sites, the design and planning, the management, the outcomes of the conservation, livelihoods, and the participation of local communities. After the KIIs, three focus group discussions (FGDs) were organized, one FGD in each Ramsar Site, participated by 12-15 rangers and officers in charge of the Ramsar site and villagers from the surrounding villages. Some questions related to indicators of the IUCN Green List were asked during the FGDs. In addition, participants in the KIIs and FGDs were volunteered and informed consent. Information collected was highly kept confidential and anonymous. The participants were protected any harmful effects.

Data and information collected were entered into the Excel sheet of the IUCN Green List indicators including: (1) the good governance, (2) the sound design and planning, (3) effective management and (4) the conservation outcomes. The Excel sheet was analysed using the tables, percentage and graphs to explain the governance, livelihoods and conservation in TSL. In addition, the political ecology framework is employed to analyze the territorialization of fisheries resources in TSL, to examine the spatial politics in the lake, look into the overlapping spaces, actors' involvements in different spatial dimensions, and power relations of actors over spatial differences and to explore the impacts of hydropower developments on TSL, and finally, the changing environment.

4. Results

4.1. Tonle Sap Lake and Socio-Cultural Ecological System

The Mekong River, tributaries, hydrological processes, and the flow of TSL produce numerous complex non-human and human spatial dynamics, which are never totally within or under 'human' control. TSL is characterized by a 'flood pulsed ecosystem'. The annual 'flood pulse' (timing, modality, speed, height, duration) is essential to fisheries productivity and fish migrations in the Lower Mekong Basin and Tonle Sap system (Rainboth, 1996; Poulsen et al., 2002). The 'flood pulse' submerges vast areas of TSL and its floodplains during the wet season, exposing those areas in the dry season, creating inundated forests with unique species adapted to the rhythms and cycles of the pulsing ecosystem, and providing a natural habitat for many species of flora and fauna. (Lambert 2006 & 2013)

TSL naturally comprises three ecological functioning zones—the terrestrial, the floodplain, and the aquatic. The terrestrial zone covers large areas—rice fields and flooded forest areas. The floodplain is an area where it is flooded for six months and dry for another six months. However, the aquatic zone is an area where it is covered year-round by water (Bailleux,

2003).

The natural ecological system of TSL influences how people live in the lake and use the lake resources to sustain their livelihoods. The human in TSL has organized their settlements in three different categories, located in three different ecological zones: land-based, water-based, and water–land-based communities. The land-based community (LBC) is a community that is situated on land for the entire year, whereby community member is engaged in farming and fishing, depending on the distance between the lake and the community. The water-based community (WBC) refers to a floating community, whereby the community is floated year-round on water, and fishing is a primary occupation of community members. The third is a water–land-based community (WLC), whereby the community physically is situated for six months on the water and six months on land. These communities are located in the ecological zone, affected by seasonal water levels (Sithirith, 2011).

People living in water-based and water-land-based communities are called '*neak tonle*' (river people). They are engaged in fishing as their main livelihood activity. The '*neak tonle*' lives adapting to the lake and river system; particularly to the hydrological regime of the lake between the dry and wet seasons including the rising and falling water; the vertical and horizontal motilities; seasonality; resources; and spaces.

Neak tonle catches fish for consumption but needs paddy rice to supplement their diets. Rice and fish are the backbones of household economies. The surplus of fish catches is thus bartered paddy rice with '*neak leu*' or highlander from the land-based community. The bartering of rice and fish between *neak leu* and *neak tonle* is similar to what James Scott (1976) describes as "equal exchange" in the moral economy of peasant society. More specifically, it means that "a gift or service received creates, for the recipients, a reciprocal obligation to return a gift or service of at least comparable value at some future dates. The notion of equal exchange was a general moral principle of peasant society" (Scott, 1976: 167). 'Reciprocity' is viewed as a moral principle underlying social action in the *neak leu* and *neak tonle* society.

The social relations of *neak tonle* and *neak leu* are deeply rooted in the rice-fish economy, allowing them to exchange different products such as fresh fish, dried and fermented fish for rice, and other agricultural products. This relationship enables the *neak tonle* and *neak leu* to complement one another and specialize in fishing and farming respectively. *Neak tonle* and *neak leu* live together as one system, where one respects the function and roles of the others. These have created a dependency and reciprocal system between *neak leu* and *neak tonle* to share their resources for their livelihoods (Sithirith, 2016).

Villager living in a land-based community around TSL is named *neak leu* (highlander), and they organize their farming fields in the Tonle Sap floodplain into *Sreleu* (rainfed lowland ricefield), *Srekandal* (medium deep-water rice-field), and *Srekrom* (Deep-water rice/floating rice field). *Sreleu* is located in the upper area of the Tonle Sap floodplain between 8 and 10m above sea level (asl). *Srekandal* is located deeper inside the floodplain, 6 and 8m asl, lower than *Sreleu* but higher than *Srekrom*. *Srekrom* (Deep-water rice/floating rice field) is located deeper inside the Tonle Sap floodplain area between 4 and 6m asl (Sithirith, 2016).

The rice cultivation in *Sreleu* has relied on rainfall, not the lake's hydrology. In *Srekandal*, farmers cultivate floating rice in

the past but have changed to planting receding rice at present, after the receding water in the lake. *Srekrom* (*Deepwater rice/floating rice field*) is located deeper inside the Tonle Sap floodplain. The rising water in TSL influences the rice productivity of *Srekrom*. Thus, farmers cultivate rice varieties that grow following the rising and falling water levels, known as a '*Srove Leung Tuk*' translated as 'rising water rice' in English. In the dry season, farmers cultivate in this area a 'dry season rice.' However, farming is traditional, small-scale, and subsistent, largely dependent on rainfalls and sometimes on lake water (Sithirith, 2016).

Farming in *Sreleu*, *Srekandal*, and *Srekrom* has been affected by too much and too little water. During the wet season, there is too much water in the TSL floodplain, and the too-much water sometimes causes flooding and damages rice farming and people's properties. During the dry season, water from TSL recedes this floodplain and flows into the Mekong River with no uses, as there are no storage systems, reservoirs, or basins that could extract water out of too-much water situation and store it for use in the season of too-little water.

4.2. The 'Global Space' of Biodiversity

Space in TSL is constructed and re-constructed, and actors are involved in the process of spatial construction. Different actors, community, state, and non-state actors; from different levels—local, national, regional, and global—construct various spaces in the same geographical areas, and much of the spatial exercise involves power (Sithirith, 2015; Keskien, 2006). First, space comprises the resources fundamental to human life on this planet. Space encompasses land, water, and the atmosphere that provide the resources for human survival, including the food we eat, the water we drink, the air we breathe, and the resources for protecting ourselves. Second, when the substantive qualities of space are filtered through human experiences of time and process, strong attachments to space can invoke or release an emotional response (Penrose, 2002; Lebel, Garden & Imamura, 2005).

4.2.1. Biodiversity Hotspots

TSL is highly productive. The annual fish catch from the lake is estimated at between 180,00–250,000 tons, while the dai fishery on Tonle Sap River annually harvests about 12,000 tons of fish migrating from the lake to the Mekong River early in the dry season (Campbell et al., 2006; van Zalinge et al., 2004). Cambodia ranks 5th globally in terms of inland fish production after China, India, Bangladesh, and Myanmar, estimated at around 500,000 tons annually, of which about 50% come from TSL. Fish and aquatic resources have been a source of food and protein for the Cambodian population. Also, fish production has been central to the Cambodian economy (Degen et al., 2000; Degen and Thouk, 2000; Bruce and Yim, 2004).

Overall it is home to many species, including more than 200-370 species of plant, 197 species of phytoplankton, 46 species of zooplankton, 210-225 species of bird, 20-46 species of mammal, 30-42 species of reptile, 300 species of invertebrates, and 05 species of amphibians. At least 44 species are globally threatened or endangered (MRC, 2010). Fish is the largest group of vertebrates, which are abundant in the Tonle Sap ecosystem (Lamberts, 2006). Although the number of fish species in the Tonle Sap is unknown, it is estimated. Approximately 500 species of fish have habituated in the Mekong River system in Cambodia, and at least 143 species have been recorded in the Tonle Sap Lake and

surrounding lowlands. Twenty-two of these species are known to be of international conservation concern (Davidson, 2006). For the above reasons, TSL was designated as Tonle Sap Biosphere Reserve (TSBR) under UNESCO auspices in 1997 for biodiversity conservation (RGC, 2001).

Given the biodiversity and some rare species in TSL, UNESCO has worked with RGC to designate the lake as a Biosphere Reserve. In 1997, TSL was declared a Biosphere Reserve. The Tonle Sap Biosphere Reserve was established by Cambodia and supported by the Royal Decree issued in April 2001. The 'Biosphere Reserve' classifies TSL into three zones—the Transitional, the Buffer, and the Core Zones. The transition zone covers an area of approximately 899,600 hectares, surrounded by 541,482 hectares of buffer zone of permanent and seasonal wetlands, including the lake water bodies. The Core Zone is classified into three Core Areas—Prek Toal, Boeung Tonle Chmar, and Stung Sen—covering an area of 58,635ha. Furthermore, since 1996, these three Core Areas have been designated as Ramsar sites to promote the value of biodiversity conservation in the region internationally. These include (1) Prek Toal covering 21,342ha, (2) Beung Tonle Chmar (BTC) covering 28,000ha, and (3) Stung Sen covering 9293ha (Table 1).

4.2.2. *Human Dimension*

The three Core Areas are also home to 5,426 households living in 19 water-based villages (floating communities) and dependent on fishing and lake resources for their livelihoods. Five villages are situated in Prek Toal with a total population of 12,424 people, equivalent to 2,704 families, most of whom live in water-based communities. Eight water-based villages are in the BTC Ramsar Site—five in Kampong Thom Province and three in Siem Reap Province—which is home to 1,222 households with a total population of 6,044 people living in floating houses, moving from one place to another place during the dry and rainy season and do fishing somewhere near the BTC for their living. There are no farmlands, only water bodies and wetlands, where fishing is the main livelihood activity. On average, poorer families spend 2-3 hours for fishing per day and can catch 5-10 kg of fish or more if they are lucky. Villagers keep some fish for their consumption and sell the rest of their catch immediately (Meynell et al., 2019). About 1,500 households live in Stung Sen Ramsar Site (SSRS), with a total population of about 7,500 people. The SSRS was under the fishing lot no.2 located in Kampong Thom Province, but in 2012, RGC canceled the fishing and returned it to the fishery conservation area (FCA), covering 921 ha (Herranz, Muñoz, and Vong, 2022).

4.2.3. *Biosphere Reserves and Ramsar Sites*

Prek Toal, BTC, and Stung Sen are the Core Areas of Tonle Sap Biosphere Reserves in 1997. Prek Toal is a Core Zone of Biosphere Reserves issued by a Sub-decree in 2001, covering 21,342ha. In October 2015, RGC and IUCN granted the Prek Toal status as the Ramsar site by Sub-Decree No. 139, issued on October 2, 2015. It is the largest Ramsar site or the core area in the TSBR, which is considered a vital area for water bird breeding and foraging habitats and supports many endangered species of water birds. The site is an area for water bird breeding and foraging habitats, among other areas in Cambodia. Prek Toal supports endangered species of water birds, such as the Spot-billed Pelican, Painted Stork, Lesser Adjutant, Greater Adjutant, three species of cormorants, Asian Openbill, Black-headed Ibis, Back-necked Stork, Oriental Darter, Grey-headed Fish-eagle, Glossy Ibis, and Masked Finfoot. Apart from the above, many water bird species, such as Great Egrets, Herons, etc., still nest, breed, and live here almost year-round. Mammals in the Prek Toal

Ramsar Site are diverse, and many species that are listed as globally threatened, and those species' populations are of international importance (Davidson, 2006).

Beung Tonle Chma (BTC) is the second largest Core Area in TSL after the Prek Toal, covering 14,560ha. However, in 1996, it was designated as the largest Ramsar site in TSL, covering 28,000ha, including 3,800ha of a permanent water body known as a Boeung Tonle Chhmar, surrounded by a complex creek system, flood plains, and flooded forests. The maximum elevation of the site is 10 m above sea level (Meynell et al., 2019). BTC Ramsar Site supports a large assemblage of plant, fish, reptile, mammal, and water bird species, many of which are vulnerable or endangered. It regularly is home to more than 20,000 individuals of large water birds, being one of the feeding grounds for the breeding colonies of birds at Prek Toal, such as Asian open-bill, Oriental darter, spot-billed pelican, Indian cormorant, lesser adjutant, and greater adjutant. There are also 296 fish species in the Tonle Sap area (43% grey fish, 40% white fish, and 17% black fish species), of which 17 are threatened species. When inundated, the Boeung Tonle Chhmar site provides a rich habitat for fish feeding and breeding.

Stung Sen is the third Core Area in Tonle Sap Lake, covering an area of 6,365ha. It has been designated as a Ramsar Site in November 2018, covering an area of 9,293ha. It is home to six villages in Phat Sanday Commune in Kampong Thom Province.

Site	Biosphere Reserves Core Area (ha)	Ramsar Area (ha)
Prek Toal	21,342	21,342
Beung Tonle Chmar	14,560	28,000
Stung Sen	6,365	9,293
Total	58,635	42,267

Table 1. The Biosphere Reserves, Ramsar Sites, and Fishery Conservation Area

Source: MoE, 2022

4.3. National Space—Fishing Lots and Fishery Conservation Areas

TSL is a national space. The state has an 'absolute power' to manage that space. Indeed, it is concerned with how TSL could contribute to the national economy and local livelihoods. The French Protectorate Administration territorialized TSL into commercial fishing lots in 1908. The fishing lot system was maintained by various government regimes between 1954 and 2012 to generate revenues for the state following the departure of the French (Degen and Thouk, 2000; Chevyv and Le Poulain, 1940; Degen et al., 2000). By the 2000s, 56 fishing lots, covering about 507,731ha, were operated in TSL (Bruce and Yim, 2004; Sithirith and Vikrom, 2008). While it brought benefits and revenues to the state, it also created conflicts and tensions. The fishing conflicts between fishing communities and fishing lot operators intensified following the 2000s, leading to its abolishment in the 2012s. The fishing lot system prevailed in Cambodia for about 150 years (Sithirith, 2011).

The abolishment of the commercial fishing lot system has released large fishing areas under the fishing lot management system into “open access”, leading to over-exploitation of fisheries and conflicts between fishermen to compete for access and maximization of fisheries resources. Faced with such a challenging situation, the RGC has issued a decree to re-territorialize the open access area into a fishery conservation area and community fishery to reduce anarchic fishing activity in the TSL. About 222 community fisheries around the lake have been established in six provinces around the Tonle Sap, with 94,033 households as members. It covers about 538,739ha around the TSL. Most of CFis in TSL have registered with MAFF. At the same time, each community fishery (CFi) has demarcated a boundary and drawn the map to differentiate one CFi from the others. Members of the CFis are allowed to fish for subsistence only, using small-scale fishing gear.

4.3.1. *Fishery Conservation Area (FCA)*

After abolishing the fishing lot system in 2012, RGC has transformed 93,246ha, or 35 percent of abolished fishing lot areas into fishery conservation areas. The fishery conservation areas were further territorialized into 23 fishery conservation areas similar to fish sanctuaries. FiA is the sole agent responsible for the fish sanctuary and fishery conservation area management. Large canceled fishing lot areas in Battambang, Pursat, and Kampong Thom Provinces were converted into fishery conservation areas, accounting for 51 percent, 44 percent, and 26 percent respectively. These include the former fishing lot no.2 (50,134ha) in Prek Total in Battambang, the former fishing lot no.2 (7476ha) in Pursat, and the former fishing lot no.6 in BTC in Kampong Thom provinces.

Before the 2012s, three out of 37 fishing lots in TSL overlapped the Core Areas. FiA considered these fishing lots as the richest fishery areas in TSL. The Prek Toal was under the fishing lot no.2 in TSL in Battambang Province, which covers 50,134ha. The Boeung Tonle Chmar was under the fishing lot no.6 in TSL in Kampong Thom Province, which covers 8.325ha. Also, the Stung Sen was under the fishing lot no.2 in TSL in Kampong Thom Province, which covers 921ha. These areas were rich in the fishery and the most productive fishing lot areas in TSL.

FiA has a mandate to manage the fishery conservation areas (FCA). The Core Areas and the Ramsar Sites in TSBR overlaps with the FCAs in each site. In Prek Toal, the FCA covers 50,134ha, the BTC’s FCA covers 8,325ha, and Stung Seng’s FCA covers 921ha. The FCA areas were former fishing lot areas, rich in fisheries, abolished in 2012.

4.3.2. *Community Protected Areas (CPAs) and Community Fisheries (CFis)*

In the TSL, many CFis and CPAs were established. In 2001, RGC removed part of the fishing lot areas for community use, and in 2012, the entire fishing lot system was abolished and turned large fishing lot areas into open-access fishing areas and community fisheries (CFis). Five CFis were established in the Prek Toal area, covering 19,713ha.

Also, there is one Community Protected Area (CPA) in BTC Ramsar Site that was established in 2006-2007 through the UNDP Tonle Sap Conservation Project (TSCP). The CPA covers 65ha, including 8ha of the Fishery Conservation Area (FCA). After a short period, the CPA was not functioning, and then it was revived by MoE with technical and financial support from the EU-NSA IUCN project (2013-2016) (IUCN, 2016). It is managed by the MoE/PDoE of Kampong Thom

Province. In addition, in 2012, 37 fishing lots in TSL, including Fishing lot no.6 in BTC, were permanently abolished and handed back to fishermen's families for their daily livelihood. Following this decision, two Community Fisheries (CFis) (including small FCAs with strict protection) were established in Peam Bang Commune, inside BTC Ramsar Site (BTC-RS), covering 2,883ha. These CFis are located in the floating villages of Pov Veuy, covering 2676 ha and Peam Bang 208 ha. The CFis fall under the Ministry of Agriculture, Forests and Fisheries (MAFF) and are managed by elected CFI Committees under the Fisheries Administration (FiA).

Within Stung Sen Ramsar Site (SSRS), a Community Protected Area (CPA) was formed in Toul Neang Sav Plov village (Phat Sanday commune) in 2009, covering 1,495ha. NatureLife Cambodia / BirdLife International (NL/BL) works with Toul Neang Sav Plov CPA to strengthen CPA management and effectiveness. NL/BL also established one additional CPA in Phat Sanday commune and built their capacity and expertise for site management to support livelihoods and habitat conservation in SSRS. In addition, fisheries in SSRS are managed by the Phat Sanday Community Fishery (CFi), established by FiA in 2007, covering 8,810ha, and registered with MAFF in 2008. Five villages in Phat Sanday Commune are members of the Phat Sanday CFI. The Phat Sanday CFI is administered by the Fisheries Administration (FiA) under the jurisdiction of MAFF. SSRS also overlaps with Lower Stung Sen Key Biodiversity Area (KBA) and Important Bird Area (IBA). Fish Conservation Areas (FCAs) or fish sanctuaries have been established within the IBA and nearby open waters of TSL.

4.4. The Regional Space and Hydropower

The Tonle Sap River acts as a valve or artery connecting the Mekong River to TSL, and thus, the Tonle Sap's biophysical characteristics cannot be considered without reference to the Mekong hydrological regime. TSL owes its uniqueness to the natural phenomenon of reverse water flow, with approximately half of an annual pulse absorbed by the Lake area from the Mekong River during the wet season (May to October) and released back during the dry season¹ (Nikula, 2005:13; Kammu *et al.*, 2008). Hence, the Tonle Sap is part of the Mekong River Basin, and the 1995 MRC Agreement recognizes TSL as a valuable area in the Lower Mekong River Basin. Further, four Lower Mekong Countries signed the 1995 MRC Agreement to protect it (MRC, 1995). Article 5 of the Agreement guarantees the equitable and reasonable utilization of the Mekong water. First, the Mekong can be dammed and diverted; second, the average flows can be maintained; and third, maintaining average flows in the Mekong during the rainy season is a healthy optimum. The question is how to protect TSL using this article given its vague meaning and confused terms (Ojendal, 2000; Sithirith, 2007). Article 6 also raises two concrete points in maintaining flows in the mainstream: a) Of not less than the acceptable minimum monthly natural flow during each month of the dry season; b) To enable the acceptable natural reverse flow of the Tonle Sap to take place during the wet season (MRC, 1995). Despite the assurance to protect TSL, it is very vague and unclear on protecting the dry season flow and the flow levels in general, but what is the dry season? How much is the acceptable low flow? How can we ensure the 'acceptable' dry and wet season flow under the scenarios of hydropower dam development? Hitherto, the real meaning of these terms is left out with no clarification or explanation (Sneddon, 2003).

However, hydropower is still possible under the 1995 MRC Agreement. These have put TSL under threat. Since the 1950s, nearly 6,000 large and small dams were built in the lower Mekong River Basin (FACT and EJV, 2001). Between

1965 and 2005, 22 major dams, hydropower, and irrigation schemes were built in four lower Mekong countries: Thailand, Laos, Vietnam, and China—About 40% of these dams were for irrigation and the rest for hydropower. The active storage capacity of these dams was 15,328 million cubic meters (mcm).

After the 1990s, 156 hydropower projects have existed in the Mekong Region—some built, some under construction, and some under planning. These dam projects could potentially produce a total capacity of 52,043 MW with a storage capacity of 126,890 MCM. Some 24 hydropower projects are being planned and built on the mainstream of the Mekong River, 13 dams in the Lancang River in Chinam, and 11 dams in the Lower Mekong River Basin (LMB). Eleven out of 13 Chinese hydropower dams were built between 1990 and 2020 and operated to generate electricity. Of 11 mainstream dams in the LMB, 9 are in Laos, and two are in Cambodia. In Laos, Xayaburi Dam was completed and operational in 2019; Don Sahong was completed in 2020; Pak Beng Dam has completed the PNCA and is under planning; and two other Laos dams are under the PNCA.

Some 132 hydropower projects have been planned and built on the tributaries of the Mekong River in the LMB, of which 25 dams are operational, 13 are under construction, 23 are under license, and 74 dams are planned (MRC, 2017). In the 3S river basin, there are 42 dams—three major hydropower dams were completely built on the Sekong, eight on the Sesan and seven on the Srepok, and 23 are under planning (Constable, 2015; Piman et al., 2016; Piman et al., 2013).

Laos has planned to build 100 hydropower dams, of which 91 are on the tributaries and nine on the mainstems of the Mekong River, with a total capacity of more than 20,000MW and the storage capacity of 57,477 MCM. Cambodia has planned 21 dams in the Mekong basin in Cambodia, of which two are on the mainstream (Sambo and Stung Treng), and the rests are in the 3S region and around TSL. Vietnam, a downstream country in the Mekong River, has planned for 20 dams, of which 15 are built in the 3S rivers², while Thailand continues to finance the hydropower projects in Laos to import electricity to Thailand.

Dams would release water in the dry season to maintain their hydroelectricity outputs and release the water to protect dams during floods. In addition, dams possess flow regulation with a total regulation storage capacity. The flow regulation entails a few seasonal floods downstream during the usual years. A few seasonal floods, however, are likely to result in a decline in soil fertility over the large areas of rice cultivation in the Lower Mekong Basin. Fewer seasonal floods would also mean less natural capacity to constrain saltwater intrusion from the sea into the Mekong Delta. As a result, aquatic life long adapted to the Mekong ecosystems could be seriously jeopardized by a changing flow regime. Fish migration could be blocked by dam building. If the Mekong's biodiversity dropped, this would likely be accompanied by falling productivity in the wild fisheries.

Hydropower dams alter the flow and the volume in TSL. According to MRC, in 2018, the wet season flow has reduced by 7%, and the dry season flow increased by 7% at all key stations, including the station in Kratie in Cambodia (MRC, 2018; Baran and Ratner, 2007; Halls and Kshatriya, 2009; Kirby and Mainuddin, 2009; Baird, 2009; Baran and Myschowoda, 2009). The flow at Kratie would reduce to 4,000m³/s in the wet season, and the dry season flow would increase to 2,200m³/s (Piman et al., 2013). These will further alter the flow and volume of water in the TSL and affects fisheries productivity. In the 3S Basin, if all planned dams are built and operated, the dry season water flow is increased by 98% at

the 3S outlet (Piman et al., 2016), and the wet season flow is decreased by 22% (Pinan et al., 2013).

The decline in water level significantly started in 2014. During 2014-2015, Cambodia experienced a severe drought, and the Tonle Sap water level was relatively low. In 2015, a forest fire in TSL occurred in Battambang Province due to the long and severe drought and the low water level. The severe drought in Cambodia continued in 2016, and the Royal Government of Cambodia declared a state emergency due to a lack of water for human consumption, and the State responded by distributing water to its populations across the country. In the Mekong Delta, more than 2 million Vietnamese and major Vietnamese rice production areas were impacted by low water levels and severe saline intrusion in 2016, resulting in over USD 670 million in losses. In March 2016, the Chinese government released water from upstream dams to relieve drought in the LMB and southern Vietnam.

The reverse flow from the MR to TSL occurs from mid-May to mid-October. The long-term average (LTA) annual Reverse Flow (RF) volume from the MR to TSL is about 40.38 km³ between 1996 and 2015. However, the trend of the RF volume of TSL declined between 1997 and 2020. In the 2019s, the RF started until the first week of August, three months late. The total RF volume to TSL was estimated at 31.48 Km³, critically lower than the LTA of TSL (40-42 Km³). In the 2020s, the RF of TSL took place intermittently; the 1st RF started on 7 July and ended on 15 July, with a total volume of just 0.21 Km³; the 2nd RF occurred in the last week of July, or early August 2020, resulting in an accumulated RF volume of 12 Km³; the 3rd and 4th RF events occurred in late September and the third week of October 2020. The total volume of the RF in 2020 was only 18.89 Km³ or about 44% of the acceptable annual volume of 43 Km³ (average condition for 1997–2005).

The changes in the RF from the MR to TSL between 2018 and 2020 and the reduction in the inundation around the lake degrade the lake's productivity. The Dai fishery production in TSR declined from 16,975 tons in 2018 to 9,900 tons in 2020. The decline in inundated areas around the lake, the low volumes of floodwaters entering the lake, the degradation of fish habitats due to lower water levels, and the destruction of flood forests between 2018 and 2020 contributed to changes in fish catches in TSL.

The annual fish catch from TSL accounts for 250,000 tons. However, between 2018 and 2020, it dropped from 291,260 tons in 2018 to 144,635 tons in 2020—a 50% reduction. The decline of fish production in TSL is associated with the lowering water level, volume, inundated areas, and flood duration. Of course, it cannot deny other factors that contributed to the decline in fish catches in TSL, such as the destruction of flooded forests, illegal fishing, and the disappearance of some fish species (Figure 2). Figure 2 also demonstrates the decline in fish catch by provinces in TSL. In Siem Reap, Kampong Chhnang, and Kampong Thom provinces, fish catch has contributed significantly to the provincial economies and livelihoods of communities around TSL. However, they recently experienced a drop in fish catch. The reductions in fish catches have transformed the human connection with the lake.




Figure 2. Inland fish production in TSL by years and by provinces in TSL (FiA, 2021)

The decline in inland fish production in TSL has affected fishing households and the fishing population. In the Tonle Sap provinces, fishing households declined by 10% between 2018 and 2019. Also, the fishing population by sex has dropped by 10% for male and 9% for female fishers. In total, the fishing population has declined by 10%. Banteay Meanchey has a high percentage of declining fishing households (25%) and fishing population (28%), followed by Kampong Thom and Kampong Chhnang Provinces (**Table 2**).

Province	No. of fishing households		Changes (%)	Male fisher		Changes (%)	Female fisher		Changes (%)
	2018	2019		2018	2019		2018	2019	
Kompong Chhnang	51,760	41,516	-20	62,112	49,932	-20	93,168	74,898	-20
Pursat	35,338	35,338	0	67,700	67,700	0	66,357	66,357	0
Battambang	33,040	29,731	-10	46,565	41,908	-10	17,896	16,106	-10
Kompong Thom	45,466	44,466	-2	41,136	41,136	0	4,330	3,330	-23
Siem Reap	72,044	72,044	0	160,706	160,706	0	112,044	112,044	0
Banteay Meanchey	58,030	43,411	-25	120,100	86,500	-28	26,900	19,460	-28
Total	295,678	266,506	-10	498,319	447,882	-10	320,695	292,195	-9

Table 2. The decline in the fishing population in TSL

Source: FiA, 2021

At the Core Areas (Prek Toal, BTC & Stung Sen), various actors have engaged in TSL-MUA's conservation and protection of global species such as birds, Mekong Catfish, and other species. UNESCO has been a leading UN organization in conserving the global endangered and rare species in TSL. UNESCO and the Ministry of Environment are working together to strengthen the sustainable management of the Biosphere Reserve in TSL. On the other hand, IUCN is an active global organization in monitoring the implementation of the Ramsar Convention and promotes the conservation of wetland areas. IUCN has worked with MoE and NGOs such as FACT, CI, and others in Ramsar Sites in TSL to conserve wetland resources in Prek Toal, BTC, and Stung Sen sites.

At the regional level, TSL has been identified by MRC as Mekong regional environmental asset where MRC works with MRC's member countries to protect and conserve it for sustainable Mekong development. MRC has implemented the Mekong-Wetland Project in the Lower Mekong River Basin, financed by KfW. In Cambodia, the Mekong-Wetland Project is implemented by MRC, focusing on Prek Toal. WCS is a conservation NGO subcontracted by MRC/KfW to carry out conservation activities at the community level.

At national levels, MoE/PDoE is an agency responsible for managing the Biosphere Reserves and Ramsar sites in TSL. There is an Office in charge of Biosphere Reserve and Ramsar management with several rangers under the direct supervision of the PDoE. Also, NGOs such as IUCN, WCS, BirdLife, and others have supported each Office in carrying out conservation activities. IUCN, FAO, and other conservation NGOs may provide technical and financial support on landscape protection and biodiversity and improve the livelihood of local conservation communities dependent on the wetland and lake resources. In addition, World Bank has worked with MoE to implement a Cambodia Sustainable Landscape and Ecotourism Project (CSLEP) in the Cardamom Mountain and Tonle Sap Landscape (CMTS). It has implemented the project in BTC and Stung Sen and supported the management for both sites.

Also, FiA has received financial support from EU-CAPFISH Capture Project to manage the fishery conservation area and community fisheries in TSL-MUA. IUCN, WCS, and BirdLife have provided technical and financial support to CFis and CPAs in all three Ramsar sites in Tonle Sap Lake. Communities (local people in the buffer zone) participate in forest plantation, protection, and ecotourism development; cooperate on forest fire precaution and fighting; wise use of wetland resources. In Prek Toal Ramsar Site, WCS and MoE/PDoE have supported the establishment of the Community Based Ecotourism (CBE). So far, the CBE's Prek Toal has received many tourists who have visited Prek Toal annually, around 2 million tourists.

However, there is no private sector (Tourism/Private companies). There is potential to engage the private sector in resource management, such as fresh water, natural landscapes, and the environment to provide ecotourism services or purchasing, processing, and trading NTFPs (aquatic and vegetable products).

5. Discussion and Conclusion

TSL possesses three scales—national, regional and global (Sithirith, 2021). At the national scale, fishery is a key resource for national economy and livelihoods of its people. At the regional scale (the Mekong River Basin), TSL is the heart of the Mekong, where flood pulse plays a critical role in absorbing the floods in the wet season and releasing it in the dry season (Campbell, Say & Beardall, 2009). At the global scale, TSL is a hotspot of global species. Concerted efforts have been organized around these scales to protect and conserve TSL, such as UN Biosphere Reserves, Ramsar Convention, the 1995 MRC Agreement and the national conservation program such as the flooded forest zoning. While these bring supports to TSL, it also influences the governance and conservation of TSL from institutional, technical and financial capacities from different scales and levels—global, regional and national.

At each scale, TSL is a space. The TSL space encompasses land, water, fishery, biodiversity and natural resources. For local communities, these resources form the dependent space for their living (Cox, 1998), a latent material and a latent emotional power (Penrose, 2002), and a representational space (Lefebvre, 1991). For the Government agencies; these resources are sectoral, institution, technical specialization, and power defined by the legal frameworks. They are institutionalized with mandates, roles and responsibilities to manage these resources (Keskinen, & Sithirith, 2009). Different institutions compete for different spaces, as of resources, powers, and influences, and so, spaces in TSL are territorialized, deterritorialized and reterritorialized, as Peluso (2005a: 6) has stressed: “territorialisation produces places in relation to claimants”, and there are numerous territories and places in and around the TSL. In addition, problems exist over overlapping claims, multiple functions in the same zones, boundary disputes between commercial, middle-scale, and family fishers, and there is great ambiguity over the specific territorial and resource access rights afforded to different communities. These have implicated the governance and conservation of TSL (Sithirith, 2022).

Space has been constructed, and reconstructed by different actors at different levels. Fishery as a space has been organized into CFIs, public fishing areas (open access) and conservation areas, after the abolishment of the commercial fishing lot system. The conservation area has been reconstructed into the fishery conservation area (FCA) and fishery sanctuary, under the responsibility of Fishery Administration. As for biodiversity conservation, TSL is managed as a conservation area under the Ministry of Environment, reinforced the Royal Decree in 2001. Furthermore, UNESCO has reconstructed the TSL as a Biosphere Reserve for conserving globally endangered species and IUCN has granted the three Core Areas of TSBR as Ramsar Sites for wetland conservation. In the Core Areas and Ramsar Sites, MOE has established the CPAs to engage local communities in the conservation activities. There are overlapping spaces, producing the implications for the governance and conservation of resources in the lake. One space, zone or territory may be managed by two or three institutions with different policy, legal frameworks and technicality, inducing the complexities of space, zone and territories and causing hardship for local people adaptation. These bring more competition than cooperation between concerned institutions and with local communities.

On the other hand, the states in the Mekong region view the Mekong as a resource of economic growth, and modernization. The intensification of economic activity in the basin precedes plans for large hydropower dams. In Laos, hydropower was envisioned as a means of generating foreign exchange and state revenue through electricity export, first to Thailand and later to Vietnam and China. Many private hydropower proponents were from Europe, Australia, and the US, with an influential financing and policy role for Western bilateral donors and multilateral development banks, namely

the World Bank and the ADB (Bakker, 1999). Later in Laos, private project developers and commercial financiers have primarily been from Thailand, Vietnam, and China, although actors from other countries, including Japan, Malaysia, and Korea, are also present (Middleton et al., 2009). In Cambodia, investment in both electricity generation and transmission comes primarily from the private sector and, in the case of large hydropower dams, has primarily come from Chinese developers and financiers (Siciliano et al., 2016). Although there are impacts of hydropower dams on TSL and tensions of over hydropower dams, riparian states compete, cooperate and share the benefits from hydropower dams. The regional and bilateral cooperation over the Mekong, the regional politics and the national interests have shed more lights on hydropower power developments than the conservations of TSL.

In the geographical scale of the MRB, as indicated by Zimmerman and Bassett (2003), the development elsewhere in the MRB such as hydropower dams would cause hydrological changes and impacts on TSL, biodiversity and livelihoods. Although the 1995 MRC Agreement ensures the maintenances of the reverse flows and dry season flow to TSL, this is only rhetoric and it will never happens in the context of increasing hydropower development and climate change. As Mekong riparian states struggle to graduate to middle income countries and reduce poverty, more hydropower dams are being built to generate electricity to boost its economy. The increased Chinese presence in the Mekong Cooperation and riparian states through Chinese investments has brought more development interventions in the Mekong River Basin, which has led to more development of hydropower dams, causing changing hydrological regimes and impacts downstream, particularly TSL. These have further impacted the water, fisheries, natural resources, and livelihoods of local communities dependent on the lake resources. Despite these and breaching international agreements, Cambodia may not choose the way in which to complain about its neighbors who cause the impacts on TSL. Instead, Cambodia may empower itself in the Mekong cooperation through also building dams and infrastructures. Thus, more development projects including hydropower dams and navigation projects have been built and have made impacts to the conservation and management of the lake.

The only to address the above challenges and impacts are through improving coordination across scales and levels and reinforcing global, regional, and national agreements and commitments for TSL. Also, the conservation of global species has benefited TSL, the Mekong Region, and the Mekong countries. At the same time, the development in the region has to be balanced and avoid the consequences following the commitments assured by the regional agreement. Coordination and cooperation between government agencies are second level of important. Coordinated policy, planning, implementation, and management over spaces, zones and territories bring the significant outcomes for the sustainability of TSL.

Acknowledgments

The author would like to extend his thanks to WorldFish and the Dean of the Faculty of Development Studies, Royal University of Phnom Penh for their support.

Declaration of Conflicting Interests

The author declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this

article.

Funding

The author received no financial support for the research, authorship, and/or publication of this article.

Notes

¹ In the wet season, the surface area of the Lake increases from 250,000-300,000 ha to approximately 1.0-1.6 million ha, with depth increasing correspondingly from 1–2 m amsl to 9–11 m amsl, and storage capacity reaching a maximum of 80 million cubic meters. It absorbs 20 percent of the Mekong River's floodwaters and serves as a flood regulator (MRC, 2004; ADB, 2002). The drop in the water level in the Mekong in the dry season creates the “reverse flow” from the Lake into the Mekong.

² This report is taken from the CGIAR Report. The CGIAR Research Program on Water, Land, and Ecosystem. <https://wle-mekong.cgiar.org/wp-content/uploads/unnamed-11.jpg>. Accessed on 13 July 2018.

References

- Andreas B, Fernie S and Dainty A (2022) “Understanding Policy and Change: Using a Political Economy Analysis Framework.” *Construction Management and Economics* 40 (11–12): 865–83.
- Bailleux, Renaud. 2003. *The Tonle Sap: A Pulse of Life*. Bangkok: Asia Horizons Books.
- Baird I (2009) *The Don Sahong Dam: Potential Impacts on Regional Fish Migrations, Livelihoods and Human Health* POLIS Project on Ecological Governance, University of Victoria, Canada.
- Bakker K (1999). The politics of hydropower: Developing the Mekong. *Political Geography* 18(2), 209-232.
- Baran E and Myschowoda C (2009). Dams and Fisheries in the Mekong Basin. *Aquatic Ecosystem Health and Management*, 12 (3):227-234.
- Baran E and Ratner B (2007) *The Don Sahong dam and Mekong fisheries*. Available at: <http://www.terraper.org/articles/DonSahong%20science%20brief.pdf>.
- Benjaminsen T A and Svarstad H (2018) *Political Ecology. Reference Module in Earth Systems and Environmental Sciences*. doi:10.1016/b978-0-12-409548-9.10608-6.
- Bruce M and Yim C (2004) *Domestic Fish Trade: A Case Study of Fish Marketing from the Great Lake to Phnom Penh* Working Paper 29. Phnom Penh: Cambodia Development Resource Institute (CDRI).
- Campbell, T.; Pin, K.; Ngor, P.B. and Hogan, Z. 2020. Conserving Mekong megafishes: Current status.
- Campbell I C, Say S, & Beardall J (2009) Tonle Sap Lake, the heart of the lower Mekong. In *The Mekong* (eds), pp. 251-272. Academic Press.
- Campbell IC, Poole C, Giesen W, Valbo-Jorgensen J (2006) Species diversity and ecology of Tonle Sap Great Lake, Cambodia. *Aquatic Sciences* 68, 355-373.
- Constable, D. (2015). Atlas of the 3S Basins. Bangkok: IUCN.
- Chevey P and F Le Poulain (1940) *La Pech Dans Les Eaux Douces du Cambodge* Travaux De L'institut

- Océanographique de L'Indochine. Le Memoire, Gouvernement General de L'Indochine. Saigon.
- Davidson P (2006) *The biodiversity of the Tonle Sap Biosphere Reserve 2005 status review*. Prep. by Wildlife Conservation Society (WCS) for the Tonle Sap Conservation Project. Phnom Penh: WCS.
 - Degen P, Van AF, Van Zalinge N, Thouk N and Vuthy L (2000). *Taken for Granted Conflict over Cambodia's Freshwater Fisheries Resources*. Written for the 8th IASCP Conference, Bloomington, Indiana, 31 May to 4 June, 2000. Phnom Penh: MRC, DoF and University of Antwerp, Belgium.
 - Degen P and Thouk N (2000) Historical, Cultural and Legal Perspectives on the Fishing Lot System in Cambodia. In *Common Property in the Mekong: Issues of Sustainability and Subsistence*, eds. M. Ahmed & P. Hirsh, 49-60. Penang: Australian Mekong Resources Center, Sydney University, and World Fish Center.
 - Fisheries Action Coalition Team (FACT) & Environmental Justice Foundation (EJF). 2001. *Feast or Famine? A Solution to Cambodia Fisheries Conflict*. London: UK.
 - Geheb K and Suhardiman D (2019) The political ecology of hydropower in the Mekong River Basin. *Current Opinion in Environmental Sustainability* 37: 8-13.
 - Halls A S and Kshatriya M (2009) *Modelling the cumulative effects of mainstream hydropower dams on migratory fish populations in the lower Mekong basin*. MRC Technical Paper No.25, Vientiane: Mekong River Commission Secretariat.
 - Herranz Muñoz V and Vong V (2022) *Climate Change Vulnerability Assessment Stung Sen Ramsar Site, Cambodia* Bangkok, Thailand: IUCN.
 - International Union for Conservation of Nature (IUCN) (2016) Strengthening Capacity of Fishing Communities in the Tonle Sap to Manage their Natural Resources Sustainably. EU-NSA Project Report (2013-2016). Phnom Penh: IUCN.
 - Johnstone G, Puskur R, Declerck F, Mam K, Sithirith M, Pech B, Seak S, Chan S, Hak S (2013) *Tonle Sap Scoping Report-CGIAR Research Program on Aquatic Agricultural Systems*. Montpellier: Consultative Group for International Agricultural Research.
 - Jones M, Jones R and Wood M (2004) *An Introduction to Political Geography: Space, Place and Politics*. London and New York: Rutledge.
 - Keskinen, M., & Sithirith, M. (2009) *Tonle Sap lake and its management: the diversity of perspectives & institutions. Research paper on improving Mekong water allocation project (PN67)*, <http://www.mpower.net.org/mweb.php>.
 - Kirby M and Mainuddin M (2009) Water and agricultural productivity in the Lower Mekong Basin: trends and future prospects. *Water International*, 34(1), 134-143.
 - Kumm M, Tes S, Yin S, Adamson P, Józsa J, Koponen J & Sarkkula J (2014) Water balance analysis for the Tonle Sap Lake–floodplain system. *Hydrological Processes*, 28(4), 1722-1733.
 - Kumm M, Lu X, Rasphone A, Sarkkula J & Koponen J (2008) Riverbank changes along the Mekong River: Remote sensing detection in the Vientiane–Nong Khai area. *Quaternary International*, 186(1), 100-112.
 - Lamberts D (2013) *The role and significance of the flood pulse in the functioning and management of the Tonle Sap ecosystem, Cambodia* (PhD dissertation). KU Leuven, Belgium.
 - Lamberts D, Koponen J (2008) Flood pulse alterations and productivity of the Tonle Sap ecosystem: a model for impact

- assessment. *AMBIO: A Journal of the Human Environment* 37 (3) :178-184.
- Lamberts D (2006) The Tonle Sap Lake as a productive ecosystem. *International Journal of Water Resources Development* 22(3),481-495.
 - Lebel L, Garden P, & Imamura M (2005) The politics of scale, position, and place in the governance of water resources in the Mekong region. *Ecology and society*, 10(2).
 - Lefebvre H (1991) *The Production of Space*. Translated by Donald Nicholson-Smith. Malden, Massachusetts: Blackwell.
 - Marston SA & Smith N (2001) States, scales and households: limits to scale thinking? A response to Brenner. *Progress in Human Geography*, 25(4), 615-619.
 - Meynell PJ, Kong K, Sorn P and Lou V (2019) *Climate Change Vulnerability Assessment Boueng Chhmar Ramsar Site, Cambodia*. Bangkok, Thailand: IUCN.
 - Mekong River Commission (MRC) (2018) THE COUNCIL STUDY: *The Study on the Sustainable Management and Development of the Mekong River Basin, including Impacts of Mainstream Hydropower Projects*. Vientiane/Lao PDR: The Mekong River Commission.
 - Mekong River Commission (MRC) (2010) *State of the Basin Report 2010 Summary*. Vientiane: Mekong River Commission.
 - Mekong River Commission (MRC) (1995) *Agreement on the Cooperation for the Sustainable Development of the MR Basin*. MRC Secretariat. Available on <https://www.mrcmekong.org/assets/Publications/95-agreement.pdf>.
 - Middleton C (2022) The political ecology of large hydropower dams in the Mekong Basin: A comprehensive review. *Water Alternatives* 15(2): 251-289.
 - Middleton C; Garcia J and Foran T (2009) Old and new hydropower players in the Mekong Region: Agendas and strategies. In Molle, F.; Foran, T. and Käkönen, M. (Eds), *Contested waterscapes in the Mekong Region: Hydropower, livelihoods and governance*, pp. 23-54. London, Sterling, VA: Earthscan.
 - Neumann R (2009) Political ecology: theorizing scale. *Progress in human geography* 33(3), 398-406. DOI: 10.1177/0309132508096353.
 - NGO Forum on Cambodia (NGOF) (2023) *Impacts of a Low Flow of Water from the Mekong River to Tonle Sap Lake on the Livelihoods of Different Communities and Natural Environments*. NGOF: Phnom Penh, Cambodia.
 - North D (1991) Institutions, Ideology, and Economic Performance. *Cato J.* 11: 477.
 - Ojendal J (2000) *Sharing the Good: Mode of Managing Water Resources in the Lower Mekong River Basin*. Department of Peace and Development Research. Sweden: Goteborg University.
 - Peluso NL (2005) Seeing property in land use: local territorialisations in West Kalimantan, Indonesia. *Danish Journal of Geography* 105(1), 1-15.
 - Penrose J (2002) Nations, States, and Homeland: Territory and Territoriality in Nationalist Thought. *Journal of Nations and Nationalism*, 8 (3), 277-297.
 - Piman T, Cochrane T A and Arias M E (2016) Effect of proposed large dams on water flows and hydropower production in the Sekong, Sesan and Srepok rivers of the Mekong Basin. *River research and applications* 32(10), 2095-2108.

- Piman T, Lennaerts T, and Southalack P (2013) Assessment of hydrological changes in the lower Mekong Basin from Basin-Wide development scenarios. *Hydrological Processes* 27(15), 2115-2125.
- Poulsen AF, Ouch P, Sintavong V, Ubolratana S and Nguyen T (2002) *Fish migrations of the Lower Mekong River Basin: implications for development, planning and environmental management*. MRC Technical Paper No. 8. Phnom Penh: Mekong River Commission.
- Rainboth WJ (1996) *FAO species identification field guide for fishery purpose Fish of the Cambodian Mekong*, Rome: FAO, 265.
- Rangan H and Kull C (2009) What makes ecology political: rethinking scale in political ecology. *Progress in Human Geography* 33(1), 28-45.
- Räsänen TA, Someth P, Lauri H, Koponen J, Sarkkula J, Kummu M (2016) Observed river discharge changes due to hydropower operations in the Upper Mekong Basin. *Journal of Hydrology*.
Doi: <http://dx.doi.org/10.1016/j.jhydrol.2016.12>.
- Ribot J and Peluso N (2003) A Theory of Access. *Rural Sociology* 68:153-181.
- Royal Government of Cambodia (RGC) (2007) *Royal Decree on the Establishment of Tonle Sap Authority, Draft, Unofficial Translation*. Available Online At: <http://www.foodsecurity.gov.kh/Otherdocs/Royal-Decree-tsba-Eng.pdf>.
- Royal Government of Cambodia (RGC) (2001) *Royal Decree on the Establishment and Management of the Tonle Sap Biosphere Reserve*, Translation by Neou Bonheur, Available online at: http://www.tsbred.org/docs/law_and_regulation/Royal_Decree_On_creation_and_managemet_of_tsbr_Eng.Pdf.
- Scott J (1976) *The Moral Economy of Peasant: Rebellion and Subsistence in Southeast Asia*. New Haven and London: Yale University Press.
- Siciliano G; Urban F; Tan-Mullins M; Pichdara L and Kim S (2016) The political ecology of Chinese Large dams in Cambodia: Implications, challenges and lessons learnt from the Kamchay Dam. *Water* 8(9), 405.
- Sithirith M (2022) Analysing transboundary water governance: A global-regional-national water governance framework of Tonle Sap Lake. *Lakes & Reservoirs: Research & Management*, 27(1), e12396.
- Sithirith M (2021) Downstream state and water security in the Mekong region: a case of Cambodia between too much and too little water. *Water*, 13(6), 802.
- Sithirith M (2016) Political economy of fishing villages: A case study in the Tonle Sap Lake, Cambodia. *Journal of Environmental Science and Engineering B* 5(6):299-313.
- Sithirith M (2011) Political geographies of the Tonle Sap: Power, space and resources (Doctoral Dissertation, National University of Singapore, Singapore).
- Sithirith M and Mathur V (2008) Entitlements and the Community Fishery in The Tonle Sap: Is the Fishing Lot System Still an Option for Inland Fisheries Management? In *Sustaining Tonle Sap: An Assessment of Development Challenges Facing The Great Lake*, eds. Chadwick, M., Juntopas, M., and Sithirith, M., 99-121. Bangkok: Stockholm Environment Institute (SEI).
- Sithirith M (2007) *Cooperation in the Mekong River Basin: A Reflection of Cambodia's Experiences in the Development of the Mekong Region*. Chiang Mai: Regional Center for Social Science and Sustainable Development (RCSD), Faculty of Social Science, Chiang Mai University, Thailand.

- Sneddon C (2003) Conservation Initiatives and Transnationalization in the Mekong River Basin. In *Globalization and New Geography of Conservation*, ed. K. Zimmerer, 191-212. [The University of Chicago Press](#).
- Vanderveest P. and Peluso N. L (1995) Territorialization and State Power in Thailand. *Theory and Society* 24: 385-426. Netherlands: Kluwer Academic Publisher.
- Van Zalinge N, Degen P, Pongsri C, Nuov S, Jensen J, Nguyen VH, Choulamany X (2004) The Mekong River System. In *Proceedings of the Second International Symposium on the Management of Large Rivers for Fisheries* eds. Welcomme, R.L., Petr, T., 1:333–355. Bangkok, Thailand: FAO Regional Office for Asia and the Pacific.
- Wittayapak C and Vanderveest P (2010) *The Politics of Decentralization: Natural Resource Management in Asia*. Silkworm: Combined Academic, Chiang Mai: Chesham.
- Watts M & Peet R (2004) Liberating political ecology. *Liberation ecologies: Environment, development, social movements*, 2, 3-43.
- Zimmerer KS & Bassett, TJ (2003) Approaching political ecology: Society, nature, and scale in human-environment studies. In Zimmerer, K. S., Bassett, T.J. (Eds), *Political Ecology: An Integrative Approach to Geography and Environment-Development Studies*, pp. 1-25. New York: Guilford Publications,